

# Decodability of Group Homomorphisms Beyond the Johnson Bound

Irit Dinur <sup>\*</sup>    Elena Grigorescu <sup>†</sup>    Swastik Kopparty <sup>‡</sup>    Madhu Sudan <sup>§</sup>

November 19, 2007

## Abstract

Given an pair of finite groups  $G$  and  $H$ , the set of homomorphisms from  $G$  to  $H$  form an error-correcting code where codewords differ in at least  $1/2$  the coordinates. We show that for every pair of abelian groups  $G$  and  $H$ , the resulting code is (locally) list-decodable from a fraction of errors arbitrarily close to its distance. At the heart of this result is the following combinatorial result: There is a fixed polynomial  $p(\cdot)$  such that for every pair of abelian groups  $G$  and  $H$ , if the maximum fraction of agreement between two distinct homomorphisms from  $G$  to  $H$  is  $\Lambda$ , then for every  $\epsilon > 0$  and every function  $f : G \rightarrow H$ , the number of homomorphisms that have agreement  $\Lambda + \epsilon$  with  $f$  is at most  $p(1/\epsilon)$ .

We thus give a broad class of codes whose list-decoding radius exceeds the “Johnson bound”. Examples of such codes are rare in the literature, and for the ones that do exist, “combinatorial” techniques to analyze their list-decodability are limited. Our work is an attempt to add to the body of such techniques. We use the fact that abelian groups decompose into simpler ones and thus codes derived from homomorphisms over abelian groups may be viewed as certain “compositions” of simpler codes. We give techniques to lift list-decoding bounds for the component codes to bounds for the composed code. We believe these techniques may be of general interest.

---

<sup>\*</sup>Hebrew University, Jerusalem, [dinuri@cs.huji.ac.il](mailto:dinuri@cs.huji.ac.il).

<sup>†</sup>MIT CSAIL, [elena.g@mit.edu](mailto:elena.g@mit.edu).

<sup>‡</sup>MIT CSAIL, [swastik@mit.edu](mailto:swastik@mit.edu).

<sup>§</sup>MIT CSAIL, [madhu@mit.edu](mailto:madhu@mit.edu).

# 1 Introduction

It is a well-known fact that distinct homomorphisms between a fixed pair of groups  $G$  and  $H$  are *far* from each other in the Hamming norm. Thus homomorphisms between groups form “error-correcting codes”, giving rise to questions: How well can these codes be list-decoded? In this paper we give new, strong, combinatorial and algorithmic results for list-decoding of codes derived from group homomorphisms, for abelian groups.

The classical example of a code based on group homomorphisms are the Hadamard codes, which can be viewed as homomorphisms from  $\mathbb{Z}_2^n$  to  $\mathbb{Z}_2$ . The list-decoding problem for this class of groups was considered in the seminal paper of Goldreich and Levin [3], where they used this result to get a generic construction of a hardcore predicates for any one-way function. Subsequently this result has formed the basis of many results in learning, average-case complexity and cryptography.

In terms of technical generalizations, the work of Goldreich-Levin has led to the study of list-decoding for polynomial functions over finite fields (see, for instance, [4, 12]), as also the task of learning Fourier coefficients ([8, 2, 1]). Again both these directions have led to important applications in pseudorandomness, cryptography, and average-case complexity.

Our work is motivated by yet a third interpretation of the Goldreich-Levin result, as a list-decoder for homomorphisms. This interpretation was already studied in a previous work by some of the authors [5], where it was shown that the algorithmic techniques of, say [3, 4, 12], can typically be extended to the case of group homomorphisms also, provided one can get convincingly strong combinatorial bounds on the “list-decodability” of these codes. Unfortunately, bounds on the list-decodability of these codes were too weak, and indeed it seems there are few techniques to bound the list-decodability of error-correcting codes. Our paper is motivated mainly by this need to augment the analytic tools in this setting. In what follows we attempt to clarify the problem, the prior state of knowledge, and our contributions.

**Combinatorics of List-decoding Homomorphisms:** Let  $G, H$  be finite groups and let  $\Lambda = \Lambda_{G,H}$  denote the maximum relative agreement between any pair of homomorphisms between  $G$  and  $H$ . Thus the class of homomorphisms between  $G$  and  $H$  may be viewed as error-correcting codes over the alphabet  $\Sigma = H$  of length  $N = |G|$  and relative (Hamming) distance  $1 - \Lambda$ . The *combinatorial* question at the heart of this paper is the following:

Given a function  $f : G \rightarrow H$  and a real number  $\epsilon > 0$ , what is the number of homomorphisms  $\phi$  such that  $f$  and  $\phi$  agree in at least  $\Lambda_{G,H} + \epsilon$  fraction of inputs?

As we argue below this simple question was vastly misunderstood, at least till this work.

For the sake of concreteness, let  $p$  be a prime and  $n, m$  be positive integers. Now consider the class of homomorphisms between  $G = \mathbb{Z}_p^n$  and  $H = \mathbb{Z}_p^m$ . Then  $\Lambda = \Lambda_{G,H} = 1/p$ . Our combinatorial question asks: How many homomorphisms can have agreement  $1/p + \epsilon$  with any given function  $f : \mathbb{Z}_p^n \rightarrow \mathbb{Z}_p^m$ . In the case of  $m = 1$ , the works of [3, 4], assert that the number of homomorphisms is at most  $O(1/\epsilon^2)$ . Such a result can also be shown to be

essentially tight, for any choice of  $p$ ,  $n$ ,  $m$ , and  $\epsilon$ . These results essentially follow from the Johnson bound for codes over a  $p$ -ary alphabet (see, for instance, [6]).

What about the case of general  $m$ ? Here, our state of knowledge tapers off quickly. Since the codes no longer have relative distance  $1 - 1/|\Sigma|$ , the Johnson bound is no longer strong and only gives bound for the number of homomorphisms that have agreement more than  $\sqrt{1/p}$  with the given function. In other words this bound requires  $\epsilon > \sqrt{1/p} - 1/p$  and does not work for every  $\epsilon > 0$ . A more ad-hoc analysis, obtained by viewing a homomorphism from  $\mathbb{Z}_p^n$  to  $\mathbb{Z}_p^m$  as a tuple of  $m$  independent homomorphisms from  $\mathbb{Z}_p^n$  to  $\mathbb{Z}_p$ , gives a bound of  $1/\epsilon^{2m}$  on the number of homomorphisms. Till our work it was unclear if this exponential dependence on  $m$  was necessary.

A similar state of affairs is also seen when we consider, say, homomorphisms from  $\mathbb{Z}_{p^n}$  (integers modulo  $p^n$ ) to  $\mathbb{Z}_{p^m}$ . In this case, Fourier analysis can be used to get some bounds on the number of homomorphisms that have agreement  $1/p + \epsilon$  with a fixed function [5], but again this bound has exponential dependence on  $m$ .

In this paper we fix these gaps in our knowledge. We show that there is a fixed polynomial  $g$  such that for all abelian groups  $G, H$  and for every function  $f : G \rightarrow H$  and every  $\epsilon > 0$ , the number of homomorphisms with agreement  $\Lambda_{G,H} + \epsilon$  with the fixed function  $f$  is at most  $g(\epsilon)$ . (The bound is thus a fixed polynomial, independent of  $G$  and  $H$ . See Theorem 2.2.)

**Algorithmic Consequences:** Our combinatorial result turns immediately into efficient, local, algorithms for list-decoding homomorphisms. Specifically, we show that the techniques from [3, 4, 12, 5] immediately yield an algorithm that behaves as follows: It takes as input an explicit description of groups  $G$  and  $H$  (in terms of their prime decomposition) and a real number  $\epsilon > 0$ , and has oracle access to a function  $f : G \rightarrow H$ . In time polynomial in  $1/\epsilon$ ,  $\log |G|$  and  $\log |H|$  it outputs an explicit description of all homomorphisms that have agreement  $\Lambda + \epsilon$  with  $f$ .

We remark that a polynomial dependence on each is necessary to represent the output. In previous work [5] considered a fixed group  $H$  and for every such group, gave a polynomial time algorithm (i.e., its running time was polynomial in  $\log |G|$  and  $\frac{1}{\epsilon}$ ) to recover the nearby homomorphisms. Their running time grew exponentially in  $\log |H|$ .

**Techniques:** We stress that despite the widespread use of list-decoding, general techniques to establish bounds are lacking. There are relatively few codes known whose list-decodability goes beyond the Johnson bound: some examples include the Ta-Shma-Zuckerman codes [13], and the Parvaresh-Vardy-Guruswami-Rudra codes [11, 7]. The latter codes are similar in the sense that they also consider codes over an alphabet that is a vector space (rather than a field), but in their case the only proof of the list-decoding properties is algorithmic. We feel this is partly due to the lack of combinatorial techniques to analyze the decoding properties of such codes.

We start by noticing that any abelian group, in particular,  $H$  can be decomposed as  $H_1 \times \dots \times H_k$  where  $H_i = \mathbb{Z}_{p_i}^{e_i}$  for some prime  $p_i$  and positive integer  $e_i$ . Thus a homomorphism  $\phi$  from  $G$  to  $H$  can be decomposed into simpler ones using this decomposition. For instance if  $H = H_1 \times H_2$  then a homomorphism  $\phi$  from  $G$  to  $H$  is just a pair of homomorphisms  $(\phi_1, \phi_2)$  where  $\phi_i : G \rightarrow H_i$  for  $i \in \{1, 2\}$ . This decomposition allows for some weak list-decoding

bounds. To see this, fix some agreement parameter  $\rho$  and let's consider the number of homomorphisms that have agreement  $\rho$  with some function  $f : G \rightarrow H$ . Suppose we have a bound  $B_i$  on the number of homomorphisms from  $G$  to  $H_i$  that have agreement  $\rho$  with any single function. Then, viewing  $f$  as a pair of functions  $(f_1, f_2)$ ,  $f_i : G \rightarrow H_i$ , and noticing that for any homomorphism  $\phi = (\phi_1, \phi_2) : G \rightarrow H$  that has agreement  $\rho$  with  $f$  it must be the case that  $\phi_i$  and  $f_i$  also have agreement at least  $\rho$ , we conclude that the number of homomorphisms that have agreement  $\rho$  with  $f$  is at most  $B_1 \cdot B_2$ . Unfortunately, this bound grows exponentially with  $\log |H|$  and this is too weak to get our claimed theorem.

Our improvements come by noticing that agreements between homomorphisms are quite restricted. For instance, in the case where  $H_1 = H_2 = \mathbb{Z}_p$  and  $H = H_1 \times H_2$ , homomorphisms  $\phi, \psi : G \rightarrow H_1$  either agree on a set of density  $\frac{1}{p}$  or a set of density at most  $\frac{1}{p^2}$ . To leverage this insight in our setting, say we have a function  $f = (f_1, f_2)$  and we wish to bound homomorphisms with agreement  $\rho$  with  $f$ . Suppose  $\phi_1 : G \rightarrow H_1$  has agreement at least  $\rho$  with  $f_1$  and suppose the set of agreements is the set  $S \subseteq G$ . We now consider all homomorphisms  $\psi_1, \dots, \psi_\ell$  that agree with  $f_2$  on subsets of  $S$  of size at least  $\rho \cdot |G|$ . We consider agreements of “triples” of homomorphisms  $\psi_i, \psi_j$  and  $\psi_k$  and note that if every such triple has a large mutual agreement in the set  $S$ , then they satisfy a *sunflower* like property, forcing  $S$  to be very large. If on the other hand every triple has a small agreement then we manage to show that  $S$  is still large, by using an inclusion-exclusion count. We then use a standard argument to show that either the given collection of homomorphisms always contain a large subcollection in which every triple has a large intersection, or a large subcollection in which every triple has a small intersection. This gives us a nontrivial lower bound on  $S$  in all cases and then we perform some very careful accounting to show that this leads to an (absolute) polynomial upper bound on the list size.

While all these arguments may seem very specific to this special case of homomorphism from  $G$  to  $\mathbb{Z}_p \times \mathbb{Z}_p$ , we show that this is not the case. Indeed we abstract a nice property of families of sets (in our case the points of agreement between  $f_2$  and  $\psi_1, \dots, \psi_k$  within a specified set  $S$ ) that allows us to give non-trivial bounds on the size of their union. We then apply this same bound in three different cases to lift list-decoding bounds for somewhat simple groups to list decoding bounds for more complex groups. We start with homomorphisms to  $\mathbb{Z}_p$  and lift this to homomorphisms to  $\mathbb{Z}_p^r$ , then to  $\mathbb{Z}_{p^r}$  and then to arbitrary groups  $\prod_i \mathbb{Z}_{p_i^{e_i}}$ . This leads us to the final theorem claimed above.

The combinatorial results turn into algorithmic results in a straightforward manner based on previous works [3, 4, 12, 5]. If anything, our algorithms become even simpler because our combinatorial bounds are stronger. See Appendix C.

**Related prior works:** The task of list-decoding homomorphisms is very closely related to the task of determining significant Fourier coefficients in abelian groups. Previous works [8, 10, 2, 1] have consider the latter task in different contexts and in particular Akavia et al. [1] give comprehensive results on “list-decoding” for the significant Fourier coefficients. Here we point out why their results do not subsume ours. Even though significant Fourier coefficients correspond exactly to list decoding when  $H = \mathbb{Z}_2$  or  $H = \mathbb{Z}_3$ , the situation changes significantly over other groups. The complex inner product, the norm underlying Fourier coefficients, is very different from the Hamming distance and indeed it is possible to

find functions with few large Fourier coefficients that are not close to any homomorphism in the Hamming norm. Conversely, it is also possible to find functions that are very close to homomorphisms in the Hamming norm that have no significant Fourier coefficients. Thus in these cases, list-decoding is a very different problem than that of finding Fourier coefficients, and combinatorial bounds such as the ones we provide were not known.

**Organization of this paper:** In Section 2 we introduce some necessary definitions and state our main results. In Section 3, we prove our main combinatorial bounds, modulo a theorem about certain set systems with restricted intersections. Finally in Section 4, we prove this theorem. Some proofs are deferred to the Appendix.

## 2 Definitions and Main Results

We start with some basic terminology about homomorphisms and introduce the combinatorial and algorithmic list decoding problems and state our results.

For abelian groups  $G, H$  let  $\text{Hom}(G, H) = \{h : G \rightarrow H \mid h(x) + h(y) = h(x + y), \forall x, y \in G\}$  be the set of homomorphisms from  $G$  to  $H$ . Also let  $\text{aHom}(G, H) = \{h + a : G \rightarrow H \mid h \in \text{Hom}(G, H), a \in H\}$  be the set of affine homomorphisms from  $G$  to  $H$ . Notice that  $\text{Hom}(G, H) \subset \text{aHom}(G, H)$ .

For two functions  $f, g : G \rightarrow H$ , define  $\text{agree}(f, g) = \Pr_{x \in G}[f(x) = g(x)]$ , and  $\Lambda_{G, H} = \max_{f, g \in \text{Hom}(G, H), f \neq g} \{\text{agree}(f, g)\}$ . In the case when  $\text{Hom}(G, H)$  contains only the zero homomorphism we define  $\Lambda_{G, H} = 0$ .

**Definition 2.1 (Combinatorial List Decodability)** *The code  $\text{aHom}(G, H)$  is  $(\delta, l)$ -list decodable if for every function  $f : G \rightarrow H$ , there exist at most  $l$  homomorphisms  $h \in \text{aHom}(G, H)$  such that  $\text{agree}(f, h) \geq \delta$ .*

The principal question that we address is: For which function  $l(\delta)$  can we conclude that  $\text{aHom}(G, H)$  is  $(\delta, l(\delta))$ -list-decodable. Our theorem below shows that this is true for  $l(\delta) = \text{poly}((\delta - \Lambda_{G, H})^{-1})$  for some polynomial independent of  $G$  and  $H$ .

**Theorem 2.2** *There is a universal constant  $C$  such that the following holds: Let  $G, H$  be abelian groups. Then for every  $\epsilon > 0$ ,  $\text{aHom}(G, H)$  is  $(\Lambda_{G, H} + \epsilon, 1/\epsilon^C)$ -list-decodable.*

We augment this combinatorial result with an algorithmic one that finds the nearby homomorphisms efficiently. We first define this algorithmic problem below and then state our algorithmic theorem.

Our algorithmic goal is to produce a list of all homomorphisms with agreement  $\Lambda_{G, H} + \epsilon$  with some function  $f : G \rightarrow H$  efficiently, i.e., in time  $\text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon})$ . In order to do so, the algorithm requires oracle access to the function  $f$  (or else reading the function will take time  $\text{poly}(|G|)$ ). Our solution requires a further assumption that the groups  $G$  and  $H$  are given explicitly, in a sense described next.

Recall that the structure theorem for finite abelian groups [9] states that every finite abelian group can be decomposed into cyclic groups of the form  $\mathbb{Z}_{p_1^{r_1}} \times \dots \times \mathbb{Z}_{p_k^{r_k}}$ , where all  $p_i$ 's are prime. We will assume that our input groups  $G$  and  $H$  are presented in this cyclic decomposition. Our algorithm returns all such homomorphisms explicitly by specifying its values on a set of generators of  $G$ .

**Definition 2.3 (Algorithmic Local List Decoding)** *A probabilistic oracle algorithm  $\mathcal{A}$  for list decoding homomorphisms takes as input two groups  $G$  and  $H$  represented explicitly, and a parameter  $\delta > 0$  (where the explicit representation of  $G = \mathbb{Z}_{p_1^{r_1}} \times \dots \times \mathbb{Z}_{p_k^{r_k}}$  is the sequence of pairs  $\langle (p_1, r_1), \dots, (p_k, r_k) \rangle$ , and similarly for  $H$ ) and has oracle access to a function  $f : G \rightarrow H$ . We say that  $\mathcal{A}$  is a  $(\delta, T)$ -local list decoder for groups  $G, H$ , if for every function  $f : G \rightarrow H$ ,  $\mathcal{A}^f$  runs in time  $T$  and outputs a list  $\{\phi_1, \phi_2, \dots, \phi_L\} \subset \text{aHom}(G, H)$  s.t. with probability at least  $3/4$ , it is the case that for every affine homomorphism  $h \in \text{aHom}(G, H)$  such that  $\text{agree}(f, h) \geq \delta$ ,  $h = \phi_j$  for some  $j \in [L]$ .*

**Theorem 2.4** *There exists an algorithm  $\mathcal{A}$ , such that for every pair of abelian groups  $G, H$ , and every  $\epsilon > 0$ ,  $\mathcal{A}$  is a  $(\Lambda_{G,H} + \epsilon, \text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon}))$ -local list decoder for  $\text{aHom}(G, H)$ .*

### 3 Analysis of Combinatorial List Decodability

In this section we prove Theorem 2.2 above, which upper bounds the number of homomorphisms that have agreement  $\Lambda_{G,H} + \epsilon$ , for some  $\epsilon > 0$ , with a given function  $f : G \rightarrow H$ .

We begin by considering two special cases. In both special cases  $G$  is a somewhat special group, which we will elaborate on later. In the first case  $H = \mathbb{Z}_p^r$ , while in the second case  $H = \mathbb{Z}_{p^r}$ . Finally we analyze the case of general  $G$  and  $H$ , by reducing it to the special cases.

All three settings (the two special cases and the general case) are analyzed by a common technique by considering set systems with very special intersection properties and bounding their cardinality (in a very specific way). In our analysis these sets will represent sets of agreement between a function at hand and some nearby homomorphisms. We define this special intersection property and state our crucial theorem about these families next.

#### 3.1 Special Intersecting Families

Below we consider a collection  $S_1, \dots, S_\ell$  of subsets of a universe  $X$ . For a set  $T \subset X$ , its density is denoted  $\mu(T) \stackrel{\text{def}}{=} |T|/|X|$ . For a set of indices  $I \subseteq [\ell]$  let  $S_I = \cap_{i \in I} S_i$  (if  $I = \emptyset$  then  $S_I = X$ ).

**Definition 3.1** *For  $0 < \tau \leq \rho \leq 1$  and  $c < \infty$  we say that a family of sets  $S_1, \dots, S_\ell \subseteq X$  is a  $(\rho, \tau, c)$ -special intersecting family if the following four properties hold:*

1. For every index  $i \in [\ell]$ ,  $\mu(S_i) \geq \rho$ .
2. For every pair of distinct indices  $i, j \in [\ell]$ ,  $\mu(S_i \cap S_j) \leq \rho$ .
3.  $\sum_{i=1}^{\ell} \alpha_i^c \leq 1$ , where  $\alpha_i = \mu(S_i) - \rho$ .
4. For subsets (of indices)  $I, J \subseteq [\ell]$ , with  $J \subseteq I$  and  $|J| \geq 2$ , if  $\mu(S_I) > \tau$  the  $S_I = S_J$ .

The first two properties above are typical conditions seen in intersecting systems, but the “restrictions” are not particularly restrictive. The third property is a typical condition that is derived when the sets  $S_i$  represent agreement sets of some fixed word with codewords of an orthogonal error-correcting code. (This condition is usually proved using Parseval’s identity, or using the Johnson bound, for  $c = 2$ .) The final condition is effectively “Helly”-like. It says, roughly, that the only way a large collection of sets  $\{S_i\}_{i \in I}$  can have a non-trivially large intersection is if every pair of intersecting sets with indices from  $I$  have the same intersection.

Our main theorem about set systems with special intersections roughly shows that the density of the union of the  $S_i$ ’s is relatively large compared to the density of the sets themselves.

**Theorem 3.2** *For every  $c < \infty$ , there exists a  $C < \infty$  such that the following holds: Let  $\rho > 0$  and let  $S_1, \dots, S_\ell \subseteq X$  be a  $(\rho, \rho^2, c)$ -special intersecting family. Let  $\alpha_i = \mu(S_i) - \rho$  and let  $\alpha = \mu(\cup_{i \in [\ell]} S_i) - \rho$ . Then  $\alpha^C \geq \sum_{i \in [\ell]} \alpha_i^C$ .*

We defer the proof of Theorem 3.2 to the next section. We first show how it can be used to prove upper bounds on the list size when decoding affine homomorphisms.

### 3.2 Upper bound for $H = \mathbb{Z}_p^r$

In this case and the next one, we will be dealing with a special class of groups  $G$  that are called  $p$ -groups. Specifically,  $G$  is a  $p$ -group if  $|G|$  is a power of a prime  $p$ . What we use below is that every subgroup of a  $p$ -group has cardinality at most  $|G|/p$ .

**Theorem 3.3** *There exists a constant  $C$  such that the following is true for every prime  $p$  and positive integer  $r$ . Let  $H = \mathbb{Z}_p^r$ . Given any function  $f : G \rightarrow H$  and agreement parameter  $\epsilon > 0$ , the number of affine homomorphisms from  $G$  to  $H$  that have agreement at least  $1/p + \epsilon$  with  $f$  is at most  $(1/\epsilon)^C$ .*

**Proof** Fix  $f : G \rightarrow H$ . Note that  $f$  can be viewed as an  $r$ -tuple of functions  $f = \langle f_1, \dots, f_r \rangle$  with  $f_i : G \rightarrow \mathbb{Z}_p$ . For  $k \in [r]$  define the  $k$ th projection of  $f$  to be the function  $f^{[k]} : G \rightarrow \mathbb{Z}_p^k$  given by  $f^{[k]}(x) = \langle f_1(x), \dots, f_k(x) \rangle$ . We extend this notation to include  $k = 0$ , by allowing a unique function from  $G \rightarrow \mathbb{Z}_p^0$  (which is considered to be an affine homomorphism).

For  $k_1 \leq k_2$ , we say that a function  $f_2 : G \rightarrow \mathbb{Z}_p^{k_2}$  extends  $f_1 : G \rightarrow \mathbb{Z}_p^{k_1}$  if  $f_1 = f_2^{[k_1]}$ .

In what follows, we fix  $k$ ,  $0 \leq k < r$  and consider the function  $f^{[k]}$ . We also fix an affine homomorphism  $\phi$  from  $G$  to  $\mathbb{Z}_p^k$  that has agreement  $\frac{1}{p} + \alpha$  with  $f^{[k]}$ , for some  $\alpha \geq \epsilon$ . We prove that the number of affine homomorphisms that extend  $\phi$  and have agreement  $1/p + \epsilon$  with  $f$  is small.

**Claim 3.4** *For every  $0 \leq k \leq r$ , and for every affine homomorphism  $\phi$  that has agreement  $1/p + \alpha$  with  $f^{[k]}$ , the number of affine homomorphisms from  $G$  to  $\mathbb{Z}_p^r$  that have agreement  $1/p + \epsilon$  with  $f$  and extend  $\phi$ , is at most  $(\alpha/\epsilon)^C$ , where  $C$  is the constant from Theorem 3.2, for  $c = 2$ .*

**Proof** We prove the claim by induction on  $r - k$ . When  $r - k = 0$ , the claim is trivially true. So assume the claim is true for  $r - (k + 1)$  and we will now prove it for the case of  $r - k$ .

Let  $S = \{x \in G \mid f^{[k]}(x) = \phi(x)\}$ . Consider  $f^{[k+1]}$  and let  $\phi_1, \dots, \phi_\ell$  be all the affine homomorphisms from  $G$  to  $\mathbb{Z}_p^{k+1}$  that extend  $\phi$  with agreement at least  $\frac{1}{p} + \epsilon$  with  $f^{[k+1]}$ . Let  $S_i = \{x \mid f^{[k+1]}(x) = \phi_i(x)\}$  and let  $\mu(S_i) = \frac{1}{p} + \alpha_i$ .

We prove below that the sets  $S_1, \dots, S_\ell \subseteq G$  form a  $((1/p), (1/p^2), 2)$ -special intersecting family. This will allow us to apply Theorem 3.2 to these sets and this in turn will allow us to derive the inductive claim easily.

**Verifying Intersection Properties:** The first property requires that  $\mu(S_i) \geq \frac{1}{p}$  which is immediate from the definition of the  $S_i$ 's. The second property required that  $\mu(S_i \cap S_j) \leq \frac{1}{p}$ . This again is true, since  $S_i \cap S_j \subseteq \{x \mid \phi_i(x) = \phi_j(x)\}$  and the latter has density at most  $1/p$ .

The third property follows from the “ $p$ -ary Johnson bound” applied to the function  $f_{k+1}^{[k+1]} : G \rightarrow \mathbb{Z}_p$ . Let  $(\phi_i)_j$  denote the  $j$ th coordinate of the function  $\phi_i$ . Then note that the functions  $(\phi_i)_{k+1}$  are distinct affine homomorphisms, with  $(\phi_i)_{k+1}$  having agreement at least  $1/p + \alpha_i$  with  $f_{k+1}$ . It follows from Lemma D.1 (in Section D) that  $\sum_{i=1}^\ell \alpha_i^2 \leq 1$ .

Finally, we come to the crucial “special-intersection” property. We need to show that if  $\mu(S_I) > 1/p^2$ , and  $J \subseteq I$  contains at least two distinct elements, then  $S_I = S_J$ . For any set  $K \subseteq [\ell]$  define  $T_K = \{x \mid \phi_i(x) = \phi_j(x), \forall i, j \in K\}$ . Note that  $T_K = G$  if and only if  $|K| \leq 1$ . Since  $T_K$  is a coset of a subgroup of  $G$ , we have that  $\mu(T_K) = 1/p^i$  for non-negative integer  $i$ . But  $S_K \subseteq T_K$  for every  $K$ , and since  $\mu(S_I), \mu(S_J) > 1/p^2$ , it must be that  $\mu(T_I), \mu(T_J) = 1/p$ . Since  $T_I \subseteq T_J$ , we get that  $T_I = T_J$ . Finally, fix some  $j \in J$  and let  $Y = \{x \mid \phi_j(x) = f^{[k+1]}(x)\}$ . Since  $\phi_i(x) = \phi_j(x)$  for every  $x \in Y \cap T_I$  and  $i \in I$ , we have  $S_I = Y \cap T_I = Y \cap T_J = S_J$  as required for the fourth property.

**Proof of Claim:** We are now ready to prove the claim. Since the sets  $S_1, \dots, S_\ell$  form a  $(1/p, 1/p^2, 2)$ -special intersecting family, we can apply Theorem 3.2 to conclude that  $(\alpha')^C \geq \sum_{i=1}^\ell \alpha_i^C$  where  $\alpha' = \mu(\cup_{i \in \ell} S_i) - 1/p$ . Since for every  $i$ ,  $S_i \subseteq S$ , we have  $\cup_{i \in \ell} S_i \subseteq S$  and so  $\alpha \geq \alpha'$ . We may thus conclude that  $\alpha^C \geq \sum_{i=1}^\ell \alpha_i^C$ . But this immediately leads to the claim as follows. By induction we have that the number of extensions of  $\phi_i$  in  $\text{aHom}(G, H)$  that have  $\frac{1}{p} + \epsilon$  agreement with  $f$  is at most  $(\alpha_i/\epsilon)^C$ . Since every affine homomorphism in



$\text{aHom}(G, H)$  that extends  $\phi$  and has  $\frac{1}{p} + \epsilon$  agreement with  $f$  must also extend one of the  $\phi_i$ , we have that the total number of such extensions (of  $\phi$ ) is at most  $\sum_{i \in [\ell]} (\alpha_i/\epsilon)^C \leq (\alpha/\epsilon)^C$  as desired. ■

The theorem statement follows immediately, by using  $k = 0$ . ■

### 3.3 Upper bound for $H = \mathbb{Z}_{p^r}$

We now consider a variant of Theorem 3.3 with  $H = \mathbb{Z}_{p^r}$ , the cyclic group of integers modulo a prime power (as opposed to the vector space  $H = \mathbb{Z}_p^r$ ).

We state the theorem below and prove it in Appendix A.1. We remark the proof is essentially the same as that of Theorem 3.3.

**Theorem 3.5** *There exists a constant  $C$  such that the following is true for every prime  $p$  and positive integer  $r$ . Let  $H = \mathbb{Z}_{p^r}$ . Given any function  $f : G \rightarrow H$  and agreement parameter  $\epsilon > 0$ , the number of affine homomorphisms from  $G$  to  $H$  that have agreement at least  $1/p + \epsilon$  with  $f$  is at most  $(1/\epsilon)^C$ . Furthermore, if  $\phi_1, \dots, \phi_\ell : G \rightarrow H$  are affine homomorphisms with  $\text{agree}(f, \phi_i) = 1/p + \alpha_i \geq 1/p + \epsilon$ , then  $\sum_{i=1}^{\ell} \alpha_i^C \leq 1$ .*

### 3.4 General abelian groups

Finally we deal with the case of general  $H$ . We first deal with the case of slightly restricted  $G$  which we describe below.

Let  $H = \mathbb{Z}_{p_1^{r_1}} \times \dots \times \mathbb{Z}_{p_m^{r_m}}$ . We assume below that  $p_i$  divides the order of  $G$  for every  $i$ , and that  $p = p_1 \leq p_2 \leq \dots \leq p_m$ . Thus  $\Lambda_{G,H} = 1/p$ . The following lemma deals with this case. Later we remove the restriction on  $G$ .

**Lemma 3.6** *There exists a constant  $C_1$  such that the following holds. Let  $p = p_1 \leq \dots \leq p_m$  be primes and  $r_1, \dots, r_m$  be positive integers. Let  $H = \prod_{i=1}^m \mathbb{Z}_{p_i^{r_i}}$  and let  $G$  be a finite abelian group such that  $p_i$  divides the order of  $G$  for every  $i \in [m]$ . Given any function  $f : G \rightarrow H$  and agreement parameter  $\epsilon > 0$ , the number of affine homomorphisms from  $G$  to  $H$  that have agreement at least  $1/p + \epsilon$  with  $f$  is at most  $(1/\epsilon)^{C_1}$ .*

**Proof** The proof is obtained by essentially notational changes to the proof of Theorem 3.3. For  $i \in [m]$  let  $H_i = \mathbb{Z}_{p_i^{r_i}}$ . Fix  $f : G \rightarrow H$ . We view  $f$  as an  $m$ -tuple of functions  $f = \langle f_1, \dots, f_m \rangle$  with  $f_i : G \rightarrow H_i$ .

For  $k \in [m]$ , let  $H_{[k]} = \prod_{i=1}^k H_i$ , and let  $f^{[k]} : G \rightarrow H_{[k]}$  be given by  $f^{[k]}(x) = \langle f_1(x), \dots, f_k(x) \rangle$ . We extend this notation to include  $k = 0$ , by allowing a unique function from  $G \rightarrow \{1\}$  (which is considered to be a homomorphism).

For  $k_1 \leq k_2$ , we say that a function  $f_2 : G \rightarrow H_{[k_2]}$  extends  $f_1 : G \rightarrow H_{[k_1]}$  if  $f_1 = f_2^{[k_1]}$ .

In what follows, we fix  $k$ ,  $0 \leq k < m$  and consider the function  $f^{[k]}$ . We also fix a homomorphism  $\phi$  from  $G$  to  $H_{[k]}$  that has agreement  $\frac{1}{p} + \alpha$  with  $f^{[k]}$ , for some  $\alpha \geq \epsilon$ . We prove that the number of homomorphisms that extend  $\phi$  and have agreement  $1/p + \epsilon$  with  $f$  is small.

**Claim 3.7** *For every  $0 \leq k \leq m$ , and for every homomorphism  $\phi$  that has agreement  $1/p + \alpha$  with  $f^{[k]}$ , the number of homomorphisms from  $G$  to  $H$  that have agreement  $1/p + \epsilon$  with  $f$  and extend  $\phi$ , is at most  $(\alpha/\epsilon)^{C_1}$ , where  $C_1$  is the constant from Theorem 3.2, for  $c = C_2$ , and  $C_2$  is the constant from Theorem 3.5.*

**Proof** We prove the claim by induction on  $r - k$ . When  $r - k = 0$ , the claim is trivially true. So assume the claim is true for  $r - (k + 1)$  and we will now prove it for the case of  $r - k$ .

Let  $S = \{x \in G \mid f^{[k]}(x) = \phi(x)\}$ . Consider  $f^{[k+1]}$  and let  $\phi_1, \dots, \phi_\ell$  be all the homomorphisms from  $G$  to  $H_{[k+1]}$  that extend  $\phi$  with agreement at least  $\frac{1}{p} + \epsilon$  with  $f^{[k+1]}$ . Let  $S_i = \{x \mid f^{[k+1]}(x) = \phi_i(x)\}$  and let  $\mu(S_i) = \frac{1}{p} + \alpha_i$ .

We prove below that the sets  $S_1, \dots, S_\ell \subseteq G$  form a  $((1/(p_{k+1}), (1/p_{k+1}^2), C_2)$ -special intersecting family. This will allow us to apply Theorem 3.2 to these sets and this in turn will allow us to derive the inductive claim easily. We use  $q$  as shorthand for  $p_{k+1}$ .

**Verifying Intersection Properties:** The first property requires that  $\mu(S_i) \geq \frac{1}{q}$  which is immediate from the definition of the  $S_i$ 's. The second property required that  $\mu(S_i \cap S_j) \leq \frac{1}{q}$ . This again is true, since  $S_i \cap S_j \subseteq \{x \mid \phi_i(x) = \phi_j(x)\}$  and the latter has density at most  $1/q$ .

The third property follows from Theorem 3.5, applied to the function  $f_{k+1}^{[k+1]} : G \rightarrow H_{k+1}$ . Let  $(\phi_i)_j$  denote the  $j$ th coordinate of the function  $\phi_i$ . Then note that the functions  $(\phi_i)_{k+1}$  are distinct homomorphisms from  $G$  to  $H_{k+1}$ , with  $(\phi_i)_{k+1}$  having agreement at least  $1/q + \alpha_i$  with  $f_{k+1}$ . It follows from Theorem 3.5, that  $\sum_{i=1}^{\ell} \alpha_i^{C_2} \leq 1$ .

Finally, we come to the fourth property. We need to show that if  $\mu(S_I) > 1/q^2$ , and  $J \subseteq I$  contains at least two distinct elements, then  $S_I = S_J$ . For any set  $K \subseteq [\ell]$  define  $T_K = \{x \mid \phi_i(x) = \phi_j(x), \forall i, j \in K\}$ . Note that  $T_K = G$  if and only if  $|K| \leq 1$ . Also note that  $\phi_i(x) = \phi_j(x)$  if and only if  $(\phi_i)_{k+1}(x) = (\phi_j)_{k+1}(x)$  (since both extend  $\phi$ ). Thus  $T_K$  is the kernel of a homomorphism from  $G$  to  $H_{k+1}$ , we have that  $1/\mu(T_K)$  must divide  $|H_{k+1}|$  and so  $\mu(T_K) = 1/q^i$  for non-negative integer  $i$ . But  $S_K \subseteq T_K$  for every  $K$ , and since  $\mu(S_I), \mu(S_J) > 1/q^2$ , it must be that  $\mu(T_I), \mu(T_J) = 1/q$ . Since  $T_I \subseteq T_J$ , we get that  $T_I = T_J$ . Finally, fix some  $j \in J$  and let  $Y = \{x \mid \phi_j(x) = f^{[k+1]}(x)\}$ . Since  $\phi_i(x) = \phi_j(x)$  for every  $x \in Y \cap T_I$  and  $i \in I$ , we have  $S_I = Y \cap T_I = Y \cap T_J = S_J$  as required for the fourth property.

**Proof of Claim:** We are now ready to prove the claim. Since the sets  $S_1, \dots, S_\ell$  form a  $(1/q, 1/q^2, C_2)$ -special intersecting family, we can apply Theorem 3.2 to conclude that there exists a constant  $C_1$  such that  $(\alpha')^{C_1} \geq \sum_{i=1}^{\ell} \alpha_i^{C_1}$  where  $\alpha' = \mu(\cup_{i \in \ell} S_i) - 1/p$ . Since for every  $i$ ,  $S_i \subseteq S$ , we have  $\cup_{i \in \ell} S_i \subseteq S$  and so  $\alpha \geq \alpha'$ . We may thus conclude that  $\alpha^{C_1} \geq \sum_{i=1}^{\ell} \alpha_i^{C_1}$ . This immediately yields the claim. ■

The lemma statement follows immediately, by using  $k = 0$ . ■

Using Lemma 3.6 it is easy to complete the proof of our main theorem, Theorem 2.2. We defer the proof to Appendix A.2.

## 4 Analyzing Special Intersecting Families

In this section we give an overview of the proof of Theorem 3.2, the complete proof being deferred to Appendix B.

Fix a  $(\rho, \rho^2, c)$ -special intersecting family  $S_1, \dots, S_\ell \subseteq X$  and let  $\alpha_i = \mu(S_i) - \rho$ . Let  $\alpha = \mu(\cup_{i \in [\ell]} S_i) - \rho$ . Recall that our goal is to prove that  $\alpha^C \geq \sum_{i=1}^{\ell} \alpha_i^C$  for a sufficiently large  $C$  (that depends only on  $c$ ).

The proof has two main stages. In the first stage, we show a weak lower bound on  $\alpha$  in terms of the  $\alpha_i$ . This is done in Lemma B.5, where we show that there exist constants  $c_1 > 0$ ,  $c_2 < 1$  and  $c_3$  depending only on  $c$  such that, for every  $K \subseteq [\ell]$ ,

$$\alpha^{c_1} \geq \frac{1}{c_3 \cdot |K|^{c_2}} \cdot \sum_{k \in K} \alpha_k^{c_1}.$$

The constraint  $c_2 < 1$  makes this Lemma extremely useful. Then, in the second stage, we show that the weak bound obtained above, along with the observation  $\alpha \geq \alpha_i + \alpha_j$  for all  $i \neq j$ , together imply the existence of  $C$  depending only on  $c$  with  $\alpha_i^C \geq \sum_{i=1}^{\ell} \alpha_i^C$ . This is proved in Lemma B.6.

We prove the weak lower bound on  $\alpha$  (Lemma B.5) by exploiting the rich structure in the sets  $S_I$ . We say  $I \subseteq [\ell]$  is a *collinear set* if either  $|I| \leq 2$  or  $\mu(S_I) > \rho^2$ . Property (4) of special intersecting families gives us some interesting properties of collinear sets, that make them resemble *collinear sets of points on the plane*.

1. If  $I$  is a collinear set, and  $J \subseteq I$ , then  $J$  is a collinear set.
2. If  $I, J$  are collinear sets with  $|I \cap J| \geq 2$ , then  $I \cup J$  is a collinear set.
3. For  $i \neq j \in [\ell]$ , then the set  $L_{i,j}$  of all points  $k \in [\ell]$  such that  $\{i, j, k\}$  is a collinear set, is itself a collinear set.

Pushing the analogy further, we say  $J \subseteq [\ell]$  is *in general position* if there is no collinear set  $K \subseteq J$  with  $|K| = 3$ .

The argument for proving the weak lower bound on  $\alpha$  proceeds by separately analyzing collinear sets and sets in general position. For any collinear set  $I$ , by finding a sunflower in the set system, we obtain a lower bound on  $\alpha$  of the form  $\sum_{i \in I} \alpha_i$  (see Lemma B.3). For a set  $J$  in general position, by an inclusion exclusion argument, we obtain a lower bound on  $\alpha$  of the form  $\Omega(\sum_{j \in J} \alpha_j)$  (see Lemma B.3). The weak lower bound on  $\alpha$  (Lemma B.5) is now obtained by showing a dichotomy: either there is a collinear set  $I$  with  $\sum_{i \in I} \alpha_i$  large, or there is a set  $J$  in general position with  $\sum_{j \in J} \alpha_j$  large. The details are given in Appendix B.

## Acknowledgments

Thanks to Amir Shpilka for many valuable discussions.

## References

- [1] A. Akavia, S. Goldwasser, and S. Safra. Proving hard-core predicates using list decoding. In *FOCS: IEEE Symposium on Foundations of Computer Science (FOCS)*, 2003.
- [2] A. Gilbert, S. Guha, P. Indyk, S. Muthukrishnan, and M. Strauss. Near-optimal sparse fourier representations via sampling. In *STOC: ACM Symposium on Theory of Computing (STOC)*, 2002.
- [3] Oded Goldreich and Leonid Levin. A hard-core predicate for all one-way functions. *Proceedings of the 21st Annual ACM Symposium on Theory of Computing*, pages 25–32, May 1989.
- [4] Oded Goldreich, Ronitt Rubinfeld, and Madhu Sudan. Learning polynomials with queries: The highly noisy case. *SIAM Journal on Discrete Mathematics*, 13(4):535–570, November 2000.
- [5] Elena Grigorescu, Swastik Kopparty, and Madhu Sudan. Local decoding and testing for homomorphisms. In Josep Díaz, Klaus Jansen, José D. P. Rolim, and Uri Zwick, editors, *APPROX-RANDOM*, volume 4110 of *Lecture Notes in Computer Science*, pages 375–385. Springer, 2006.
- [6] V. Guruswami and M. Sudan. Extensions to the johnson bound, 2001.
- [7] Venkatesan Guruswami and Atri Rudra. Explicit capacity-achieving list-decodable codes. *Electronic Colloquium on Computational Complexity (ECCC)*, (133), 2005.
- [8] E. Kushilevitz and Y. Mansour. Learning decision trees using the fourier spectrum. *SICOMP: SIAM Journal on Computing*, 22, 1993.
- [9] Serge Lang. *Algebra*. Addison-Wesley, 1965.
- [10] Yishay Mansour. Randomized interpolation and approximation of sparse polynomials. *SIAM J. Comput.*, 24(2):357–368, 1995.
- [11] Farzad Parvaresh and Alexander Vardy. Correcting errors beyond the guruswami-sudan radius in polynomial time. In *FOCS*, pages 285–294. IEEE Computer Society, 2005.
- [12] Madhu Sudan, Luca Trevisan, and Salil Vadhan. Pseudorandom generators without the XOR lemma. *Proceedings of the 31st Annual ACM Symposium on Theory of Computing*, pages 537–546, 1999.
- [13] A. Ta-Shma and D. Zuckerman. Extractor codes. *IEEE Transactions on Information Theory*, 50, 2004.

# A Proofs of Combinatorial List Decodability

## A.1 Upper bound for $H = \mathbb{Z}_{p^r}$

Here we prove Theorem 3.5. Recall that  $G$  is a  $p$ -group,  $H = \mathbb{Z}_{p^r}$ , the cyclic group of integers modulo a prime power (as opposed to the vector space  $H = \mathbb{Z}_p^r$  as in Theorem 3.3).

### Proof

For any function  $g : G \rightarrow \mathbb{Z}_{p^r}$  and  $i \in [\ell]$ , let  $g^{(i)} : G \rightarrow \mathbb{Z}_{p^i}$  be given by  $g^{(i)}(x) = g(x) \bmod (p^i)$ . Also, let  $g^{(-i)} : G \rightarrow \mathbb{Z}_{p^{r-i}}$  be such that <sup>1</sup> for all  $x \in G$ ,  $g(x) = p^i \cdot g^{(-i)}(x) + g^{(i)}(x)$ .  $g^{(-i)}$  is clearly well defined. Note that if  $g$  is an affine homomorphism, then  $g^{(i)}$  is also an affine homomorphism, while  $g^{(-i)}$ , in general, need not be one. For  $j \leq i$ , we say that  $g : G \rightarrow \mathbb{Z}_{p^i}$  extends  $h : G \rightarrow \mathbb{Z}_{p^j}$  if  $g^{(j)} = h$ .

Now fix  $f : G \rightarrow H$ . Fix  $k \in \{0, \dots, r\}$ . Let  $\phi : G \rightarrow \mathbb{Z}_{p^k}$  be an affine homomorphism with agreement  $1/p + \alpha \geq 1/p + \epsilon$  with  $f^{(k)}$ . (As usual, we assume there is a single function from  $G$  to  $\mathbb{Z}_{p^0}$  and that it is a homomorphism.) We claim that the number of affine homomorphisms extending  $\phi$  is small.

**Claim A.1** *For every  $0 \leq k \leq r$ , and for every affine homomorphism  $\phi$  that has agreement  $1/p + \alpha$  with  $f^{(k)}$ , the number of affine homomorphisms from  $G$  to  $\mathbb{Z}_{p^r}$  that have agreement  $1/p + \epsilon$  with  $f$  and extend  $\phi$ , is at most  $(\alpha/\epsilon)^C$ , where  $C$  is the constant from Theorem 3.2, for  $c = 2$ .*

**Proof** The proof is identical to that of Claim 3.4 with mainly notational changes switching  $f^{[\cdot]}$  to  $f^{(\cdot)}$ . The only noticeable change is in the argument for the third property of “special intersections”.

We prove the claim by induction on  $r - k$ . When  $r - k = 0$ , the claim is trivially true. So assume the claim is true for  $r - (k + 1)$  and we will now prove it for the case of  $r - k$ .

Let  $S = \{x \in G \mid f^{(k)}(x) = \phi(x)\}$ . Consider  $f^{(k+1)}$  and let  $\phi_1, \dots, \phi_\ell$  be all the affine homomorphisms from  $G$  to  $\mathbb{Z}_{p^{k+1}}$  that extend  $\phi$  with agreement at least  $\frac{1}{p} + \epsilon$  with  $f^{(k+1)}$ . Let  $S_i = \{x \mid f^{(k+1)}(x) = \phi_i(x)\}$  and let  $\mu(S_i) = \frac{1}{p} + \alpha_i$ .

We prove below that the sets  $S_1, \dots, S_\ell \subseteq G$  form a  $((1/p), (1/p^2), 2)$ -special intersecting family. This will allow us to apply Theorem 3.2 to these sets and this in turn will allow us to derive the inductive claim easily.

**Verifying Intersection Properties:** The first property requires that  $\mu(S_i) \geq \frac{1}{p}$  which is immediate from the definition of the  $S_i$ 's. The second property required that  $\mu(S_i \cap S_j) \leq \frac{1}{p}$ . This again is true, since  $S_i \cap S_j \subseteq \{x \mid \phi_i(x) = \phi_j(x)\}$  and the latter has density at most  $1/p$ .

The third property follows from the “ $p$ -ary Johnson bound”: applied to the function  $g = (f^{(k+1)})^{(-k)} : G \rightarrow \mathbb{Z}_p$ . Let  $\psi_i = \phi_i^{(-k)} : G \rightarrow \mathbb{Z}_p$  (we warn the reader that  $\psi_i$  need not be an

<sup>1</sup>Here we abuse notation and identify an element of  $\mathbb{Z}_{p^{r-i}}$  with the least non-negative integer in its residue class mod  $p^{r-i}$ .

affine homomorphism). Note that the agreement set of  $\psi_i$  with  $g$  contains  $S_i$ . Furthermore, note that for  $i \neq j$ ,  $\psi_i \neq \psi_j$  and  $\text{agree}(\psi_i, \psi_j) = \text{agree}(\psi_i - \psi_j, 0) = \text{agree}(\phi_i - \phi_j, 0) \leq \frac{1}{p}$  (since  $\phi_i = p^k \cdot \psi_i + \phi$ ,  $\phi_j = p^k \cdot \psi_j + \phi$ ,  $\phi_i \neq \phi_j$ , and  $\text{image}(\phi_i - \phi_j) \subseteq p^k \cdot \mathbb{Z}_{p^{k+1}} \cong \mathbb{Z}_p$ ). Then  $\psi_1, \dots, \psi_\ell$  are distinct with pairwise agreement at most  $\frac{1}{p}$ , and each  $\psi_i$  has agreement at least  $1/p + \alpha_i$  with  $g$ . It follows from Lemma D.1 (in Appendix D) that  $\sum_{i=1}^{\ell} \alpha_i^2 \leq 1$ .

Finally, the fourth property can be verified exactly as in Claim 3.4, and the inductive proof of the claim is also derived exactly as in the proof of Claim 3.4. ■

The theorem statement follows immediately, by using  $k = 0$ . ■

## A.2 General abelian groups

Here we show how to use Lemma 3.6 to prove Theorem 2.2.

**Proof of Theorem 2.2** Let  $H = H' \times H''$ , where  $H' = \mathbb{Z}_{p_1}^{e_1} \times \dots \times \mathbb{Z}_{p_m}^{e_m}$  and  $H'' = \mathbb{Z}_{q_1}^{e_1} \times \dots \times \mathbb{Z}_{q_k}^{e_k}$ , such that  $p := p_1 \leq \dots \leq p_m$ ,  $p_i \mid |G|$  for each  $i \in [m]$ , while  $q_j \nmid |G|$  for each  $j \in [k]$ . Thus  $\Lambda = \frac{1}{p}$ .

Let  $f = (f', f'') : G \rightarrow H$  be any function, where  $f' : G \rightarrow H'$  and  $f'' : G \rightarrow H''$ . Let

$$\mathcal{L} = \{\phi \in \text{aHom}(G, H) : \text{agree}(\phi, f) \geq \Lambda + \epsilon\}.$$

Consider a  $\phi = (\phi', \phi'') \in \mathcal{L}$  where  $\phi' \in \text{aHom}(G, H')$  and  $\phi'' \in \text{aHom}(G, H'')$ . Then  $\text{agree}(\phi', f') \geq \Lambda + \epsilon$  and  $\text{agree}(\phi'', f'') \geq \Lambda + \epsilon$ . Thus

$$\phi' \in \mathcal{L}_1 := \{\psi' \in \text{aHom}(G, H') : \text{agree}(\psi', f') \geq \frac{1}{p} + \epsilon\},$$

and,

$$\phi'' \in \mathcal{L}_2 := \{\psi'' \in \text{aHom}(G, H'') : \text{agree}(\psi'', f'') \geq \frac{1}{p} + \epsilon\}.$$

By Lemma 3.6, the size of  $\mathcal{L}_1$  is at most  $\left(\frac{1}{\epsilon}\right)^{C_1}$ . Since  $\gcd(|G|, |H''|) = 1$ , the only functions in  $\text{aHom}(G, H'')$  are the constant functions (which pairwise disagree everywhere), and hence the size of  $\mathcal{L}_2$  is at most  $\frac{1}{\epsilon}$ .

Thus  $\mathcal{L} \subseteq \mathcal{L}_1 \times \mathcal{L}_2$ , and so

$$|\mathcal{L}| \leq |\mathcal{L}_1| \cdot |\mathcal{L}_2| \leq \left(\frac{1}{\epsilon}\right)^{C_1} \cdot \frac{1}{\epsilon}.$$

Taking  $C = C_1 + 1$ , the theorem follows. ■

## B Special Intersecting Families

We prove Theorem 3.2 by a sequence of lemmas. Fix a  $(\rho, \rho^2, c)$ -special intersecting family  $S_1, \dots, S_\ell \subseteq X$  and let  $\alpha_i = \mu(S_i) - \rho$ . Let  $\alpha = \mu(\cup_{i \in [\ell]} S_i) - \rho$ . Recall our goal is to prove that  $\alpha^C \geq \sum_{i=1}^{\ell} \alpha_i^C$  for a sufficiently large  $C$  (that depends only on  $c$ ).

The first three of the lemmas prove different lower bounds on  $\alpha$  under some special conditions, We glue these together to prove a weak variant of Theorem 3.2 in Lemma B.5. In Lemma B.6 we show that the weak condition actually implies the strong condition and this leads to the proof of Theorem 3.2 at the end of this section.

### B.1 A weak bound on the cardinality of the union

**Lemma B.1** *For all  $i \neq j \in [\ell]$ ,  $\alpha \geq \alpha_i + \alpha_j$ .*

**Proof** Simple inclusion-exclusion, using Property (2) of special intersecting families. We have

$$\rho + \alpha \geq \mu(S_i \cup S_j) = \mu(S_i) + \mu(S_j) - \mu(S_i \cap S_j) \geq (\rho + \alpha_i) + (\rho + \alpha_j) - \rho = \alpha_i + \alpha_j + \rho.$$

The lemma follows. ■

Let  $I \subseteq [\ell]$ . We say that  $I$  is a *collinear set* if either  $|I| \leq 2$  or  $\mu(S_I) > \rho^2$ . Collinear sets enjoy the following properties<sup>2</sup>:

1. If  $I$  is a collinear set, and  $J \subseteq I$ , then  $J$  is a collinear set. Indeed, if  $|J| \geq 2$ , then by property (4) of special intersecting families,  $S_J = S_I$  and  $J$  is collinear. If  $|J| < 2$ , then  $J$  is collinear by definition.
2. If  $I, J$  are collinear sets with  $|I \cap J| \geq 2$ , then  $I \cup J$  is a collinear set. To see this, notice that  $S_{I \cup J} = S_I \cap S_J$ , and by property (4) of special intersecting families,  $S_I = S_{I \cap J} = S_J$ . Thus  $S_{I \cup J} = S_I = S_J$ , and so  $I \cup J$  is a collinear set.
3. For  $i \neq j \in [\ell]$ , let  $L_{i,j} = \{i, j\} \cup \{k \in [\ell] \setminus \{i, j\} \mid \{i, j, k\} \text{ is a collinear set}\}$ . Then  $L_{i,j}$  is a collinear set. This follows easily from the previous property.

We say  $J \subseteq [\ell]$  is *in general position* if there is no collinear set  $K \subseteq J$  with  $|K| = 3$ .

First, in Lemma B.2, we obtain a lower bound on  $\alpha$  in terms of  $\sum_{i \in I} \alpha_i$ , for any collinear set  $I$ . Next, in Lemma B.3, we obtain a lower bound on  $\alpha$  in terms of  $\sum_{j \in J} \alpha_j$  for any  $J$  in general position. Finally, in Lemma B.5 we show a weak version of Theorem 3.2 by showing a dichotomy: either there is a collinear set  $I$  with  $\sum_{i \in I} \alpha_i$  large, or there is a set  $J$  in general position with  $\sum_{j \in J} \alpha_j$  large.

---

<sup>2</sup>We draw an analogy with a set of points, labelled by elements of  $[\ell]$ , on an abstract plane. Here a subset of  $[\ell]$  is called “collinear” if the corresponding points all lie on a single straight line.

**Lemma B.2** *If  $I \subseteq [\ell]$  is a collinear set, then  $\alpha \geq \sum_{i \in I} \alpha_i$ .*

**Proof** For distinct  $i_1, i_2 \in I$ , we know that  $S_{\{i_1, i_2\}} = S_I$ . Thus the sets  $\{S_i\}_{i \in I}$  form a sunflower, i.e, each set  $S_i$  is of the form  $R_i \cup S_I$ , where the  $R_i$ 's are pairwise disjoint. In turn this implies that

$$\begin{aligned}
\rho + \alpha &= \mu(\cup_{i \in [\ell]} S_i) \\
&\geq \mu(\cup_{i \in I} S_i) \\
&= \mu(S_I) + \sum_{k \in J} \mu(S_k - S_I) \\
&= \mu(S_{i_0}) + \sum_{i \in I \setminus \{i_0\}} \mu(S_i - S_I) \quad (\text{for some } i_0 \in I) \\
&\geq (\alpha_{i_0} + \rho) + \sum_{i \in I \setminus \{i_0\}} \alpha_i \\
&= \rho + \sum_{i \in I} \alpha_i.
\end{aligned}$$

We thus get  $\alpha \geq \sum_{i \in I} \alpha_i$  as claimed. ■

**Lemma B.3** *If  $J \subseteq [\ell]$  is in general position, and  $|J| \leq 1/\sqrt{\rho}$ , then  $\alpha \geq \frac{|J|-2}{3}\rho + \frac{1}{2} \sum_{j \in J} \alpha_j$ .*

**Proof** Before proving the lemma, we first state and prove a simple variant of the standard inclusion-exclusion lower bound on the cardinality of  $\cup_j S_j$ .

**Claim B.4** *For any set  $J \subseteq [\ell]$ ,*

$$\left| \bigcup_{j \in J} S_j \right| \geq \frac{1}{2} \sum_{j \in J} |S_j| - \frac{1}{2} \sum_{i, j, k \in J, i < j < k} |S_i \cap S_j \cap S_k|.$$

**Proof** Consider an element  $e \in \cup_{j \in J} S_j$  and say it is contained in  $m$  of the sets. Then  $e$  additively contributes  $m/2$  to the first term,  $\frac{1}{2} \sum_{j \in J} |S_j|$ . On the other hand in the second term it subtracts at least  $\frac{1}{2} \binom{m}{3}$ . Thus the net contribution of  $e$  to the RHS is at most  $\frac{m}{2} \left( 1 - \frac{(m-1)(m-2)}{6} \right)$ . It can be verified that this contribution is at most 1 for every positive integer  $m$ , which is also a lower bound on its contribution to the LHS. ■



We are now ready to prove the lemma. We have:

$$\begin{aligned}
\rho + \alpha &= \mu \left( \bigcup_{j \in [\ell]} S_j \right) \\
&\geq \mu \left( \bigcup_{j \in J} S_j \right) \\
&\geq \frac{1}{2} \sum_{j \in J} \mu(S_j) - \frac{1}{2} \sum_{i,j,k \in J, i < j < k} \mu(S_{\{i,j,k\}}) \quad (\text{Using Claim B.4}) \\
&\geq \frac{1}{2} \sum_{j \in J} (\rho + \alpha_j) - \frac{1}{2} \binom{|J|}{3} \rho^2 \quad (\text{Using definition of } \alpha_j \text{ and that } J \text{ is in general position}) \\
&= \frac{|J|}{2} \rho - \frac{1}{2} \binom{|J|}{3} \rho^2 + \frac{1}{2} \sum_{j \in J} \alpha_j
\end{aligned}$$

Rearranging terms above, we get

$$\begin{aligned}
\alpha &\geq \frac{|J|-2}{2} \rho - \frac{1}{2} \binom{|J|}{3} \rho^2 + \frac{1}{2} \sum_{j \in J} \alpha_j \\
&= \frac{|J|-2}{2} \cdot \rho \cdot \left( 1 - \frac{|J|(|J|-1)}{3} \rho \right) + \frac{1}{2} \sum_{j \in J} \alpha_j \\
&\geq \frac{|J|-2}{3} \cdot \rho + \frac{1}{2} \sum_{j \in J} \alpha_j,
\end{aligned}$$

as claimed. ■

We are now ready to put the above ingredients together to prove a weak version of Theorem 3.2. Specifically the following lemma claims that some positive power of  $\alpha$  is lower bounded by some multiple of the sum of the  $\alpha_i$ 's to the same power. Unfortunately, the multiple is a pretty weak one and becomes smaller as the number of summands grow (whereas in Theorem 3.2 we want this multiple to be 1!). Nevertheless this weak version is good enough, as we'll see later.

**Lemma B.5** *For every  $c < \infty$  there exist constants  $c_1, c_2, c_3$  (depending only on  $c$ ) with  $c_1, c_3 < \infty$  and  $c_2 < 1$  such that for every set  $K \subseteq [\ell]$ ,*

$$\alpha^{c_1} \geq \frac{1}{c_3 \cdot |K|^{c_2}} \cdot \sum_{k \in K} \alpha_k^{c_1}.$$

Remark: Note that if the constant  $c_2$  in the exponent of  $|K|$  had been replaced by 1, the lower bound on  $\alpha$  would have been quite trivial (with  $c_1, c_2 = 3$ ). The slightly smaller exponent turns out to be sufficiently powerful to let us derive Theorem 3.2 later.

**Proof** Fix a set  $K \subseteq [\ell]$ .

We prove the lemma for  $c_1 = c + 1$ . (The other constants will be determined later.) Let  $P = \{k \in K \mid \alpha_k < \rho^{c_1}\}$  and  $Q = \{k \in K \mid \alpha_k \geq \rho^{c_1}\}$ . We consider two cases based on whether  $\sum_{k \in P} \alpha_k^{c_1}$  is larger than  $\sum_{k \in Q} \alpha_k^{c_1}$  or not.

Case 1:  $\sum_{k \in P} \alpha_k^{c_1} \geq \sum_{k \in Q} \alpha_k^{c_1}$ : If  $|P| = 1$ , then it must be that  $|Q| = 0$  and then  $|K| = 1$  and if so this case is trivial. So  $|P|$  must be at least 2 in this case. Again we divide the analysis into two subcases.

The first subcase we consider is when there exist distinct  $i, j, k \in P$  with  $\{i, j, k\}$  not collinear. In this case, by applying Lemma B.3 to the set  $J = \{i, j, k\}$ , we get  $\alpha \geq \rho/3$ . On the other hand, we have

$$\begin{aligned} \sum_{k \in P} \alpha_k^{c+1} &\leq \max_{k \in P} \alpha_k \cdot \sum_{j \in P} \alpha_j^c \\ &\leq \max_{k \in P} \alpha_k \cdot 1 \quad (\text{Using the third property of the set systems}) \\ &\leq \rho^{c_1} \quad (\text{By definition of } P). \end{aligned}$$

So in this subcase we have

$$\begin{aligned} \alpha^{c_1} &\geq \rho^{c_1} / (3^{c_1}) \\ &\geq \frac{1}{3^{c_1}} \sum_{k \in P} \alpha_k^{c_1} \\ &\geq \frac{1}{2 \cdot 3^{c_1}} \sum_{k \in K} \alpha_k^{c_1} \quad (\text{Since } P \text{ contains more than half the weight of } K) \\ &\geq \frac{1}{c_3 |K|^{c_2}} \sum_{k \in K} \alpha_k^{c_1} \quad (\text{provided } c_3 \geq 2 \cdot 3^{c_1} \text{ and } c_2 \geq 0.) \end{aligned}$$

Now we move to the second subcase. In this case, we assume that for every triple  $i, j, k$  in  $P$ ,  $\{i, j, k\}$  is collinear. This implies that  $P$  itself is collinear. By Lemma B.2, we have  $\alpha \geq \sum_{k \in P} \alpha_k$ . Thus  $\alpha^{c_1} \geq \sum_{k \in P} \alpha_k^{c_1} \geq \frac{1}{2} \sum_{k \in K} \alpha_k^{c_1}$ . Thus again we have  $\alpha^{c_1} \geq \frac{1}{c_3 |K|^{c_2}} \sum_{k \in K} \alpha_k^{c_1}$  provided  $c_3 \geq 2$  and  $c_2 \geq 0$ . This concludes the analysis of Case 1.

Case 2:  $\sum_{k \in P} \alpha_k^{c_1} < \sum_{k \in Q} \alpha_k^{c_1}$ : Note that this is the case, where our lower bound on  $\alpha$  will start deteriorating with  $|K|$ .

Note that in this case  $|Q|$  is at most  $\frac{1}{\rho^{c_1}}$  by Property (3) of the  $(\rho, \rho^2, c)$ -special intersection condition. We divide our analysis into two subcases again. For a subset  $I \subseteq K$ , let  $\text{wt}(I) = \sum_{i \in I} \alpha_i^{c_1}$ .

Let  $c_2 = 1 - 1/(2c \cdot c_1)$ .

The first subcase is when there exists a collinear  $I \subseteq K$  such that  $\text{wt}(I) \geq \frac{\text{wt}(Q)}{|Q|^{c_2}}$ . If so, by Lemma B.2, we have that  $\alpha \geq \sum_{i \in I} \alpha_i$  and so

$$\alpha^{c_1} \geq \sum_{i \in I} \alpha_i^{c_1} \geq \frac{\text{wt}(Q)}{|Q|^{c_2}} \geq \frac{\text{wt}(K)}{2|Q|^{c_2}} \geq \frac{\text{wt}(K)}{2|K|^{c_2}}.$$

This yields the lemma for every  $c_3 \geq 2$ .

Now we consider the final remaining subcase, where for every collinear  $I \subseteq K$ , it is the case that  $\text{wt}(I) < \frac{\text{wt}(Q)}{|Q|^{c_2}}$ . We first prove that there is a  $J \subset Q$  in general position such that  $|J| = |Q|^{1/(2c \cdot c_1)}$  and  $\text{wt}(J) \geq \text{wt}(Q)/|Q|^{c_2}$ . Applying Lemma B.3 to this set then completes the analysis for this case.

We start with showing that such a set  $J$  exists. We pick  $J$  greedily starting with  $J = \emptyset$ . We maintain a set of candidates  $Q'$ , initially containing all of  $Q$ . We then add the element  $i \in Q'$  with the largest value of  $\alpha_i$  to  $J$ . We remove  $i$  from  $Q'$  as also every element  $k \in Q'$  such that for some  $j \in J \in \{i\}$ ,  $\{i, j, k\}$  is collinear. We repeat until  $|J| = |Q|^{1/(2c \cdot c_1)}$ .

To analyze this construction, we first claim that  $Q'$  does not become empty (which would preclude us from growing  $J$ ). Note that for every  $i, j \in J$ , the total weight of the elements removed from  $Q'$  on account of  $i, j$  is at most  $\text{wt}(L_{i,j}) < \text{wt}(Q)/|Q|^{c_2}$ . Since there are at most  $|J|^2/2 \leq |Q|^{1/(c \cdot c_1)}/2$  such pairs, the total weight of the removed elements is at most  $\text{wt}(Q)/(2 \cdot |Q|^{c_2 - 1/(c \cdot c_1)}) < \text{wt}(Q)/2$ , provided  $c_2 > 1/(c \cdot c_1)$ , which holds for  $c_2 \geq 1/2$  and  $c \geq 1$  and  $c_1 \geq 2$ .

Thus, the process concludes with  $|J| = |Q|^{1/(2c \cdot c_1)}$  and  $J$  is in general position by construction. To see that the weight of  $J$  is large, notice that every element of  $J$  has weight at least that of the average element, i.e.,  $\alpha_k^{c_1} \geq \text{wt}(Q')/|Q'| \geq \text{wt}(Q)/(2|Q|)$ . We thus conclude that  $\text{wt}(J) \geq |J| \cdot \text{wt}(Q)/(2|Q|) \geq \text{wt}(Q)/(2|Q|^{1 - 1/(2c \cdot c_1)}) = \text{wt}(Q)/(2|Q|^{c_2})$ .

Finally, note that  $|J| = |Q|^{1/(2c \cdot c_1)} \leq (1/(\rho^{c \cdot c_1}))^{1/(2c \cdot c_1)} = 1/\sqrt{\rho}$ . So this puts us in a position to apply Lemma B.3 to the set  $J$  to conclude that  $\alpha \geq \frac{1}{2} \sum_{j \in J} \alpha_j$ . This in turn yields  $\alpha^{c_1} \geq \frac{1}{2^{c_1}} (\sum_{j \in J} \alpha_j)^{c_1} \geq \frac{1}{2^{c_1}} \sum_{j \in J} \alpha_j^{c_1} \geq \frac{1}{2^{c_1}} \text{wt}(Q)/|Q|^{c_2} \geq \frac{1}{2^{c_1+1}} \text{wt}(K)/|K|^{c_2}$ . This yields the lemma statement for  $c_2 = 1 - 1/(2c \cdot c_1)$  and  $c_3 \geq 2^{1+c_1}$ .

Putting all the cases together, we get the lemma statement for  $c_2 = \max\{1 - 1/(2c \cdot c_1), \frac{1}{2}\}$  and  $c_3 = 2 \cdot 3^{c_1}$  (and, of course,  $c_1 = \max\{2, c + 1\}$ ). ■

## B.2 Strengthening the bound

Our final lemma shows how to convert the assertion of Lemma B.5, in conjunction with Lemma B.1, to get the assertion of Theorem 3.2.

**Lemma B.6** *For every  $c_1 < \infty$ ,  $c_2 < 1$ , and  $c_3 < \infty$  there exists  $C < \infty$  such that the following holds: Let  $\alpha, \alpha_1, \dots, \alpha_\ell$  be non-negative reals such that (1)  $\alpha \geq \alpha_i + \alpha_j$  for every  $i \neq j \in [\ell]$  and (2) For every  $K \subseteq [\ell]$ ,  $\alpha^{c_1} \geq \frac{1}{c_3 |K|^{c_2}} \cdot \sum_{k \in K} \alpha_k^{c_1}$ . Then  $\alpha^C \geq \sum_{k=1}^\ell \alpha_k^C$ .*

**Proof** In what follows, we assume without loss of generality that  $c_1 = 1$ . The reason we can do so is the following: Suppose the lemma holds for some constant  $C$  when  $c_1 = 1$  and some  $c_2, c_3$ . Then, by applying it to the sequence  $\beta, \beta_1, \dots, \beta_\ell$  for  $\beta = \alpha^{c_1}$  and  $\beta_i = \alpha_i^{c_1}$ , we get the lemma for general  $c_1$  for the constant  $C * c_1$ . Thus, from now on we assume  $c_1 = 1$ . By scaling all the  $\alpha_i$ 's and  $\alpha$  and sorting the  $\alpha_i$ 's, we may also assume  $\alpha_1 = 1$  and  $\alpha_i \geq \alpha_{i+1}$ .

We prove the lemma for (every)  $C \geq \frac{\log_2(8c_3)}{1-c_2}$ .

In what follows, we attempt to find the maximum value of  $\sum_{i=1}^{\ell} \beta_i^C$  for any sequence  $1 = \beta_1 \geq \beta_2 \geq \dots \geq \beta_{\ell}$  that satisfy the properties (1)  $\beta_1 + \beta_2 \leq \alpha$ ; and (2)  $\beta_1 + \dots + \beta_k \leq c_3 k^{c_2} \cdot \alpha$  for every  $k \in [\ell]$ . The following claim will prove to be a useful tool in this maximization.

**Claim B.7** *Suppose  $a_1 \geq \dots \geq a_{\ell} \geq 0$  and  $b_1 \geq \dots \geq b_{\ell}$  satisfy  $\sum_{i=1}^k a_i \geq \sum_{i=1}^k b_i$  for every  $k \in [\ell]$ . Then, for every  $C > 1$ ,  $\sum_{i=1}^{\ell} a_i^C \geq \sum_{i=1}^{\ell} b_i^C$ .*

**Proof** Consider the set of sequences  $d_1, \dots, d_{\ell}$  that satisfy the conditions  $\sum_{i=1}^k d_i \leq \sum_{i=1}^k a_i$  for every  $k$  and  $d_1 \geq \dots \geq d_{\ell}$ . This is a closed compact set and so there is an element in this set that maximizes the sum  $\sum_{i=1}^{\ell} d_i^C$ . We claim that this sequence is  $a_1, \dots, a_{\ell}$ .

If not, and suppose  $d_1, \dots, d_{\ell}$  maximize this sum. Let  $i_0$  be the smallest index such that  $d_{i_0} \neq a_{i_0}$ . Since  $\sum_{j=1}^{i_0-1} d_j = \sum_{j=1}^{i_0-1} a_j$ , and  $\sum_{j=1}^{i_0} d_j \leq \sum_{j=1}^{i_0} a_j$ , it must be that  $d_{i_0} < a_{i_0}$ . Since  $\sum_{j=1}^{\ell} d_j^C > \sum_{j=1}^{\ell} a_j^C$ , there must also be an index  $j$  such that  $a_j < d_j$ . Let  $j_1$  be the smallest index such that  $a_{j_1} < d_{j_1}$ . Note that  $i_0 < j_1$ . Now let  $i_1$  be the largest index less than  $j_1$  such that  $a_{i_1} > d_{i_1}$ . Then it must be that  $a_k = d_k$  for all  $k \in \{i_1 + 1, \dots, j_1 - 1\}$ . Now let  $i_2$  be the smallest index such that  $d_{i_2} = d_{i_1}$  and let  $j_2$  be the largest index such that  $d_{j_2} = d_{j_1}$ . Let  $\epsilon = \min\{a_{i_2} - d_{i_2}, d_{i_2-1} - d_{i_2}, d_{j_2} - a_{j_2}, d_{j_2} - d_{j_2+1}\}$ . Now consider the sequence  $e_1, \dots, e_{\ell}$  with  $e_{i_2} = d_{i_2} + \epsilon$  and  $e_{j_2} = d_{j_2} - \epsilon$ , and  $e_k = d_k$  otherwise. We claim that the sequence  $e_1, \dots, e_{\ell}$  is (A) non-increasing, (B) satisfies  $\sum_{i=1}^k e_i \leq \sum_{i=1}^k a_i$  for every  $k \in [\ell]$ , and (C)  $\sum_{i=1}^{\ell} e_i^C > \sum_{i=1}^{\ell} d_i^C$ . If so, this yields a contradiction to the maximality of  $d_i$ 's.

We start with the final property first. For this we note that  $\epsilon > 0$ . This follows from the facts that (i)  $a_{i_2} - d_{i_2} \geq a_{i_1} - d_{i_1} > 0$ , (ii)  $i_2$  was the smallest index with  $d_{i_2} = d_{i_1}$  and so  $d_{i_2} - d_{i_1} > 0$  and similarly (iii)  $d_{j_2} - a_{j_2} > 0$  and (iv)  $d_{j_2} > d_{j_2+1}$ . So  $\sum_{i=1}^{\ell} e_i^C - \sum_{i=1}^{\ell} d_i^C = e_{i_2}^C - d_{i_2}^C + e_{j_2}^C - d_{j_2}^C = (d_{i_2} + \epsilon)^C - d_{i_2}^C + (d_{j_2} - \epsilon)^C - d_{j_2}^C$ . From the strict convexity of the function  $f(x) = x^C$  when  $x \geq 0$ , and the fact that  $d_{i_2} \geq d_{j_2}$  it follows that  $(d_{i_2} + \epsilon)^C - d_{i_2}^C + (d_{j_2} - \epsilon)^C - d_{j_2}^C > 0$ .

We now show that the  $e_i$ 's are non-increasing. The only possible violations would involve  $e_{i_2}$  or  $e_{j_2}$ . First note that  $e_{i_2} > d_{i_2} \geq d_{i_2+1} \geq e_{i_2+1}$  (specifically  $d_{i_2+1} = e_{i_2+1}$  unless  $i_2 + 1 = j_2$  in which case  $d_{j_2} > e_{j_2}$ ). Similarly it can be checked that  $e_{j_2-1} \geq e_{j_2}$ . Finally, the choice of  $\epsilon$  ensures  $e_{i_2-1} \leq e_{i_2}$  and  $e_{j_2} \leq e_{j_2+1}$ .

Finally to verify the prefix sum conditions, note that if  $k < i_2$  or  $k \geq j_2$ , then  $\sum_{i=1}^k e_i = \sum_{i=1}^k d_i \leq \sum_{i=1}^k a_i$ . It remains to prove the condition for  $k \in \{i_2, \dots, j_2 - 1\}$ . To do so we note the following inequalities hold:  $d_i \leq a_i$  for  $i \in \{1, \dots, j_1 - 1\}$  with  $d_i \leq a_i - \epsilon$  for  $i = i_2$ . Furthermore,  $d_i = d_{j_1} > a_{j_1} \geq a_i$  for  $i \geq j_1$  with  $d_i \geq a_i + \epsilon$  for  $i = j_1$ . Thus if  $i_2 \leq k < j_1$ , we have

$$\sum_{i=1}^k e_i = \sum_{i=1}^k d_i + \epsilon \leq \sum_{i=1}^k a_i,$$

as desired. On the other hand, if  $j_1 \leq k < j_2$ , we have

$$\begin{aligned}
\sum_{i=1}^k e_i &= \sum_{i=1}^k d_i + \epsilon \\
&= \sum_{i=1}^{j_2} d_i - \sum_{i=k+1}^{j_2} d_i + \epsilon \\
&\leq \sum_{i=1}^{j_2} a_i - \sum_{i=k+1}^{j_2} d_i + \epsilon \\
&\quad \text{(Since } d_i\text{'s form a prefix sum respecting sequence)} \\
&< \sum_{i=1}^{j_2} a_i - \sum_{i=k+1}^{j_2-1} a_i - (d_{j_2} - \epsilon) \\
&\leq \sum_{i=1}^{j_2} a_i - \sum_{i=k+1}^{j_2} a_i \\
&= \sum_{i=1}^k a_i
\end{aligned}$$

Thus  $e_i$ 's form a non-increasing, prefix sum respecting sequence that achieve a strictly larger sum  $\sum_{i=1}^{\ell} e_i^C$  violating the maximality of  $C$ . This proves the claim. ■

We now return to the proof of the lemma. We consider two cases:

Case 1:  $\alpha \geq 2$ : Let  $k = \lceil (c_3\alpha)^{1/(1-c_2)} \rceil$ , so that  $c_3\alpha k^{c_2} \leq k$ . Now consider the sequence  $\beta_1, \dots, \beta_\ell$  with  $\beta_i = 1$  for  $i \leq k$  and  $\beta_i = c_3\alpha(i^{c_2} - (i-1)^{c_2})$  for  $i > k$ .

For every  $i \in [\ell]$ , we claim that  $\sum_{j=1}^i \alpha_j \leq \sum_{j=1}^i \beta_j$ . For  $i \leq k$ , this is true since  $\sum_{j=1}^i \alpha_j \leq i \cdot \alpha_1 \leq i = \sum_{j=1}^i \beta_j$ . For  $i > k$ , we have  $\sum_{j=1}^i \alpha_j \leq c_3\alpha i^{c_2}$ , while

$$\sum_{j=1}^i \beta_j = k + \sum_{j=k+1}^i \beta_j \leq c_3\alpha k^{c_2} + c_3\alpha \sum_{j=k+1}^i (j^{c_2} - (j-1)^{c_2}) = c_3\alpha i^{c_2}.$$

Thus we can apply Claim B.7 and get:

$$\begin{aligned}
\sum_{i=1}^{\ell} \alpha_i^C &\leq \sum_{i=1}^{\ell} \beta_i^C \\
&\leq k + (c_3\alpha)^C \sum_{i=k+1}^{\ell} (i^{c_2} - (i-1)^{c_2})^C \\
&\leq k + (c_3\alpha)^C \sum_{i=k+1}^{\ell} i^{-C(1-c_2)} \\
&\leq k + (c_3\alpha)^C k^{-C(1-c_2)+1} \quad (\text{Using } C(1-c_2) \geq 2) \\
&= k + (c_3\alpha/k^{1-c_2})^C k^1 \\
&\leq 2k \\
&\leq 2(1 + (c_3\alpha)^{1/(1-c_2)}) \\
&\leq 4(c_3\alpha)^{1/(1-c_2)}
\end{aligned}$$

Thus to conclude this case, it suffices to prove that for sufficiently large  $C$ ,  $4(c_3\alpha)^{1/(1-c_2)} \leq \alpha^C$ , which follows if  $4c_3^{1/(1-c_2)} \leq \alpha^{(C-1)/(1-c_2)}$ . which in turn follows if  $(4c_3) \leq 2^{(C(1-c_2)-1)}$ , which holds if  $C \geq \frac{1}{1-c_2}(1 + \log_2(4c_3)) = \frac{\log_2(8c_3)}{1-c_2}$ . This completes the analysis for Case 1.

Case 2:  $\alpha < 2$ . Let  $\tau = \alpha - 1$ . Let  $k = \lceil (c_3\alpha/\tau)^{1/(1-c_2)} \rceil$  so that  $c_3\alpha k^{c_2} \leq \tau k \leq 1 + (k-1)\tau$ . Let  $\beta_i = 1$  if  $i = 1$ ,  $\beta_i = \tau$  if  $2 \leq i \leq k$  and  $\beta_i = c_3(1 + \tau)(i^{c_2} - (i-1)^{c_2})$  for  $i \geq k+1$ . As in Case 1, it can be verified that  $\sum_{j=1}^i \alpha_j \leq \sum_{j=1}^i \beta_j$ . Applying Claim B.7, we get

$$\begin{aligned}
\sum_{i=1}^{\ell} \alpha_i^C &\leq \sum_{i=1}^{\ell} \beta_i^C \\
&\leq 1 + (k-1)\tau^C + (c_3\alpha)^C \sum_{i=k+1}^{\ell} (i^{c_2} - (i-1)^{c_2})^C \\
&\leq 1 + (k-1)\tau^C + (c_3\alpha)^C \sum_{i=k+1}^{\ell} (i^{-C(1-c_2)}) \\
&\leq 1 + (k-1)\tau^C + (c_3\alpha)^C (k^{-C(1-c_2)+1}) \\
&= 1 + (k-1)\tau^C + (c_3\alpha/k^{1-c_2})^C \cdot k \\
&\leq 1 + 2k\tau^C
\end{aligned}$$

Using the crude bound  $2k \leq (4c_3(1 + \tau)/\tau)^{1/(1-c_2)}$  we see that it suffices to prove that  $1 + (4c_3(1 + 1/\tau))^{1/(1-c_2)}\tau^C \leq (1 + \tau)^C$  which is equivalent to proving that  $(4c_3(1 + 1/\tau))^{1/(1-c_2)} \leq (1 + 1/\tau)^C - 1/\tau^C$ . Using  $(x + 1)^C - x^C \geq (x + 1)^{C-1}$ , we find that it suffices  $(4c_3)^{1/(1-c_2)} \leq (1 + 1/\tau)^{C-1/(1-c_2)}$  which holds if  $(4c_3) \leq 2^{(C(1-c_2)-1)}$ , which in turn holds if  $C \geq \frac{1}{1-c_2} \log_2(8c_3)$ .

■

**Proof of Theorem 3.2** Follows by combining Lemmas B.1, B.5 and B.6. ■

## C Algorithmic Results

In this section, we sketch a proof of Theorem 2.4. As we shall see, the combinatorial bounds that we proved make the algorithm relatively straightforward. Our algorithm can be viewed as a direct extension of the original list-decoder of Goldreich and Levin [3].

We are given a function  $f : G \rightarrow H$ . We want to locally list decode all homomorphisms  $\phi \in \text{Hom}(G, H)$  such that  $\text{agree}(f, \phi) \geq \Lambda_{G, H} + \epsilon$ .

We first assume  $H = \mathbb{Z}_{p^r}$ . We will later use the algorithm for this special case to handle the case of general  $H$ .

Let  $\mathcal{L} = \{\phi \in \text{Hom}(G, H) : \exists b \in H \text{ s.t. } \text{agree}(\phi + b, f) \geq \Lambda_{G, H} + \epsilon\}$ . Let  $G = \bigoplus_{i=1}^k \mathbb{Z}_{p_i^{r_i}}$ . For  $t \in [k]$ , let  $G_t = \bigoplus_{i=1}^t \mathbb{Z}_{p_i^{r_i}}$  and  $F_t = \bigoplus_{i=t+1}^k \mathbb{Z}_{p_i^{r_i}}$ . We say  $\phi \in \text{Hom}(G_t, H)$ , extends to  $\phi' \in \text{Hom}(G, H)$  if there exists  $\theta \in \text{Hom}(F_t, H)$  with  $\phi'(x, y) = \phi(x) + \theta(y)$ . For any homomorphism  $\phi' \in \text{Hom}(G, H)$ , there is exactly one homomorphism  $\phi \in \text{Hom}(G_t, H)$  such that  $\phi$  extends to  $\phi'$ . Let  $\Phi_t = \{\phi \in \text{Hom}(G_t, H) : \exists \phi' \in \mathcal{L} \text{ s.t. } \phi \text{ extends to } \phi'\}$ . Our first goal will be to compute  $\Phi_k$ .

We start with  $t = 1$  and set  $S_0 = \emptyset$ . For any given  $t$ , given a set  $S_{t-1}$  containing  $\Phi_{t-1}$ , we will compute a set  $S_t$  containing  $\Phi_t$ . This is done in two phases, *extending* and *pruning*.

**Extending:** Note that  $G = G_{t-1} \times \mathbb{Z}_{p_t^{r_t}} \times F_t$ . If  $p_t \neq p$ , then we just set  $S_t = S_{t-1}$  and exit. Otherwise, we proceed as follows.

For each  $\phi \in S_{t-1}$ , do the following  $\text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon})$  times:

1. Pick  $s \in F_t$  at random.
2. Repeat the following  $\text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon})$  times.
  - (a) Pick  $x_1, x_2 \in G_{t-1}$  and  $c_1, c_2 \in \mathbb{Z}_{p_t^{r_t}}$
  - (b) If  $c_1 - c_2$  is not divisible by  $p$ , solve the system of equations for  $a, s \in \mathbb{Z}_{p^r}$ :

$$\phi(x_1) + ac_1 + s = f(x_1, c_1, s)$$

$$\phi(x_2) + ac_2 + s = f(x_2, c_2, s)$$

- (c) Let  $\theta \in \text{Hom}(G_{t-1} \times \mathbb{Z}_{p_t^{r_t}}, H)$  be given by  $\theta(x, c) = \phi(x) + a \cdot c \pmod{p^r}$ . Add  $\theta$  to  $S_t$ .

It can be shown that if  $\Phi_{t-1} \subseteq S_{t-1}$ , then with high probability,  $\Phi_t \subseteq S_t$ .

**Pruning:** Now we write  $G = G_t \times F_t$ . In the pruning phase, we will cut down the size of  $S_t$  while preserving the fact that  $\Phi_t \subseteq S_t$ .

Repeat  $\text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon})$  times:

1. Pick a random  $s \in F_t$ .
2. For each  $\phi \in S_t$ , by random sampling, find all  $b \in H$  such that  $\Pr_{x \in G_t}[\phi(x, s) - f(x, s) = b] \geq \Lambda_{G,H} + \epsilon/2$ .
3. If there exists even one  $b$  as above, then add  $\phi$  to  $S'_t$ .

Upon conclusion, replace  $S_t$  with  $S'_t$ .

Our list decoding bound implies that after the pruning phase, the cardinality of  $S_t$  is at most  $\text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon})$  for some fixed polynomial. Furthermore, with high probability, all the  $\phi \in \Phi_t$  remain in  $S_t$ .

After executing these two phases for  $t = 1 \dots, k$ , we get a set  $S_k \supseteq \Phi_k$ . Now by random sampling, we may test the elements of  $S_k$  and their affine shifts for agreement with  $f$  to get the final list. This concludes the description of the list decoding algorithm for  $H = \mathbb{Z}_p^r$ .

**General  $H$**  Using the above algorithm, we can now handle the case of general  $H$ . We induct on the number of components in  $H$ . Suppose  $H = H_1 \times \mathbb{Z}_p^r$ . Let  $f = (f_1, f_2)$  where  $f_1 : G \rightarrow H_1$  and  $f_2 : G \rightarrow \mathbb{Z}_p^r$ . The algorithm list decodes from  $f_1$  (by induction) and from  $f_2$  (by the algorithm above), and takes the product of the resulting lists. This list may now be pruned by random sampling to verify agreement. By the combinatorial list decoding bound, we have that the resulting list's size is at most  $(\frac{1}{\epsilon})^C$ . This is the list that we output. It is crucial for the induction that the size of the list is a polynomial independent of the size of  $H$ .

It can be easily seen that the running time of the algorithm is  $\text{poly}(\log |G|, \log |H|, \frac{1}{\epsilon})$ .

## D $p$ -ary Johnson Bound

**Lemma D.1** *Let  $f, \phi_1, \dots, \phi_\ell : [n] \rightarrow [q]$  be functions satisfying the following properties:*

1.  $\text{agree}(f, \phi_i) = \frac{1}{q} + \alpha_i$  for  $\alpha_i \geq 0$ .
2.  $\text{agree}(\phi_j, \phi_i) \leq \frac{1}{q}$  for every  $i \neq j$ .

Then  $\sum_{i=1}^{\ell} \alpha_i^2 \leq 1$ .

**Proof** Let  $\zeta_1, \dots, \zeta_q \in \mathbb{R}^{q-1}$  be the vertices of the  $q$ -point simplex, i.e., they satisfy  $\langle \zeta_i, \zeta_i \rangle = 1$  and  $\langle \zeta_i, \zeta_j \rangle = -1/(q-1)$  if  $i \neq j$ . Fix a bijection  $\pi[q] \rightarrow \{\zeta_1, \dots, \zeta_q\}$ , and use  $\pi$  to convert the functions  $f : [n] \rightarrow [q]$  to functions  $g, \psi_1, \dots, \psi_\ell : [n] \rightarrow (\mathbb{R}^{q-1})$  using  $g = \pi \circ f$  and  $\psi_i = \pi \circ \phi_i$ . Note that for any two functions  $h, k : [n] \rightarrow [q]$ , we have  $\langle \pi \circ h, \pi \circ k \rangle \stackrel{\text{def}}{=} \mathbb{E}_{x \in [n]} [\pi(h(x)) \cdot \pi(k(x))]$  is given by  $\langle \pi \circ h, \pi \circ k \rangle = (q/(q-1))\text{agree}(h, k) - 1/(q-1)$ . Thus we get:  $\langle g, \psi_i \rangle = (q/(q-1))\alpha_i \geq 0$  and  $-1/(q-1) \leq \langle \psi_i, \psi_j \rangle \leq 0$ . This implies, via Lemma D.2,  $\sum_{i=1}^{\ell} \langle g, \psi_i \rangle^2 \leq \langle g, g \rangle^2$  and so  $\sum_{i=1}^{\ell} \alpha_i^2 \leq (q-1)/q$ . ■



**Lemma D.2** Let  $w, v_1, \dots, v_n \in \mathbb{R}^m$  such that:

1.  $\langle v_i, v_i \rangle = 1$  for each  $i \in [n]$ .
2.  $\langle w, v_i \rangle \geq 0$  for each  $i \in [n]$ .
3.  $\langle v_i, v_j \rangle \leq 0$  for any distinct  $i, j \in [n]$ .

Then

$$\sum_{i=1}^n \langle w, v_i \rangle^2 \leq \langle w, w \rangle.$$

**Proof** Let  $u_1, \dots, u_n$  be the Gram-Schmidt orthogonalization of  $v_1, \dots, v_n$ . Explicitly,

$$u_i = v_i - \sum_{j=1}^{i-1} u_j^* \langle u_j^*, v_i \rangle,$$

where for a vector  $u$ ,  $u^*$  is the scalar multiple of  $u$  with  $\langle u^*, u^* \rangle = 1$ . (Notice that each  $u_j$  has magnitude  $\leq 1$ ).

We will check the following two claims:

1. For each  $i < k$ ,  $\langle u_i, v_k \rangle \leq 0$ .

By induction on  $i$ : When  $i = 1$ , it is clearly true for any  $k$ . Assume we know it for any  $(i', k')$  where  $i' < i$  and  $k' \leq k$ .

$$\begin{aligned} \langle u_i, v_k \rangle &= \langle v_i, v_k \rangle - \sum_{j=1}^{i-1} \langle u_j^*, v_k \rangle \langle u_j^*, v_i \rangle \\ &\leq \langle v_i, v_k \rangle \quad (\text{since each term of the summation is nonnegative} \\ &\quad \text{by induction hypothesis on } (j, k) \text{ and } (j, i)) \\ &\leq 0 \end{aligned}$$

2. For each  $i$ ,  $\langle w, u_i \rangle \geq \langle w, v_i \rangle \geq 0$ .

By induction on  $i$  again. assume we know it for any  $i' < i$ .

$$\langle w, u_i \rangle = \langle w, v_i \rangle - \sum_{j=1}^{i-1} \langle w, u_j^* \rangle \langle u_j^*, v_i \rangle \geq \langle w, v_i \rangle.$$

(because each term in the sum is negative, by the first claim above and the induction hypothesis.)

Thus  $\langle w, v_i \rangle \leq \langle w, u_i \rangle \leq \langle w, u_i^* \rangle$ , (since  $\langle w, u_i \rangle \geq 0$  and  $u_i$  has magnitude at most 1).

But we know that  $\sum_{i=1}^n \langle w, u_i^* \rangle^2 \leq 1$ . So  $\sum_{i=1}^n \langle w, v_i \rangle^2 \leq 1$ .

■