

Lecture 33: Working Example for RSA Assumption

Recall: RSA Assumption

- We pick two primes uniformly and independently at random
 $p, q \xleftarrow{\$} P_n$
- We define $N = p \cdot q$
- We shall work over the group (\mathbb{Z}_N^*, \times) , where \mathbb{Z}_N^* is the set of all natural numbers $< N$ that are relatively prime to N , and \times is integer multiplication mod N
- We pick $y \xleftarrow{\$} \mathbb{Z}_N^*$
- Let $\varphi(N)$ represent the size of the set \mathbb{Z}_N^* , which is $(p - 1)(q - 1)$
- We pick any $e \in \mathbb{Z}_{\varphi(N)}^*$, that is, e is a natural number $< \varphi(N)$ and is relatively prime to $\varphi(N)$
- We give (n, N, e, y) to the adversary \mathcal{A} as ask her to find the e -th root of y , i.e., find x such that $x^e = y$

RSA Assumption. For any computationally bounded adversary, the above-mentioned problem is hard to solve

Working Example I

- We shall use $p = 3$ and $q = 11$
- So, we have $N = p \cdot q = 33$
- Moreover, we have

$$\mathbb{Z}_N^* = \{1, 2, 4, 5, 7, 8, 10, 13, 14, 16, 17, 19, 20, 23, 25, 26, 28, 29, 31, 32\}$$

- Now, $\varphi(N) = (p - 1)(q - 1) = 2 \cdot 10 = 20$. Verify that this is the size of \mathbb{Z}_N^*
- For this example, we shall choose $e = 3$ (note that 3 is relatively prime to $\varphi(N) = 20$)

Working Example II

Let us start the repeated squaring procedure. The first row represents each element of \mathbb{Z}_N^* and the second row is the square of the corresponding element in the first row.

x	1	2	4	5	7	8	10	13	14	16	17	19	20	23	25	26	28	29	31	32
x^2	1	4	16	25	16	31	1	4	31	25	25	31	4	1	31	16	25	16	4	1

Working Example III

Using repeated squaring, we compute the third row that is the fourth-power of the element in the first row.

x	1	2	4	5	7	8	10	13	14	16	17	19	20	23	25	26	28	29	31	32
x^2	1	4	16	25	16	31	1	4	31	25	25	31	4	1	31	16	25	16	4	1
x^4	1	16	25	31	25	4	1	16	4	31	31	4	16	1	4	25	31	25	16	1

Working Example IV

We add a row that computes $y = x^e$ (recall that $e = 3$ in our case). We can obtain x^3 by multiplying $x \times x^2$.

x	1	2	4	5	7	8	10	13	14	16	17	19	20	23	25	26	28	29	31	32
x^2	1	4	16	25	16	31	1	4	31	25	25	31	4	1	31	16	25	16	4	1
x^4	1	16	25	31	25	4	1	16	4	31	31	4	16	1	4	25	31	25	16	1
$y = x^e = x^3$	1	8	31	26	13	17	10	19	5	4	29	28	14	23	16	20	7	2	25	32

Working Example V

We can now verify from the table that x^3 is a bijection from \mathbb{Z}_N^* to \mathbb{Z}_N^* (because 3 is relatively prime to $\varphi(N)$)

We recall the following result (stated without proof) from the previous lecture.

Theorem

For any $e \in \mathbb{N}$ such that $\gcd(e, \varphi(N)) = 1$ and $e < \varphi(N)$, the function $x^e: \mathbb{Z}_N^ \rightarrow \mathbb{Z}_N^*$ is a bijection.*

Since x^e is a bijection, we can uniquely define $y^{1/e}$ for any $y \in \mathbb{Z}_N^*$. For example, if $y = 19$ then $y^{1/e} = 13$, where $e = 3$.

The RSA assumption states that, for a random y , finding $y^{1/e}$ is a computationally difficult task!

Working Example VI

Let d be an integer $< \varphi(N)$ such that $e \cdot d = 1 \pmod{N}$. In our case, we have $d = 7$.

Let us calculate a row corresponding to x^7 . We can calculate this by multiplying $x \times x^2 \times x^4$.

x	1	2	4	5	7	8	10	13	14	16	17	19	20	23	25	26	28	29	31	32
x^2	1	4	16	25	16	31	1	4	31	25	25	31	4	1	31	16	25	16	4	1
x^4	1	16	25	31	25	4	1	16	4	31	31	4	16	1	4	25	31	25	16	1
$y = x^e = x^3$	1	8	31	26	13	17	10	19	5	4	29	28	14	23	16	20	7	2	25	32
$x^d = x^7$	1	29	16	14	28	2	10	7	20	25	8	13	26	23	31	5	19	17	4	32

Working Example VII

Note that d is also relatively prime to $\varphi(N)$, and, hence, the mapping x^d is also a bijection.

Working Example VIII

But note that, given d , we can easily compute the e -th root of y .
Check that y^d is identical to $y^{1/e}$.

x	1	2	4	5	7	8	10	13	14	16	17	19	20	23	25	26	28	29	31	32
x^2	1	4	16	25	16	31	1	4	31	25	25	31	4	1	31	16	25	16	4	1
x^4	1	16	25	31	25	4	1	16	4	31	31	4	16	1	4	25	31	25	16	1
$y = x^e = x^3$	1	8	31	26	13	17	10	19	5	4	29	28	14	23	16	20	7	2	25	32
$x^d = x^7$	1	29	16	14	28	2	10	7	20	25	8	13	26	23	31	5	19	17	4	32
$y^d = y^7$	1	2	4	5	7	8	10	13	14	16	17	19	20	23	25	26	28	29	31	32

Quick Summary

- The function $x^e: \mathbb{Z}_N^* \rightarrow \mathbb{Z}_N^*$ is a bijection for all e such that $\gcd(e, \varphi(N)) = 1$
- Given (n, N, e, y) , where $y \stackrel{\$}{\leftarrow} \mathbb{Z}_N^*$, it is difficult for any computationally bounded adversary to compute the e -th root of y , i.e., the element $y^{1/e}$
- But given d such that $e \cdot d = 1 \pmod{\varphi(N)}$, it is easy to compute $y^{1/e}$, because $y^d = y^{1/e}$

Now, think how we can design a key-agreement scheme using these properties. Once the key-agreement protocol is ready, we can use a one-time pad to create an public-key encryption scheme.