

# Lecture 01: Mathematical Basics (Summations)

# What I am Assuming

- I am assuming that you know asymptotic notations. For example, the big-O, little-O notations

- Let us try to write a closed form expression for the following summation

$$S = \sum_{i=1}^n 1$$

- It is trivial to see that  $S = n$

# Summation II

- Now, let us try to write a closed form expression for the following summation

$$S = \sum_{i=1}^n i$$

- We can prove that  $S = \frac{n(n+1)}{2}$ 
  - How do you prove this statement? (Use Induction? Use the formula for the Sum of an Arithmetic Progression?)
- Using Asymptotic Notation, we can say that  $S = \frac{n^2}{2} + o(n^2)$

# Summation III

- Now, let us try to write a closed form expression for the following summation

$$S = \sum_{i=1}^n i^2$$

- We can prove that  $S = \frac{n(n+1)(2n+1)}{6}$ 
  - Why is the expression on the right an integer? (Prove by induction that 6 divides  $n(n+1)(2n+1)$  for all positive integer  $n$ )
  - How do you prove this statement? (Use Induction?)
- Using Asymptotic Notation, we can say that  $S = \frac{n^3}{3} + o(n^3)$


# Summation IV

- Do we see a pattern here?
- Conjecture: For  $k \geq 1$ , we have  $\sum_{i=1}^n i^{k-1} = \frac{n^k}{k} + o(n^k)$ .
  - How do we prove this statement?

# Estimating Summations by Integration I

- Let  $f$  be an increasing function
- For example,  $f(x) = x^{k-1}$  is an increasing function for  $k > 1$  and  $x \geq 0$

# Estimating Summations by Integration II





# Estimating Summations by Integration III

- Observation: “Blue area under the curve” is smaller than the “Shaded area of the rectangle”
  - Blue area under the curve is:

$$\int_{x-1}^x f(t) dt$$

- Shaded area of the rectangle is:

$$f(x)$$

- So, we have the inequality:

$$\int_{x-1}^x f(t) dt \leq f(x)$$

- Summing both side from  $x = 1$  to  $x = n$ , we get

$$\sum_{x=1}^n \int_{x-1}^x f(t) dt \leq \sum_{x=1}^n f(x)$$

# Estimating Summations by Integration IV

- The left-hand side of the inequality is


$$\int_0^1 f(t) dt + \int_1^2 f(t) dt + \cdots + \int_{n-1}^n f(t) dt = \int_0^n f(t) dt$$

- So, for an increasing  $f$ , we have the following lower bound.

$$\int_0^n f(t) dt \leq \sum_{x=1}^n f(x) \quad (1)$$

# Estimating Summations by Integration V

- Now, we will upper bound the summation expression. Consider the figure below



## Estimating Summations by Integration VI

- Observation: “Blue area under the curve” is greater than the “Shaded area of the rectangle”
- So, we have the inequality:

$$\int_{x-1}^x f(t) dt \geq f(x-1)$$

- Now we sum the above inequality from  $x = 2$  to  $x = n + 1$
- We get

$$\int_1^2 f(t) dt + \int_2^3 f(t) dt + \cdots + \int_n^{n+1} f(t) dt \geq f(1) + f(2) + \cdots + f(n)$$

- So, for an increasing  $f$ , we get the following upper bound

$$\int_1^{n+1} f(t) dt \geq \sum_{x=1}^n f(x) \quad (2)$$

# Summary: Estimation of Summation using Integration

## Theorem

For an increasing function  $f$ , we have

$$\int_0^n f(t) dt \leq \sum_{x=1}^n f(x) \leq \int_1^{n+1} f(t) dt$$

Exercise:

- Use this theorem to prove that  $\sum_{i=1}^n i^{k-1} = \frac{n^k}{k} + o(n^k)$ , for  $k \geq 1$
- Consider the function  $f(x) = 1/x$  to find upper and lower bounds for the sum  $H_n = 1 + \frac{1}{2} + \dots + \frac{1}{n}$  using the approach used to prove Theorem 1

# Differentiation and Integration

- Differentiation:  $f'(x)$  represents the slope of the curve  $y = f(x)$  at  $x$
- Integration:  $\int_a^b f(t) dt$  represents the area under the curve  $y = f(x)$  between  $x = a$  and  $x = b$
- Increasing function:
  - Observation: The slope an increasing function is positive
  - So, “ $f$  is increasing at  $x$ ” is equivalent to “ $f'(x) > 0$ ,” i.e.  $f'$  is positive at  $x$
- Suppose we want to mathematically write “Slope of a function  $f$  is increasing”
  - The “slope of a function  $f$ ” is the function “ $f'$ ”
  - So, the statement “slope of a function  $f$  is increasing” is equivalent to “ $(f')' \equiv f''$  is positive”

# Concave Upwards Functions


## Definition (Concave Upwards Function)

A function  $f$  is *concave upwards* in the interval  $[a, b]$  if  $f''$  is positive in the interval  $[a, b]$ .

- Example of functions that concave upwards:  $x^2$ ,  $\exp(x)$ ,  $1/x$  (in the interval  $(0, \infty)$ ),  $x \log x$  (in the interval  $(0, \infty)$ )
  - We emphasize that a “concave upwards” function need not be increasing, for example  $f(x) = 1/x$  (for positive  $x$ ) is decreasing

# Property of Concave Upwards Function I


- Consider the coordinates  $(x - 1, f(x - 1))$  and  $(x, f(x))$
- For a concave upwards function, the secant between the two coordinates is always (on or) above the part of the curve  $f$  between the two coordinates





## Property of Concave Upwards Function II

- So, the shaded area of the trapezium is greater than the blue area under the curve



## Property of Concave Upwards Function III

- So, we get

$$\frac{f(x-1) + f(x)}{2} \geq \int_{x-1}^x f(t) dt$$

- Now, use this new observation to obtain a better lower bound for the sum  $\sum_{x=1}^n f(x)$
- Think: Can you get even tighter bounds?
- Additional Reading: Read on the “trapezoidal rule”

# Estimating Products

- Consider the objective of estimating  $n!$  using elementary functions
- Note that one can convert this estimation of products into estimation of sums by taking log. For example,

$$\ln(n!) = \sum_{i=1}^n \ln(i).$$

- Now, one can tightly upper and lower bound the expression  $\sum_{i=1}^n \ln(i)$ . Use the techniques in the previous slides to obtain meaningful upper and lower bounds of this expression. Suppose

$$L_n \leq \sum_{i=1}^n \ln(i) \leq U_n.$$

- Therefore, one concludes that

$$\exp(L_n) \leq n! \leq \exp(U_n).$$

# Estimating Fractions

- Consider the objective of estimating a fraction  $A_n/B_n$
- Suppose we have  $A_n \leq U_n$  and  $L'_n \leq B_n$ . Note that

$$\frac{1}{B_n} \leq \frac{1}{L'_n}.$$

- Note that multiplying with  $A_n \leq U_n$ , one gets that

$$\frac{A_n}{B_n} \leq \frac{U_n}{L'_n}.$$

- To summarize, upper-bounding a fraction involves upper-bounding the numerator and lower-bounding the denominator
- Analogously, if  $L_n \leq A_n$  and  $B_n \leq U'_n$ , then we get  $\frac{L_n}{B_n} \leq \frac{A_n}{U'_n}$
- **Food for thought.** Provide meaningful upper and lower bound the expression  $\binom{2n}{n} := \frac{(2n)!}{(n!)^2}$ .