

Homework 2

Collaborators :

Practice Questions

These are practice questions. They will **NOT** be graded. We have also provided the final answers. However, it is up to you to understand how or why the given solution is correct.

You do not need to submit these on Gradescope. However, you may find it easier to just include them in the pdf. In that case, please do not mark these questions on Gradescope.

1. **Calculating Large Powers mod p (0 points).** Recall that we learned the repeated squaring algorithm in class. Calculate the following using this concept

$$10^{2025^{2025}+2025} \pmod{101}$$

(Hint: Note that 101 is a prime number, and before applying the repeated squaring algorithm, try to simplify the problem using what you learned in part (a) of Question 1 of the graded homework questions.)

(Note: as can be seen from the latex file, the entirety of $2025^{2025} + 2025$ is in the exponent of 10. Also recall that $a^{b^c} = a^{(b^c)}$, not $(a^b)^c$, which is $a^{b \cdot c}$)

Solution.

100

2. **Order of an Element in $(\mathbb{Z}_p^*, \times)$ (0 points).** The *order* of an element x in the multiplicative group $(\mathbb{Z}_p^*, \times)$ is the smallest positive integer h such that $x^h = 1 \pmod{p}$. For example, the order of 2 in $(\mathbb{Z}_5^*, \times)$ is 4, and the order of 4 in $(\mathbb{Z}_5^*, \times)$ is 2. What is the order of 3 in $(\mathbb{Z}_7^*, \times)$?

Solution.

The order of 3 in $(\mathbb{Z}_7^*, \times)$ is 6.

3. **Elliptic Curve (0 points).** In class, we have briefly discussed elliptic curves. Here we will see some concrete examples of elliptic curves on finite prime fields.

Let $Y^2 = X^3 + X$ be an elliptic curve over the field $(\mathbb{F}_{23}, +, \cdot)$. A point (X, Y) lies on the elliptic curve if it satisfies the equation $Y^2 = X^3 + X$.

- (a) (0 points) Are the two points $P = (9, 18)$ and $Q = (11, 10)$ on the curve?

Solution.

Yes, they are.

- (b) (0 points) Find the third point R (that is, the intersection point other than P and Q) where the line connecting P and Q intersects the elliptic curve $Y^2 = X^3 + X$. For $R = (x, y)$, define the “inverse of R ” to be the point $S = (x, -y)$. Find the inverse of point R . Recall from the lecture that “ $P + Q$ ” is defined to be the point $S :=$ “inverse of R .”

Solution.

The inverse of R is $S = (19, 22)$.

4. Special Group: Quotient Group $(\mathbb{Z}_p^*/\{-1, +1\}, \times)$ (0 points).

Throughout this problem, p is an odd prime.

We will define the group $(\mathbb{Z}_p^*/\{-1, +1\}, \times)$ using the quotient group and coset notation.

In general, given a group G and a subgroup H , G/H means the set of cosets formed by H in G . The cosets of H in G are $gH := \{g \times h : h \in H\}$ for $g \in G$. When G is abelian, as $\mathbb{Z}_p^*$ is, this set of cosets is itself a group under the operation inherited from G ; this is why G/H is called a quotient group.

Now, consider the elements in $\mathbb{Z}_p^*$, which are

$$\left\{1, 2, 3, \dots, \frac{p-1}{2} - 1, \frac{p-1}{2}, \frac{p-1}{2} + 1, \frac{p-1}{2} + 2, \dots, p-3, p-2, p-1\right\}.$$

The cosets $\mathbb{Z}_p^*/\{-1, +1\}$ are

$$\left\{1 \times \{-1, +1\}, 2 \times \{-1, +1\}, 3 \times \{-1, +1\}, \dots, \frac{p-1}{2} \times \{-1, +1\}\right\}.$$

Observe that $1 \times \{-1, +1\}$ contains both 1 and $-1 = p-1$. Similarly, $2 \times \{-1, +1\}$ contains both 2 and $p-2$. Finally, $\frac{p-1}{2} \times \{-1, +1\}$ contains both $\frac{p-1}{2}$ and $\frac{-p+1}{2} = p + \frac{-p+1}{2} = \frac{p-1}{2} + 1$. The final set of elements in $\mathbb{Z}_p^*/\{-1, +1\}$ are $\left\{1, 2, \dots, \frac{p-1}{2} - 1, \frac{p-1}{2}\right\}$.

Intuitively, you can think of it as follows: for any $x \in \mathbb{Z}_p^*$, both x and $-x \in \mathbb{Z}_p^*$ represent the same element in $\mathbb{Z}_p^*/\{-1, +1\}$. The same way that the integers x and $x + p$ represent the same element in $\mathbb{Z}_p$. For this reason, $\mathbb{Z}_p$ can also be written as $\mathbb{Z}/p\mathbb{Z}$, the quotient of the group $(\mathbb{Z}, +)$ by its subgroup $p\mathbb{Z}$.

Let us look at a concrete example: $(\mathbb{Z}_{11}^*/\{-1, +1\}, \times)$. The elements in $\mathbb{Z}_{11}^*$ are $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$. The elements in $\mathbb{Z}_{11}^*/\{-1, +1\}$ are $\{1, 2, 3, 4, 5\}$.

$\times$ is multiplication in $\mathbb{Z}_p^*$, followed by replacing the product with whichever of it and its negation lies in $\left\{1, 2, \dots, \frac{p-1}{2}\right\}$. For example, $2 \times 3 = 6 = -5 = 5$.

Answer the following questions in $(\mathbb{Z}_{11}^*/\{-1, +1\}, \times)$:

- (a) (0 points) What is 4×5 ?

Solution.

$$4 \times 5 = 2.$$

- (b) (0 points) What is 3×4 ?

Solution.

$$3 \times 4 = 1, \text{ the identity of the group.}$$

Homework Questions

These are homework questions and will be graded. Please make sure to clearly mark each problem on Gradescope.

1. **Some properties of $(\mathbb{Z}_p^*, \times)$ (5+5 points).** Recall that $\mathbb{Z}_p^*$ is the set $\{1, \dots, p-1\}$ and $\times$ is integer multiplication mod p , where p is a prime. For example, if $p = 5$, then 2×3 is 1. In this problem, we shall prove that $(\mathbb{Z}_p^*, \times)$ is a group when p is any prime. The only part missing in the lecture was the proof that every $x \in \mathbb{Z}_p^*$ has an inverse. We will find the inverse of any element $x \in \mathbb{Z}_p^*$.

(a) (5 points) For $x \in \mathbb{Z}_p^*$, prove that the inverse of $x \in \mathbb{Z}_p^*$ is given by

$$\overbrace{x \times x \times \dots \times x}^{(p-2)\text{-times}}$$

That is, prove that $x^{p-1} = 1 \pmod p$, for any prime p and $x \in \mathbb{Z}_p^*$. (Hint: Use Homework 1, Problem 5(b), which states that $x^p \pmod p = x$ for any prime p and $x \in \{1, 2, \dots, p-1\}$.)

Solution.

- (b) (5 points) Let p , and q be two distinct primes. Prove that $(p^q - p) + (q^p - q)$ is divisible by pq .

Solution.

2. **Understanding Groups: Part One (6+8+5+6+5 points).** Recall that when we defined a group $(G, \circ)$, we stated that there exists an element e such that for all $x \in G$ we have $x \circ e = x$. Note that e is “applied on x from the right.” Similarly, for every $x \in G$, we are guaranteed that there exists $\text{inv}(x) \in G$ such that $x \circ \text{inv}(x) = e$. Note that $\text{inv}(x)$ is again “applied to x from the right.”

In this problem, however, we shall explore the following questions: (a) Is there an “identity from the left?” and (b) Is there an “inverse from the left?”

We shall formalize and prove these results in this question.

- (a) (6 points) Prove that it is impossible that there exists $a, b, c \in G$ such that $a \neq b$ but $a \circ c = b \circ c$.

Solution.

- (b) (8 points) Prove that $e \circ x = x$, for all $x \in G$.

Solution.

- (c) (5 points) Prove that if there exists an element $\alpha \in G$ such that for **some** $x \in G$, we have $\alpha \circ x = x$, then $\alpha = e$. (Remark: Note that these two steps prove that the “left identity” is identical to the right identity e .)

Solution.

- (d) (6 points) Prove that $\text{inv}(x) \circ x = e$, for all $x \in G$.

Solution.

- (e) (5 points) Prove that if there exists an element $\alpha \in G$ and $x \in G$ such that $\alpha \circ x = e$, then $\alpha = \text{inv}(x)$.

(Remark: Note that these two steps prove that the “left inverse of x ” is identical to the right inverse $\text{inv}(x)$.)

Solution.

3. **Understanding Groups: Part Two (9+6 points).** In this part, we will prove that distinct elements of a group have distinct inverses. And finally, using the results of Question 2, we will prove a crucial result for the security of the one-time pad over the group $(G, \circ)$.

- (a) (9 points) Suppose $a, b \in G$. Let $\text{inv}(a)$ and $\text{inv}(b)$ be the inverses of a and b , respectively (i.e., $a \circ \text{inv}(a) = e$ and $b \circ \text{inv}(b) = e$). Prove that $\text{inv}(a) = \text{inv}(b)$ if and only if $a = b$.

Solution.

- (b) (6 points) Suppose $m \in G$ is a message and $c \in G$ is a ciphertext. Prove that there exists a unique $\mathbf{sk} \in G$ such that $m \circ \mathbf{sk} = c$.

Solution.

4. **Order of an Element in $(\mathbb{Z}_p^*, \times)$ (10+5+10+5 points).** Throughout this problem, p is a prime. The *order* of an element x in the multiplicative group $(\mathbb{Z}_p^*, \times)$ is the smallest positive integer h such that $x^h = 1 \pmod p$. For example, the order of 2 in $(\mathbb{Z}_5^*, \times)$ is 4, and the order of 4 in $(\mathbb{Z}_5^*, \times)$ is 2.

- (a) (10 points) Let x be an element in $(\mathbb{Z}_p^*, \times)$ such that $x^n = 1 \pmod p$ for some positive integer n and let h be the order of x in $(\mathbb{Z}_p^*, \times)$. Prove that h divides n .

Solution.

- (b) (5 points) Let $x \in \mathbb{Z}_p^*$ and let h be the order of x in $(\mathbb{Z}_p^*, \times)$. Prove that h divides $(p - 1)$.

Solution.

- (c) (10 points) Let $x \in \mathbb{Z}_p^*$, let h be the order of x in $(\mathbb{Z}_p^*, \times)$, and let k be a positive integer. Let r denote the order of $y = x^k \bmod p \in \mathbb{Z}_p^*$. Show that $r = \frac{h}{d} \in \mathbb{Z}$ where d denotes the greatest common divisor of h and k .

Hint: Use part (a) and prove that r divides $\frac{h}{d}$ and $\frac{h}{d}$ divides r .

Solution.

- (d) (5 points) Let $p > 2$ be a prime, and $a \in (\mathbb{Z}_p^*, \times)$. Then, show that the element $b = a^{\frac{p-1}{2}} \in (\mathbb{Z}_p^*, \times)$ is equal to 1 or has order 2.

Solution.

5. **Defining Multiplication over $\mathbb{Z}_{27}^*$ (5+10+10 points).** In the class, we had considered the group $(\mathbb{Z}_{26}, +)$ to construct a one-time pad for a one-alphabet message. Can we define a group with 26 elements using a “multiplication”-like operation? This problem will help you define the $(\mathbb{Z}_{27}^*, \times)$ group, which has 26 elements.

The first attempt from class. Recall that in class we saw that the following is also a group.

$$(\mathbb{Z}_{27} \setminus \{0, 3, 6, 9, 12, 15, 18, 21, 24\}, \times),$$

where $\times$ is integer multiplication mod 27. However, the set had only 18 elements.

In this problem, we shall define $(\mathbb{Z}_{27}^*, \times)$ in a different manner such that the set has 26 elements.

A new approach. Interpret $\mathbb{Z}_{27}^*$ as the set of all triplets (a_0, a_1, a_2) such that $a_0, a_1, a_2 \in \mathbb{Z}_3$ and at least one of them is non-zero. Intuitively, you can think of the triplets as the ternary representation of the elements in $\mathbb{Z}_{27}^*$. We interpret the triplet (a_0, a_1, a_2) as the polynomial $a_0 + a_1X + a_2X^2$. So, every element in $\mathbb{Z}_{27}^*$ has an associated non-zero polynomial of degree at most 2, and every non-zero polynomial of degree at most 2 has an element in $\mathbb{Z}_{27}^*$ associated with it.

The multiplication ($\times$ operator) of the element (a_0, a_1, a_2) with the element (b_0, b_1, b_2) is defined as the element corresponding to the polynomial

$$(a_0 + a_1X + a_2X^2) \times (b_0 + b_1X + b_2X^2) \mod 2 + 2X + X^3$$

The multiplication ($\times$ operator) of the element (a_0, a_1, a_2) with the element (b_0, b_1, b_2) is defined as follows.

Input (a_0, a_1, a_2) and (b_0, b_1, b_2) .

- i. Define $A(X) := a_0 + a_1X + a_2X^2$ and $B(X) := b_0 + b_1X + b_2X^2$
- ii. Compute $C(X) := A(X) \times B(X)$ (interpret this step as “multiplication of polynomials with integer coefficients”)
- iii. Compute $R(X) := C(X) \mod 2 + 2X + X^3$ (interpret this as step as taking a remainder where one treats both polynomials as polynomials with integer coefficients). Let $R(X) = r_0 + r_1X + r_2X^2$
- iv. Return $(c_0, c_1, c_2) = (r_0 \mod 3, r_1 \mod 3, r_2 \mod 3)$

For example, the multiplication $(0, 1, 1) \times (1, 1, 2)$ is computed in the following way.

- i. $A(X) = X + X^2$ and $B(X) = 1 + X + 2X^2$.
- ii. $C(X) = X + 2X^2 + 3X^3 + 2X^4$.
- iii. $R(X) = -6 - 9X - 2X^2$.

iv. $(c_0, c_1, c_2) = (0, 0, 1)$.

According to [this definition](#) of the $\times$ operator, solve the following problems.

(a) (5 points) Evaluate $(0, 2, 2) \times (2, 1, 2)$.

Solution.

- (b) (10 points) Note that $e = (1, 0, 0)$ is an identity element. Find the inverse of $(1, 0, 2)$.

Solution.

- (c) (10 points) Assume that $(\mathbb{Z}_{27}^*, \times)$ is a group. Find the order of the element $(2, 1, 0)$.

(Recall that, in a group $(G, \circ)$, the order of an element $x \in G$ is the smallest positive integer h such that $\overbrace{x \circ x \circ \cdots \circ x}^{h\text{-times}} = e$)

Solution.

6. **Special Group: Elliptic Curve Group (1+2+4+3+1+2 points).** Throughout this problem, K is a field whose characteristic is neither 2 nor 3, and $a, b \in K$ satisfy $4a^3 + 27b^2 \neq 0$. An elliptic curve over K is the graph E of an equation $Y^2 = X^3 + aX + b$ where $X, Y \in K$. The condition $4a^3 + 27b^2 \neq 0$ says that the cubic $X^3 + aX + b$ has no repeated root; without it the graph has a singular point, at which the addition rule below breaks down. The elliptic curve group is the set $E \cup \{\infty\}$, where ∞ is one extra point, the point at infinity, which serves as the identity of the group. Note that the point ∞ is not a point on the graph $Y^2 = X^3 + aX + b$.

(a) **Addition Rule.** Let $P = (x, y)$ be a point on the elliptic curve E : $Y^2 = X^3 + aX + b$, and define $-P := (x, -y)$, the reflection of P in the x -axis. Let $P = (x_p, y_p)$ and $Q = (x_q, y_q)$ be two points on E . If $x_p \neq x_q$, so that the two points are not on the same vertical line, define $P + Q := -R$, where R is the third point at which the straight line through P and Q meets E . If $x_p = x_q$ and $y_p \neq y_q$, so that the two points are distinct and lie on the same vertical line, then $y_q = -y_p$ and $Q = -P$; in this case, define $P + Q := \infty$, the identity of the elliptic curve group. The one remaining case, $P = Q$, is handled by the tangent line rule in part (iii) below. Moreover, define $P + \infty = \infty + P = P$ for every $P \in E \cup \{\infty\}$.

- i. (1 point) Consider E : $Y^2 = X^3 - 4X - 5$ over $\mathbb{F}_{17}$. Let $P = (4, 3)$ and $Q = (9, 12)$. Check that P and Q are points on E .

Solution.

- ii. (2 points) Compute $P + Q$.

Solution.

- iii. (4 points) To add a point $P \neq \infty$ to itself, draw a tangent line on the elliptic curve E at point P . If the line is vertical, then $P + P = \infty$. If the tangent line is not vertical, then it intersects E at exactly one more point. For $P = (4, 3)$, $Q = (9, 12)$ and $-R = P + Q$, compute $P + P$, $Q + Q$ and $R + R$.

Solution.

(b) **Deriving the addition formulas.** Consider elliptic curve $E: Y^2 = X^3 + aX + b$ over field K . Let $P = (x_p, y_p), Q = (x_q, y_q)$ be two points on E . Let $-R = P + Q = (x_r, -y_r)$. The formulas below, together with the case analysis of part (a), show that $+$ is a well defined operation on $E \cup \{\infty\}$, that ∞ is its identity, and that $-P$ is the inverse of P . Associativity holds as well, but proving it is a longer computation that we do not carry out here. We will show the following.

- i. (3 points) Let $Y = mX + t$ be the line intersecting P and Q . Assume $x_p \neq x_q$. Let $m = \frac{y_p - y_q}{x_p - x_q}$ be the slope of the line. Show that $x_r = m^2 - x_p - x_q$ and $y_r = y_p - m(x_p - x_r)$.
(Hint: Recall Vieta's formulas for a degree 3 polynomial $a_3X^3 + a_2X^2 + a_1X + a_0 = 0$ over a field K , where $a_0, a_1, a_2, a_3 \in K$ and $a_3 \neq 0$. Let $r_1, r_2, r_3 \in K$ be the roots of the polynomial. The following equalities hold.

$$\begin{cases} r_1 + r_2 + r_3 = -\frac{a_2}{a_3} \\ r_1r_2 + r_2r_3 + r_1r_3 = \frac{a_1}{a_3} \\ r_1r_2r_3 = -\frac{a_0}{a_3} \end{cases} .$$

)

Solution.

- ii. (1 point) Show that if $x_p = x_q$ and $y_p = -y_q$, then $P + Q = \infty$.

Solution.

- iii. (2 points) Show that if $x_p = x_q$ and $y_p = y_q \neq 0$, then $x_r = m^2 - 2x_p$ and $y_r = y_p - m(x_p - x_r)$ with $m = \frac{3x_p^2 + a}{2y_p}$.

Solution.