

Solution of Homework 1: *Basic Logic*

Q.1 Make truth tables for the following statement:

- $p \vee (\overline{r \vee q})$;

Answer

p	q	r	$r \vee q$	$\overline{r \vee q}$	$p \vee (\overline{r \vee q})$
T	T	T	T	F	T
T	T	F	T	F	T
T	F	T	T	F	T
T	F	F	F	T	T
F	T	T	T	F	F
F	T	F	T	F	F
F	F	T	T	F	F
F	F	F	F	T	T

- $(p \wedge \neg q) \rightarrow r$.

Answer

p	q	r	$\neg q$	$p \wedge \neg q$	$(p \wedge \neg q) \rightarrow r$
T	T	T	F	F	T
T	T	F	F	F	T
T	F	T	T	T	T
T	F	F	T	T	F
F	T	T	F	F	T
F	T	F	F	F	T
F	F	T	T	F	T
F	F	F	T	F	T

Q.2 Using *logical equivalences* discussed in class prove that

$$(p \wedge q) \rightarrow (p \vee q)$$

is a tautology, that is, prove that

$$(p \wedge q) \rightarrow (p \vee q) \equiv T.$$

Answer

$$\begin{aligned}
 (p \wedge q) \rightarrow (p \vee q) &\equiv \neg(p \wedge q) \vee (p \vee q) \\
 &\equiv (\neg p \vee \neg q) \vee (p \vee q) \\
 &\equiv (\neg p \vee p) \vee (\neg q \vee q) \\
 &\equiv T \vee T \\
 &\equiv T
 \end{aligned}$$

Note: Another way to solve this question is by constructing the truth table for the given logical expression and showing that it always yields T for all values of p and q .

Q.3 Let

$P(x, y) : x + y \geq 5$ where x, y are positive integers.

Tell whether the following statements are true or false:

- $\forall_x \forall_y P(x, y)$
- $\forall_x \exists_y P(x, y)$.

Answer

- $\forall_x \forall_y P(x, y)$: **“False”**
Counterexample: $P(1, 2) : 1 + 2 \not\geq 5$.
- $\forall_x \exists_y P(x, y)$: **“True”**
If we pick an arbitrary value for x , say a , then there always exists a value for y (for example, $a + 5$) such that $x + y = 2a + 5 \geq 5$.

Q.4 Which of the following is equivalent to $\overline{\forall_x \exists_y P(x, y)} \equiv \neg \forall_x \exists_y P(x, y)$:

- (a) $\overline{\exists_x \forall_y P(x, y)}$;
- (b) $\overline{\forall_x \exists_y P(x, y)}$;
- (c) $\overline{\exists_x \forall_y \overline{P(x, y)}}$;
- (d) $\overline{\exists_x \exists_y \overline{P(x, y)}}$.

Answer

- (c) $\overline{\exists_x \forall_y \overline{P(x, y)}}$.