

# Robust Online Classification: From Estimation to Denoising

Changlong Wu

Ananth Grama

Wojciech Szpankowski

CSol, Purdue University

WUCHANGL@HAWAII.EDU

AYG@CS.PURDUE.EDU

SZPAN@PURDUE.EDU

## Abstract

We study online classification in the presence of noisy labels. The noise mechanism is modeled by a general *kernel* that specifies, for any feature-label pair, a (known) *set* of distributions over noisy labels. At each time step, an *adversary* selects an *unknown* distribution from the distribution set specified by the kernel based on the actual feature-label pair, and generates the noisy label from the selected distribution. The learner then makes a prediction based on the actual features and *noisy* labels observed thus far, and incurs loss 1 if the prediction differs from the underlying truth (and 0 otherwise). The prediction quality is quantified through *minimax risk*, which computes the cumulative loss over a finite horizon  $T$ . We show that for a wide range of natural noise kernels, adversarially selected features, and finite class of labeling functions, minimax risk can be upper bounded *independent* of the time horizon and logarithmic in the size of labeling function class. We then extend these results to infinite classes and stochastically generated features via the concept of stochastic sequential covering. Our results extend and encompass findings of Ben-David et al. (2009) through substantial generality, and provide *intuitive* understanding through a novel reduction to *online conditional distribution estimation*.

**Keywords:** Online classification, noisy label, sequential probability assignment, mixable losses

## 1. Introduction

Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be a class of functions mapping the features (instances)  $\mathcal{X}$  to the labels  $\mathcal{Y}$ , where  $|\mathcal{Y}| = N$  is a finite set of size  $N$ . The *realizable* online classification problem is defined as a game with the following protocol: at the beginning, Nature selects  $h \in \mathcal{H}$ ; at each time step  $t$ , nature chooses  $\mathbf{x}_t \in \mathcal{X}$  and reveals to a *learner*; the learner then makes a prediction  $\hat{y}_t \in \mathcal{Y}$ , potentially using the history observed thus far; Nature then reveals the true label  $y_t = h(\mathbf{x}_t)$  and the learner incurs the loss  $1\{\hat{y}_t \neq y_t\}$ . The goal is to minimize the cumulative loss for a given horizon  $T$ . It was shown in the seminal work of Littlestone (1988) that such cumulative loss can be completely characterized by the Littlestone dimension  $\text{Ldim}(\mathcal{H})$  of  $\mathcal{H}$  if the features  $\mathbf{x}_t$  are selected adversarially. Attempts to generalize this *realizable* case primarily focus on the *regret* formulation (Ben-David et al., 2009; Daniely et al., 2015; Rakhlin et al., 2010), and assume the *observable* labels  $\tilde{y}^T = \{\tilde{y}_1, \dots, \tilde{y}_T\}$  to be generated *adversarially*. Here, we do not evaluate the *actual* cumulative loss for the underlying truth, rather, we evaluate the prediction quality through the following regret:

$$\sum_{t=1}^T 1\{\tilde{y}_t \neq \hat{y}_t\} - \inf_{h \in \mathcal{H}} \sum_{t=1}^T 1\{h(\mathbf{x}_t) \neq \tilde{y}_t\}.$$

The *expected* regret will then be upper bounded by  $O(\sqrt{T \text{Ldim}(\mathcal{H}) \log(TN)})$  Daniely et al. (2015).

While the sub-linear regret bounds for the adversarially generated observable labels are intriguing from a mathematical point of view, they do not necessarily reveal the *actual* cumulative errors

a learner might incur, especially when there are *correlations* between the observable labels and the underlying truth. Indeed, a notable result by Ben-David et al. (2009) showed that if the hypotheses are binary valued and the *observable* (noisy) labels  $\tilde{y}_t$  are generated by flipping the *true* label  $y_t = h(\mathbf{x}_t)$  w.p.  $\eta_t \leq \eta < \frac{1}{2}$  (and w.p.  $1 - \eta_t$  remain unchanged), then one can achieve the *actual* expected cumulative loss of order  $O(\log |\mathcal{H}|/(2\eta - 1)^2)$ . Surprisingly, the best approximating error of hypotheses in  $\mathcal{H}$  to noisy labels is of order  $\eta T$ ; the *actual* cumulative error is independent of the time horizon  $T$ ! As remarkable as this result is, its proof, as provided in Ben-David et al. (2009), is based on a somewhat non-intuitive *backward induction*.

This paper generalizes the noisy-label scenario of Ben-David et al. (2009) to broader *stochastic* noisy models. Our approach not only provides results that go beyond Ben-David et al. (2009) through substantial generality, but also provide an *intuitive* understanding of the underlying paradigm. Let  $\tilde{\mathcal{Y}}$  be the set of noisy observations, which we assume is finite (this assumption can be relaxed) and of size  $M$ . We model the noise process as a *noisy kernel*  $\mathcal{K} : \mathcal{X} \times \mathcal{Y} \rightarrow 2^{\mathcal{D}(\tilde{\mathcal{Y}})}$ , where  $\mathcal{D}(\tilde{\mathcal{Y}})$  is the set of all distributions over  $\tilde{\mathcal{Y}}$ . That is, the kernel  $\mathcal{K}$  maps each pair  $(\mathbf{x}, y)$  to a *subset*  $\mathcal{Q}_y^{\mathbf{x}} = \mathcal{K}(\mathbf{x}, y) \subset \mathcal{D}(\tilde{\mathcal{Y}})$  of distributions over  $\tilde{\mathcal{Y}}$ . We consider the following *robust (noisy) online classification scenario*: at the beginning, *Nature* selects  $h \in \mathcal{H}$ ; at each time step  $t$ , *Nature* chooses  $\mathbf{x}_t \in \mathcal{X}$  and reveals to the learner; the learner then makes a prediction  $\hat{y}_t$ , based the feature  $\mathbf{x}^t$  and *noisy* labels  $\tilde{y}^{t-1}$ ; An *adversary* then selects a distribution  $\tilde{p}_t \in \mathcal{Q}_{h(\mathbf{x}_t)}^{\mathbf{x}_t}$ , samples  $\tilde{y}_t \sim \tilde{p}_t$  and reveals  $\tilde{y}_t$  to the learner. Let  $\Phi$  and  $\Psi$  be the strategies of the learner and adversary, respectively. The goal of the learner is to minimize the following expected minimax risk:

$$\tilde{r}_T(\mathcal{H}, \mathcal{K}) = \inf_{\Phi} \sup_{h \in \mathcal{H}, \mathbf{x}^T \in \mathcal{X}^T, \Psi} \mathbb{E} \left[ \sum_{t=1}^T 1\{h(\mathbf{x}_t) \neq \hat{y}_t\} \right], \quad (1)$$

where  $\hat{y}_t = \Phi(\mathbf{x}^t, \tilde{y}^{t-1})$ ,  $\tilde{y}_t \sim \tilde{p}_t$  and  $\tilde{p}_t = \Psi(\mathbf{x}^t, \hat{y}^t) \in \mathcal{Q}_{h(\mathbf{x}_t)}^{\mathbf{x}_t}$ . In addition, we also consider scenarios that hold with *high probability* and cases for which the features are generated from *general stochastic* process. We refer to Section 2 for a more complete specification of our setting.

### 1.1. Results and Techniques

Our main results in this paper establish tight minimax risk upper bounds for the *robust online classification problem* for a wide range of hypotheses classes  $\mathcal{H}$ , noisy kernel  $\mathcal{K}$ , as well as the feature generating mechanisms, both in expectation and with high probability. The main technique for achieving these results is a novel reduction to the *online conditional distribution estimation* problem under exponentially concave losses, see Section 2.1.

At a high level, our approach for online conditional distribution estimation is to find the *noisy* distribution  $\tilde{p}_t$  at each time step  $t$ . Suppose we are able to do it, then the prediction of the true labels will be simply estimated as the  $\hat{y}_t \in \mathcal{Y}$  such that  $\tilde{p}_t \in \mathcal{Q}_{\hat{y}_t}^{\mathbf{x}_t}$ , where  $\mathcal{Q}_{\hat{y}_t}^{\mathbf{x}_t} = \mathcal{K}(\mathbf{x}_t, \hat{y}_t) \subset \mathcal{D}(\tilde{\mathcal{Y}})$ . Under mild assumptions, e.g., when  $\mathcal{Q}_y^{\mathbf{x}_t}$  and  $\mathcal{Q}_{y'}^{\mathbf{x}_t}$  are *well-separated* for any  $y \neq y' \in \mathcal{Y}$ , the prediction quality of  $\hat{y}_t$  will then be bounded, provided our estimation on the distribution  $\tilde{p}_t$  is accurate.

The main challenge here is that, the distribution  $\tilde{p}_t$  is selected *arbitrarily* (even adversarially) from a *set* of distributions  $\mathcal{Q}_{h(\mathbf{x}_t)}^{\mathbf{x}_t}$ , which is *unknown* a-priori to the learner. Our main idea for resolving this issue is to estimate  $\tilde{p}_t$  in a more *structural* way. More precisely, assume  $\mathcal{Y} = \{0, 1\}$ , we will construct for any  $h \in \mathcal{H}$  a *distribution-valued* function  $f_h$ , such that  $\forall \mathbf{x} \in \mathcal{X}$ ,  $f_h(\mathbf{x}) = q_{h(\mathbf{x})}^{\mathbf{x}}$ , where  $\forall y \in \mathcal{Y}$ ,  $q_y^{\mathbf{x}} \in \mathcal{Q}_y^{\mathbf{x}}$  are distributions satisfying  $\|q_0^{\mathbf{x}} - q_1^{\mathbf{x}}\|_2^2 = \inf_{p \in \mathcal{Q}_0^{\mathbf{x}}, q \in \mathcal{Q}_1^{\mathbf{x}}} \{\|p - q\|_2^2\}$ .

We then show that there exist predictors  $\hat{p}_t = \Phi(\mathbf{x}_t, \tilde{y}^{t-1}) \in \mathcal{D}(\tilde{\mathcal{Y}})$  such that

$$\sup_{h \in \mathcal{H}} \mathbb{E} \left[ \sum_{t=1}^T L^2(\tilde{p}_t, \hat{p}_t) - L^2(\tilde{p}_t, f_h(\mathbf{x}_t)) \right] \leq O(\log |\mathcal{H}|), \quad (2)$$

where  $L^2(p, q) = \|p - q\|_2^2$ . This is satisfied by, for example, the *Exponential Weighted Average* (EWA) predictor with exp-concave losses; see Lemma 9. The *classifier* is now given by

$$\hat{y}_t = \arg \min_y \{L(q_y^{\mathbf{x}_t}, \hat{p}_t) : y \in \mathcal{Y}\}.$$

Using a *geometric* property of  $L^2$  divergence (see Lemma 13), we then arrive at our first main result (Theorem 12):

**Theorem 1 (Informal)** *Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be any finite class with  $|\mathcal{Y}| = 2$ ,  $\mathcal{K}$  be any noisy kernel that satisfies  $\forall \mathbf{x} \in \mathcal{X}, y \neq y' \in \mathcal{Y}, L^2(\mathcal{Q}_y^{\mathbf{x}}, \mathcal{Q}_{y'}^{\mathbf{x}}) \stackrel{\text{def}}{=} \inf_{p \in \mathcal{Q}_y^{\mathbf{x}}, q \in \mathcal{Q}_{y'}^{\mathbf{x}}} \{\|p - q\|_2^2\} \geq \gamma > 0$  and  $\mathcal{Q}_y^{\mathbf{x}} = \mathcal{K}(\mathbf{x}, y) \subset \mathcal{D}(\tilde{\mathcal{Y}})$  is closed and convex. Then  $\tilde{r}_T(\mathcal{H}, \mathcal{K}) \leq \frac{16 \log |\mathcal{H}|}{\gamma}$ .*

This result recovers (Ben-David et al., 2009, Theorem 15) up to a constant factor. However, our result holds for *any* noisy kernel whenever it exhibits a bounded gap under  $L^2$  divergence. We further extend this argument in Section 3.3 to special noisy kernels with improved bounds. This includes Theorem 21, which extends (Ben-David et al., 2009, Theorem 15) to the *multi-label* case and with *high probability* minimax risk of form  $O\left(\frac{\log |\mathcal{H}| + \log(1/\delta)}{\gamma}\right)$ , and in Theorem 23 we describe an extension to the *Tsybakov* type noise.

Theorem 1, while widely applicable, does not extend directly to the *multi-label* case for a *general* noisy kernel, since for the multi-label case there is no *canonical* way of defining distribution-valued functions  $f_h$ . Our main idea for resolving this issue is a reduction to *pairwise comparison* of two-hypotheses. This leads to our second main result (Theorem 17 & 18):

**Theorem 2 (Informal)** *Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be a finite class of size  $K$  and  $\mathcal{K}$  be any noisy kernel. If for any  $h_i, h_j \in \mathcal{H}$ , the sub-class  $\{h_i, h_j\}$  admits a predictor that achieves a cumulative error of  $C(\delta/2K)$  w.p.  $\geq 1 - \delta/2K$ , then the class  $\mathcal{H}$  admits a predictor that achieves a cumulative error of  $O(C(\delta/2K) \log K + \log(2/\delta))$  w.p.  $\geq 1 - \delta$ . Specializing to the kernels in Theorem 1, we achieve high probability minimax risk of form  $O\left(\frac{\log^2 K + \log K \log(1/\delta\gamma)}{\gamma^2}\right)$ .*

Note that Theorem 2 holds with *high probability* and with *exponentially* decaying error probability. Moreover, the error bound is *independent* to the size of  $\mathcal{Y}$  and  $\tilde{\mathcal{Y}}$ , as well as the time horizon  $T$ .

Our next main result (Theorem 26) establishes tight minimax risk for *infinite* class  $\mathcal{H}$  and stochastically generated features from a *general random process*, through the concept of *stochastic sequential covering* introduced in Wu et al. (2022a, 2023a) (see also Ben-David et al. (2009)).

**Theorem 3 (Informal)** *Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be any (infinite) class and  $\mathcal{K}$  be a kernel satisfying conditions in Theorem 1. Suppose  $\mathcal{H}$  admits a finite stochastic sequential covering  $\mathcal{G}$  w.r.t. a random process class  $\mathbf{P}$  over  $\mathcal{X}^T$ . Then, the high probability minimax risk is upper bounded by  $O\left(\frac{\log^2 |\mathcal{G}| + \log |\mathcal{G}| \log(1/(\delta\gamma))}{\gamma^2}\right)$ . In particular, if  $\text{Ldim}(\mathcal{H}) < \infty$ ,  $|\mathcal{Y}| = N$  and  $\mathbf{x}^T$  are generated adversarially, we have error bound  $O\left(\frac{\text{Ldim}(\mathcal{H})^2 \log^2(TN/\delta\gamma)}{\gamma^2}\right)$ .*

We refer to Section 3.4 for more results with general class  $\mathcal{H}$  and random process class  $\mathbf{P}$ .

**Summary of contributions.** In summary, our main contributions are as follows: (i) we introduce a novel formulation of *robust online classification* that goes beyond classical adversarial models to include broader *stochastic noisy* scenarios and with tighter guarantees; (ii) by leveraging classical results of prediction under *exp-concave* losses, we show that for a wide range of noisy scenarios, the classification problem can be effectively reduced to the *estimation* of the *noisy distributions*, even if such distributions are *unknown* a priori, leading to tight minimax upper bounds that extend and subsume prior known results; (iii) for noisy scenarios where such a reduction is not available, we provide a novel reduction to the *pairwise comparison* of two hypotheses, which provides nearly tight upper bounds for *general* models and may be of independent interest; (iv) finally, we prove tight minimax upper bounds for a wide range of (possibly infinite) hypothesis classes and feature-generating processes through the concept of stochastic sequential cover.

## 1.2. Related Work

Our approach is conceptually similar to the *estimation to decision* framework of Foster et al. (2021), where the authors use online conditional probability estimation in the context of *online decision making*. However, a distinguishing feature of our work is that our conditional probability estimation problem is necessarily *miss-specified*, since our noisy distributions are selected *adversarially* and unknown a-priori to the learner. We are able to solve this problem by using the  $L^2$  divergence (instead of Hellinger distance as in Foster et al. (2021)) since it allows us to use *geometric* properties of  $L^2$  divergence to handle miss-specified noisy distributions. Our problem setup is also related to the *differentially private* conditional distribution learning as in Wu et al. (2023b) and *robust hypothesis testing* discussed in (Polyanskiy and Wu, 2022, Chapter 16). Online conditional probability estimation has been extensively studied in literature, see Rakhlin and Sridharan (2015); Bilodeau et al. (2020); Bhatt and Kim (2021); Bilodeau et al. (2021); Wu et al. (2022b,a). Conditional density estimation in the *batch* setting has also been extensively studied, see Grünwald and Mehta (2020) for KL-divergence with miss-specification and Efromovich (2007) for  $L^2$  loss. Learning from noisy labels in the *batch* case was discussed in Natarajan et al. (2013) (see also the references therein) by leveraging suitably defined proxy loss. There has been a long line of research on online prediction with *adversarial* observable labels in the *regret* formulation, see Cesa-Bianchi and Lugosi (2006); Ben-David et al. (2009); Rakhlin et al. (2010); Daniely et al. (2015).

## 2. Problem Formulation and Preliminaries

Let  $\mathcal{X}$  be a set of features (or instances),  $\mathcal{Y}$  be a set of labels, and  $\tilde{\mathcal{Y}}$  be a set of *noisy observations*. We assume throughout the paper that  $|\mathcal{Y}| = N$  and  $|\tilde{\mathcal{Y}}| = M$  for some integers  $N, M \geq 2$ . We denote  $\mathcal{D}(\tilde{\mathcal{Y}}) = \left\{ p = (p[1], \dots, p[M]) \in [0, 1]^M : \sum_{m=1}^M p[m] = 1 \right\}$  as the set of all *probability distributions* over  $\tilde{\mathcal{Y}}$ . A *noisy kernel* is defined to be a map:

$$\mathcal{K} : \mathcal{X} \times \mathcal{Y} \rightarrow 2^{\mathcal{D}(\tilde{\mathcal{Y}})},$$

where  $2^{\mathcal{D}(\tilde{\mathcal{Y}})}$  is the set of all *subsets* of  $\mathcal{D}(\tilde{\mathcal{Y}})$ , i.e., the kernel  $\mathcal{K}$  maps each  $(\mathbf{x}, y) \in \mathcal{X} \times \mathcal{Y}$  to a *subset of distributions*  $\mathcal{K}(\mathbf{x}, y) \subset \mathcal{D}(\tilde{\mathcal{Y}})$ . We will write  $\mathcal{Q}_y^{\mathbf{x}} = \mathcal{K}(\mathbf{x}, y)$  for notational convenience.

For any time horizon  $T$ , we denote by  $\mathbf{P}$  a set of general *random processes* over  $\mathcal{X}^T$  that models the generating mechanism of the features. For any  $t \in [T]$ , we write  $\mathbf{x}^t = \{\mathbf{x}_1, \dots, \mathbf{x}_t\}$ ,  $y^t = \{y_1, \dots, y_t\}$  and  $\tilde{y}^t = \{\tilde{y}_1, \dots, \tilde{y}_t\}$ . Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be a class of *hypotheses*,  $\mathbf{P}$  be a class of random

processes over  $\mathcal{X}^T$ , and  $\mathcal{K}$  be a noisy kernel as defined above. We consider the following *robust online classification* scenario:

1. At the beginning, *Nature* selects some  $h \in \mathcal{H}$  and  $\boldsymbol{\nu}^T \in \mathcal{P}$ , and samples  $\mathbf{x}^T \sim \boldsymbol{\nu}^T$ ;
2. At each time step  $t$ , *Nature* reveals  $\mathbf{x}_t$  to a *learner*, where  $\mathbf{x}_t$  is the  $t$ th sample in  $\mathbf{x}^T$ ;
3. *Learner* predicts  $\hat{y}_t \in \mathcal{Y}$ , based on (noisy) history observed thus far (i.e.,  $\mathbf{x}^t, \tilde{y}^{t-1}$ );
4. An *adversary* then selects  $\tilde{p}_t \in \mathcal{K}(\mathbf{x}_t, h(\mathbf{x}_t))$ , samples a *noisy* sample  $\tilde{y}_t \sim \tilde{p}_t$ , and reveals  $\tilde{y}_t$  to the learner.

The goal of the *learner* is to minimize the *cumulative error*  $\sum_{t=1}^T 1\{h(\mathbf{x}_t) \neq \hat{y}_t\}$ .

Note that the cumulative error is a *random variable* that depends on all the randomness associated with the game. To remove the dependency on such randomness and to assess the fundamental limits of the prediction quality, we consider the following two measures:

**Definition 4** Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be the set of hypotheses,  $\mathcal{P}$  be a set of random processes over  $\mathcal{X}^T$ , and  $\mathcal{K} : \mathcal{X} \times \mathcal{Y} \rightarrow 2^{\mathcal{D}(\tilde{\mathcal{Y}})}$  be a noisy kernel. We denote by  $\Phi$  and  $\Psi$  the (possibly randomized) strategies of the learner and adversary, respectively. The expected minimax risk is defined as:

$$\tilde{r}_T(\mathcal{H}, \mathcal{P}, \mathcal{K}) = \inf_{\Phi} \sup_{h \in \mathcal{H}, \boldsymbol{\nu}^T \in \mathcal{P}, \Psi} \mathbb{E} \left[ \sum_{t=1}^T 1\{h(\mathbf{x}_t) \neq \hat{y}_t\} \right], \quad (3)$$

where  $\mathbf{x}^T \sim \boldsymbol{\nu}^T$ ,  $\hat{y}_t = \Phi(\mathbf{x}^t, \tilde{y}^{t-1})$ ,  $\tilde{y}_t \sim \tilde{p}_t$  and  $\tilde{p}_t = \Psi(\mathbf{x}^t, \hat{y}^t) \in \mathcal{K}(\mathbf{x}_t, h(\mathbf{x}_t))$ . Moreover, the expectation is over all the randomness involved.

**Definition 5** Let  $\mathcal{H}$ ,  $\mathcal{P}$ ,  $\mathcal{K}$ ,  $\Phi$  and  $\Psi$  be as in Definition 4. For any confidence parameter  $\delta > 0$ , the high probability minimax risk at confidence  $\delta$  is defined to be the minimum cumulative error  $B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}) \geq 0$  such that there exists predictor  $\Phi$  satisfying:

$$\sup_{h \in \mathcal{H}, \boldsymbol{\nu}^T \in \mathcal{P}, \Psi} \Pr \left[ \sum_{t=1}^T 1\{h(\mathbf{x}_t) \neq \hat{y}_t\} \geq B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}) \right] \leq \delta. \quad (4)$$

**Example 1 (Robust Hypothesis Testing)** Let  $\mathcal{Y} = \{0, 1\}$ ,  $\mathcal{X} = \{\mathbf{x}\}$  be a singleton set,  $\mathcal{H} = \{h_i(\mathbf{x}) = i : i \in \mathcal{Y}\}$  and  $\tilde{\mathcal{Y}}$  be any finite set of size  $M$ . We define a kernel  $\mathcal{K}$  such that  $\mathcal{K}(\mathbf{x}, i) = \mathcal{Q}_i \subset \mathcal{D}(\tilde{\mathcal{Y}})$  with  $i \in \mathcal{Y}$ . Our robust online classification game is reduced to the following hypothesis testing problem: at the beginning *Nature* fixes some  $i \in \{0, 1\}$ ; at each time step  $t$ , an adversary selects some  $\tilde{p}_t \in \mathcal{Q}_i$  and samples  $\tilde{y}_t \sim \tilde{p}_t$ . The goal of the predictor is to identify the index  $i$  from the observation of  $\tilde{y}^T$ . Note that the distribution  $\tilde{p}_t$  changes at every step with the only restriction that it must be selected from the same distribution class  $\mathcal{Q}_i$ . This differs from classical hypothesis testing, where the distribution is selected once and used for all time steps.

**Example 2 (Binary Classification with Bernoulli Noise)** Let  $\mathcal{Y} = \tilde{\mathcal{Y}} = \{0, 1\}$  and  $\mathcal{H} \subset \{0, 1\}^{\mathcal{X}}$  be any hypothesis class with instance space  $\mathcal{X}$ . For any given  $\eta \in [0, \frac{1}{2})$ , we define the following kernel  $\mathcal{K}$ : for  $i \in \{0, 1\}$ ,  $\mathcal{K}(\mathbf{x}, i) = \{\text{Bernolli}(|i - \eta'|) : \eta' \in [0, \eta]\} \subset \mathcal{D}(\tilde{\mathcal{Y}})$ . Then, our robust online classification game recovers the bounded noisy setting as introduced in [Ben-David et al.](#)

(2009). More generally, we can also consider the following Tsybakov type noise kernel<sup>1</sup>: for any  $\mathbf{x}^T \in \mathcal{X}^T$ ,  $i \in \{0, 1\}$ , we have  $\mathcal{K}(\mathbf{x}_t, i) = \{\text{Bernolli}(|i - \eta'|) : \eta' \in [0, \eta_t]\} \subset \mathcal{D}(\tilde{\mathcal{Y}})$ , such that for some  $A > 0$  and  $0 \leq \alpha < 1$ , we set  $\forall r \in (0, 1/2]$ ,  $\frac{1}{T} \sum_{t=1}^T \mathbb{1}\{\eta_t \geq \frac{1}{2} - r\} \leq Ar^{\frac{\alpha}{1-\alpha}}$ . We refer to Section 3.3 for a detailed analysis of such scenarios.

**Remark 6** Note that we have assumed that the underlying truth of our robust online classification game is selected from the class  $\mathcal{H}$ . This is justified by the fact that our guarantee of minimax risk is also evaluated for the underlying truth when observing only noisy labels. In Section 3.4 (Remark 31) we discuss how our approach can be extended to non-realizable cases.

## 2.1. Bregman Divergence, Exp-concavity and Conditional Distribution Estimation

We now introduce several key technical concepts and results with proofs deferred to Appendix C. Let  $\mathcal{D}(\tilde{\mathcal{Y}})$  be the set of probability distributions over  $\tilde{\mathcal{Y}}$ . A function  $L : \mathcal{D}(\tilde{\mathcal{Y}}) \times \mathcal{D}(\tilde{\mathcal{Y}}) \rightarrow \mathbb{R}^{\geq 0}$  is referred to as a *divergence*. We say a divergence  $L$  is a *Bergman divergence* if there exists a strictly convex function  $F : \mathcal{D}(\tilde{\mathcal{Y}}) \rightarrow \mathbb{R}$  such that for any  $p, q \in \mathcal{D}(\tilde{\mathcal{Y}})$ ,  $L(p, q) = F(p) - F(q) - (p - q)^\top \nabla F(q)$ . Note that both KL-divergence  $\text{KL}(p, q) = \sum_{\tilde{y} \in \tilde{\mathcal{Y}}} p[\tilde{y}] \log \frac{p[\tilde{y}]}{q[\tilde{y}]}$  and the  $L^2$ -divergence  $L^2(p, q) = \|p - q\|_2^2$  are Bergman divergences (Cesa-Bianchi and Lugosi, 2006, Chapter 11.2).

**Proposition 7** Let  $P$  be a random variable over  $\mathcal{D}(\tilde{\mathcal{Y}})$  (i.e., a random variable with values in  $\mathbb{R}^M$ ) and  $L$  be a Bergman divergence. Then for any  $q_1, q_2 \in \mathcal{D}(\tilde{\mathcal{Y}})$

$$\mathbb{E}_{p \sim P}[L(p, q_1) - L(p, q_2)] = L(\mathbb{E}_{p \sim P}[p], q_1) - L(\mathbb{E}_{p \sim P}[p], q_2).$$

A function  $\ell : \tilde{\mathcal{Y}} \times \mathcal{D}(\tilde{\mathcal{Y}}) \rightarrow \mathbb{R}^{\geq 0}$  is referred to as a *loss function*. For instance, the *log-loss* is defined as  $\ell^{\log}(\tilde{y}, p) = \text{KL}(e_{\tilde{y}}, p)$ , and the *Brier loss* is defined as  $\ell^{\text{B}}(\tilde{y}, p) = L^2(e_{\tilde{y}}, p)$ , where  $e_{\tilde{y}}$  is the probability distribution that assigns probability 1 to  $\tilde{y}$ . We say a loss  $\ell$  is  $\alpha$ -*Exp-concave* if for any  $\tilde{y} \in \tilde{\mathcal{Y}}$ , the function  $e^{-\alpha \ell(\tilde{y}, p)}$  is concave w.r.t.  $p$  for some  $\alpha \in \mathbb{R}^{\geq 0}$ .

**Proposition 8** The log-loss is 1-Exp-concave and the Brier loss is 1/4-Exp-concave.

**Online conditional distribution estimation:** The key to our noisy classification problem is a distribution estimation that we discuss next. Let  $\mathcal{F} \subset \mathcal{D}(\tilde{\mathcal{Y}})^{\mathcal{X}}$  be a class of functions mapping  $\mathcal{X}$  to distributions in  $\mathcal{D}(\tilde{\mathcal{Y}})$ . The *online conditional distribution estimation* is a game between *Nature* and *predictor* that follows the following protocol: (1) at each times step  $t$ , Nature selects some  $\mathbf{x}_t \in \mathcal{X}$  and reveals it to the predictor; (2) the predictor then makes a prediction  $\hat{p}_t \in \mathcal{D}(\tilde{\mathcal{Y}})$ , based on  $\mathbf{x}^t, \tilde{y}^{t-1}$ ; (3) Nature then selects some  $\tilde{p}_t \in \mathcal{D}(\tilde{\mathcal{Y}})$ , samples  $\tilde{y}_t \sim \tilde{p}_t$  and reveals  $\tilde{y}_t$  to the predictor. The goal of the predictor is to minimize the following *regret*

$$R_T(\mathcal{F}) = \sup_{\mathbf{x}^T \in \mathcal{X}^T, f \in \mathcal{F}, \tilde{p}^T \in \mathcal{D}(\tilde{\mathcal{Y}})^T} \mathbb{E}_{\tilde{y}^T} \left[ \sum_{t=1}^T L(\tilde{p}_t, \hat{p}_t) - L(\tilde{p}_t, f(\mathbf{x}_t)) \right],$$

where  $\tilde{y}_t \sim \tilde{p}_t$ ,  $\hat{p}_t = \Phi(\mathbf{x}^t, \tilde{y}^{t-1})$  and  $L$  is any divergence. Note that  $\tilde{p}_t$  is understood as the *conditional distribution* of  $\tilde{p}^T$  conditioning on  $\tilde{y}^{t-1}$ . We should emphasis that the distributions  $\tilde{p}^T$

1. Note that the noisy kernel  $\mathcal{K}$  in general is assumed to be *known* by the predictor. However, for the Tsybakov type noise, we do not assume the parameters  $\eta_t$ s are known.

are *not* necessarily realizable by  $f$  and are selected completely arbitrarily. This is in contrast to the well-specified case employed in Foster et al. (2021), and is the key that allows us to deal with the *unknown* noisy distribution (which is necessarily being *miss-specified*).

**Lemma 9** *Let  $\mathcal{F}$  be a finite class and  $L$  be a Bergman divergence such that the induced loss  $\ell(\tilde{y}, p) \stackrel{\text{def}}{=} L(e_{\tilde{y}}, p)$  is  $\alpha$ -Exp-concave. Then for the EWA predictor  $\Phi$  we have*

$$\sup_{\mathbf{x}^T \in \mathcal{X}^T, f \in \mathcal{F}, \tilde{p}^T \in \mathcal{D}(\tilde{\mathcal{Y}}^T)} \mathbb{E}_{\tilde{y}^T} \left[ \sum_{t=1}^T L(\tilde{p}_t, \hat{p}_t) - L(\tilde{p}_t, f(\mathbf{x}_t)) \right] \leq \frac{\log |\mathcal{F}|}{\alpha},$$

where  $\hat{p}_t = \Phi(\mathbf{x}^t, \tilde{y}^{t-1})$ ,  $\tilde{y}_t \sim \tilde{p}_t$ , and  $\hat{p}_t$  is a convex combination of  $\{f(\mathbf{x}_t) : f \in \mathcal{F}\}$  for all  $t \in [T]$ .

We refer to Appendix B for construction of the EWA predictor (Algorithm 2). Note that the property of  $L$  being Bregman divergence plays a central rule in the proof of Lemma 9 (see Appendix C), thanks to Proposition 7.

**Lemma 10** *Let  $\mathcal{F}$  and  $L$  be as in Lemma 9 and  $L$  be bounded by  $M$ . Then for the EWA predictor  $\Phi$  we have for all  $\delta > 0$*

$$\sup_{\mathbf{x}^T \in \mathcal{X}^T, f \in \mathcal{F}, \tilde{p}^T \in \mathcal{D}(\tilde{\mathcal{Y}}^T)} \Pr_{\tilde{y}^T} \left[ \sum_{t=1}^T L(\tilde{p}_t, \hat{p}_t) - L(\tilde{p}_t, f(\mathbf{x}_t)) \geq \frac{\log |\mathcal{F}|}{\alpha} + M\sqrt{(T/2) \log(1/\delta)} \right] \leq \delta,$$

where  $\hat{p}_t = \Phi(\mathbf{x}^t, \tilde{y}^{t-1})$ ,  $\tilde{y}_t \sim \tilde{p}_t$ , and  $\hat{p}_t$  is a convex combination of  $\{f(\mathbf{x}_t) : f \in \mathcal{F}\}$  for all  $t \in [T]$ .

Note that Lemma 10 requires the loss to be bounded by  $M$ , which generally is not satisfied by the KL-divergence. However, for KL-divergence, one can always truncate  $\hat{p}_t$  as in Wu et al. (2022b) to ensure the probability mass is bounded away from 0.

### 3. Main Results

This is main section of our paper. We present results for the *minimax risk* of our *robust online classification game* as defined in Section 2 for various hypothesis classes  $\mathcal{H}$  and noisy kernel  $\mathcal{K}$ , with guarantees both in expectation and with high probability.

#### 3.1. The Binary Label Case

We first consider a simpler case where we assume the label space  $\mathcal{Y} = \{0, 1\}$  is binary valued. For any  $\mathbf{x} \in \mathcal{X}$  and  $y \in \mathcal{Y}$  we write  $\mathcal{K}(\mathbf{x}, y) = \mathcal{Q}_y^{\mathbf{x}} \subset \mathcal{D}(\tilde{\mathcal{Y}})$ . We can assume, w.l.o.g., that the  $\mathcal{Q}_y^{\mathbf{x}}$ s are *convex* and *closed* sets, since the adversary can select arbitrary distribution from  $\mathcal{Q}_y^{\mathbf{x}}$ s at each time step, including randomized strategies that effectively sample from a mixture (i.e., convex combination) of distributions in  $\mathcal{Q}_y^{\mathbf{x}}$ s.

Clearly, one must introduce some constraints on the kernel  $\mathcal{K}$  in order to obtain meaningful results. To do so, we introduce the following *well-separation* condition:

**Definition 11** *Let  $L : \mathcal{D}(\tilde{\mathcal{Y}}) \times \mathcal{D}(\tilde{\mathcal{Y}}) \rightarrow \mathbb{R}^{\geq 0}$  be a divergence, we say the kernel  $\mathcal{K}$  is well-separated w.r.t.  $L$  at scale  $\gamma > 0$ , if*

$$\forall \mathbf{x} \in \mathcal{X}, \forall y \neq y' \in \mathcal{Y}, L(\mathcal{Q}_y^{\mathbf{x}}, \mathcal{Q}_{y'}^{\mathbf{x}}) \stackrel{\text{def}}{=} \inf_{p \in \mathcal{Q}_y^{\mathbf{x}}, q \in \mathcal{Q}_{y'}^{\mathbf{x}}} L(p, q) \geq \gamma.$$

We are now ready to state our first main result:

**Theorem 12** *Let  $\mathcal{H} \subset \{0, 1\}^{\mathcal{X}}$  be any finite binary valued class,  $\mathcal{K}$  be any noisy kernel that is well-separated at scale  $\gamma$  w.r.t. the  $L^2$  divergence, and  $\mathbf{P}$  be the class of all singleton distributions over  $\mathcal{X}^T$  (i.e., we assume  $\mathbf{x}^T$  is generated adversarially). Then, the expected minimax risk, as in Definition 4, is upper bounded by*

$$\tilde{r}_T(\mathcal{H}, \mathbf{P}, \mathcal{K}) \leq \frac{16 \log |\mathcal{H}|}{\gamma}.$$

Moreover, for any  $\delta > 0$ , the high probability minimax risk, as in Definition 5, is upper bounded by

$$B^\delta(\mathcal{H}, \mathbf{P}, \mathcal{K}) \leq \frac{16 \log |\mathcal{H}| + \sqrt{8T \log(1/\delta)}}{\gamma}.$$

It is important to point out that our minimax risk bound in Theorem 12 is *independent* of the size of the noisy observations  $\tilde{\mathcal{Y}}$ . Before we present a formal proof of Theorem 12, we first establish the following simple geometry fact that is crucial for our proof.

**Lemma 13** *Let  $\mathcal{Q} \subset \mathcal{D}(\tilde{\mathcal{Y}})$  be a convex and closed set,  $p$  be a point outside of  $\mathcal{Q}$  with  $\gamma \stackrel{\text{def}}{=} \inf_{q \in \mathcal{Q}} L^2(p, q)$ . Denote by  $q^* \in \mathcal{Q}$  the (unique) point that attains  $L^2(p, q^*) = \gamma$ . Then for any  $q \in \mathcal{Q}$ , we have*

$$L^2(q, p) - L^2(q, q^*) \geq L^2(p, q^*) = \gamma.$$

**Proof** By the *hyperplane separation theorem*, the hyperplane perpendicular to line segment  $p - q^*$  at  $q^*$  separates  $\mathcal{Q}$  and  $p$ . Therefore, the degree  $\theta$  of angle formed by  $p - q^* - q$  is greater than  $\pi/2$ . By the law of cosines, we have  $L^2(p, q) \geq L^2(q, q^*) + L^2(q^*, p) = L^2(q, q^*) + \gamma$ , as needed. ■

**Proof [Proof of Theorem 12]** Our key idea of proving Theorem 12 is to reduce the noisy classification problem to a suitable conditional distribution learning problem leveraging the results from Section 2 to derive the desired minimax risk bounds. To achieve this, we define the following distribution valued function class  $\mathcal{F}$  using the hypothesis class  $\mathcal{H}$  and noisy kernel  $\mathcal{K}$ . For any  $\mathbf{x} \in \mathcal{X}$ , we denote by  $\mathcal{Q}_0^{\mathbf{x}}$  and  $\mathcal{Q}_1^{\mathbf{x}}$  the noisy distribution sets corresponding to labels 0 and 1, respectively. Since the kernel  $\mathcal{K}$  is well-separated at scale  $\gamma$  under  $L^2$  divergence, we have, by the *hyperplane separation theorem*, that there must be  $q_0^{\mathbf{x}} \in \mathcal{Q}_0^{\mathbf{x}}$  and  $q_1^{\mathbf{x}} \in \mathcal{Q}_1^{\mathbf{x}}$  such that

$$L^2(q_0^{\mathbf{x}}, q_1^{\mathbf{x}}) = L^2(\mathcal{Q}_0^{\mathbf{x}}, \mathcal{Q}_1^{\mathbf{x}}) \geq \gamma.$$

We now define for any  $h \in \mathcal{H}$  the function  $f_h$  such that  $\forall \mathbf{x} \in \mathcal{X}$ ,  $f_h(\mathbf{x}) = q_{h(\mathbf{x})}^{\mathbf{x}}$ . Let  $\mathcal{F} = \{f_h : h \in \mathcal{H}\}$  and  $\Phi$  be the predictor from Lemma 9 with class  $\mathcal{F}$  and  $L^2$  divergence (using  $\mathbf{x}^T, \tilde{y}^T$  from the *original* noisy classification game). We define our *classification* predictor as follows:

$$\hat{y}_t = \arg \min_y \{L^2(q_y^{\mathbf{x}^t}, \hat{p}_t) : y \in \{0, 1\}\},$$

i.e., we predict the label  $y$  so that  $q_y^{\mathbf{x}^t}$  is closer to  $\hat{p}_t$  under  $L^2$  divergence, where  $\hat{p}_t = \Phi(\mathbf{x}^t, \tilde{y}^{t-1})$ .

Let  $h^* \in \mathcal{H}$  be the underlying true classification function. We have by Lemma 9 and 1/4-Exp-concavity of  $L^2$  divergence that

$$\sup_{\mathbf{x}^T \in \mathcal{X}^T} \mathbb{E}_{\tilde{y}^T} \left[ \sum_{t=1}^T L^2(\tilde{p}_t, \hat{p}_t) - L^2(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \right] \leq 4 \log |\mathcal{F}|, \quad (5)$$

where  $\tilde{y}_t \sim \tilde{p}_t$  and  $\tilde{p}_t \in \mathcal{Q}_{h^*(\mathbf{x}_t)}^{\mathbf{x}_t}$  is the noisy distribution selected by the adversary (this is true since Lemma 9 holds for *any* distributions  $\tilde{p}^T$ ).

For any time step  $t$ , we denote  $y_t = h^*(\mathbf{x}_t)$  as the true label. We now observe that  $\tilde{p}_t \in \mathcal{Q}_{y_t}^{\mathbf{x}_t}$ ,  $f_{h^*}(\mathbf{x}_t) = q_{y_t}^{\mathbf{x}_t}$  and  $\hat{p}_t$  is a convex combination of  $q_0^{\mathbf{x}_t}$  and  $q_1^{\mathbf{x}_t}$ , where the last assertion follows by the last part of Lemma 9 and the construction of  $\mathcal{F}$ . By the construction of  $q_y^{\mathbf{x}_t}$ , we have  $q_y^{\mathbf{x}_t}$  is the closest point in  $\mathcal{Q}_y^{\mathbf{x}_t}$  to  $\hat{p}_t$  under  $L^2$  divergence for  $y \in \{0, 1\}$ . Therefore, by Lemma 13, we have

$$L^2(\tilde{p}_t, \hat{p}_t) - L^2(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \geq L^2(\hat{p}_t, f_{h^*}(\mathbf{x}_t)).$$

Denote  $a_t = L^2(\tilde{p}_t, \hat{p}_t) - L^2(\tilde{p}_t, f_{h^*}(\mathbf{x}_t))$  for notational convenience. We have:

1. For all  $t \in [T]$ ,  $a_t \geq 0$  since  $L^2$  divergence is always positive;
2. If  $\hat{y}_t \neq y_t$ , then  $a_t \geq \gamma/4$ . This is because the event  $\{\hat{y}_t \neq y_t\}$  implies that  $\hat{p}_t$  is closer to  $q_{1-y_t}^{\mathbf{x}_t}$  than  $q_{y_t}^{\mathbf{x}_t}$  under  $L^2$ , and hence  $L^2(\hat{p}_t, f_{h^*}(\mathbf{x}_t)) = L^2(\hat{p}_t, q_{y_t}^{\mathbf{x}_t}) \geq (\sqrt{\gamma}/2)^2 = \gamma/4$ , since  $L^2(\hat{p}_t, q_{y_t}^{\mathbf{x}_t}) \geq L^2(\hat{p}_t, q_{1-y_t}^{\mathbf{x}_t})$  and  $\sqrt{L^2(\hat{p}_t, q_{y_t}^{\mathbf{x}_t})} + \sqrt{L^2(\hat{p}_t, q_{1-y_t}^{\mathbf{x}_t})} = \sqrt{L^2(q_{y_t}^{\mathbf{x}_t}, q_{1-y_t}^{\mathbf{x}_t})} \geq \sqrt{\gamma}$ .

Putting all together we arrive at

$$\sum_{t=1}^T 1\{\hat{y}_t \neq y_t\} \leq \frac{4}{\gamma} \sum_{t=1}^T L^2(\tilde{p}_t, \hat{p}_t) - L^2(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)).$$

The expected minimax risk will now follow from (5). To prove the high probability minimax risk, we can replicate the argument above but replacing (5) with the result in Lemma 10 and noting that  $L^2$  divergence is bounded by 1.  $\blacksquare$

**Remark 14** We should remark that the selection of  $L^2$  divergence plays a central rule in the proof of Theorem 12 thanks to its nice geometry property. We note that a naive extension to the stronger KL-divergence does not work, mainly due to the fact that if  $q$  is a projection of point  $p$  onto a convex set under KL-divergence, it does not necessarily imply that  $q$  is the projection of any point along the line segment of  $p$  and  $q$ . Therefore, our central argument in the proof of Theorem 12 that relates  $1\{\hat{y}_t \neq y_t\}$  and  $L(\tilde{p}_t, \hat{p}_t) - L(\tilde{p}_t, f_{h^*}(\mathbf{x}_t))$  will not go through. Nonetheless, we will show in Section 3.3 how to overcome these limitations for certain special, yet important, noisy kernels.

**Remark 15** Note that our approach in the proof of Theorem 12 is robust enough to accommodate the case when there is a small number  $D$  of time steps where the noisy samples are generated arbitrarily. This will result in an  $O\left(\frac{\log |\mathcal{H}| + D}{\gamma}\right)$  expected risk using the same approach, see also Theorem 23 for an concrete example of such scenarios.

### 3.2. Multi-Label Case through Pairwise Comparison

In this section, we consider the case when the label space  $\mathcal{Y}$  is of size  $N \geq 2$ . The main difficulty of generalizing our approach from the proof of Theorem 12 is the definition of the class  $\mathcal{F}$ . Note that in the binary label case we define  $\mathcal{F}$  with values  $q_0^{\mathbf{x}}, q_1^{\mathbf{x}}$  that attain  $L^2(q_0^{\mathbf{x}}, q_1^{\mathbf{x}}) = L^2(\mathcal{Q}_0^{\mathbf{x}}, \mathcal{Q}_1^{\mathbf{x}})$ . In the multi-label case, this selection is less obvious, since for any  $y \in \mathcal{Y}$  the closest points in  $\mathcal{Q}_y^{\mathbf{x}}$  to different sets  $\mathcal{Q}_{y'}^{\mathbf{x}}$  are *different*. There is no canonical way of assigning the value  $f_h(\mathbf{x})$  if  $h(\mathbf{x}) = y$ . Our key idea is to reduce the multi-label classification problem to the problem of pairwise comparison that can be handled using our techniques from Section 3.1.

To begin, we first introduce the following technical concepts. Recall that our robust online classification problem is completely determined by the tuple  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$  of hypothesis class  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$ , random process class  $\mathbf{P}$ , and noisy kernel  $\mathcal{K}$ .

**Definition 16** A robust online classification problem  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$  is said to be pairwise testable with confidence  $\delta > 0$  and error bound  $C(\delta) \geq 0$ , if for any pair  $h_i, h_j \in \mathcal{H}$ , the sub-problem  $(\{h_i, h_j\}, \mathbf{P}, \mathcal{K})$  admits a high probability minimax risk  $B^\delta(\{h_i, h_j\}, \mathbf{P}, \mathcal{K}) \leq C(\delta)$  at confidence  $\delta$  (see Definition 5).

Suppose that a robust online classification problem  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$  is *pairwise testable* and the class  $\mathcal{H} = \{h_1, \dots, h_K\}$  is finite of size  $K$ . Let  $\Phi_{i,j}$  be the predictor for the sub-problem  $(\{h_i, h_j\}, \mathbf{P}, \mathcal{K})$  with error bound  $C(\delta/(2K))$  and confidence  $\delta/(2K) > 0$ . Let  $\mathbf{x}^T, \tilde{y}^T$  be any realization of problem  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$ . We define for any  $h_i \in \mathcal{H}$  and  $t \in [T]$  a *surrogate loss* vector

$$\forall j \in [K], \mathbf{v}_t^i[j] = 1\{\Phi_{i,j}(\mathbf{x}^t, \tilde{y}^{t-1}) \neq h_i(\mathbf{x}_t) \text{ and } h_i(\mathbf{x}_t) \neq h_j(\mathbf{x}_t)\}, \quad (6)$$

i.e., the loss  $\mathbf{v}_t^i[j] = 1$  if and only if  $h_i(\mathbf{x}_t) \neq h_j(\mathbf{x}_t)$  and the predictor  $\Phi_{i,j}(\mathbf{x}^t, \tilde{y}^{t-1})$  differs from  $h_i(\mathbf{x}_t)$ . The purpose of such a surrogate loss will be revealed shortly. We now construct our prediction rule for  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$  as in Algorithm 1.

We now ready to state our main result of this section:

**Theorem 17** Let  $\mathcal{H} = \{h_1, \dots, h_K\}$  be a finite hypothesis class of size  $K$ ,  $\mathbf{P}$  be any random processes over  $\mathcal{X}^T$ , and  $\mathcal{K}$  be any noisy kernel. If the robust online classification problem  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$  is pairwise testable, and  $\Phi_{i,j}$ s are the predictors that achieves confidence  $\delta/(2K) > 0$  and error bound  $C(\delta/(2K))$  as in Definition 16, then the predictor in Algorithm 1 with input  $\mathcal{H}$ ,  $\Phi_{i,j}$ s and  $C = C(\delta/(2K))$  achieves high probability minimax risk at confidence  $\delta$  upper bounded by

$$B^\delta(\mathcal{H}, \mathbf{P}, \mathcal{K}) \leq O(C(\delta/(2K)) \log K + \log(2/\delta)), \quad (7)$$

where  $O$  hides an absolute constant.

**Proof** Let  $h_{k^*} \in \mathcal{H}$  be the underlying true classification function and  $\mathbf{x}^T$  be any realization of the features. By the definition of *pairwise testability* and union bound, we have for any  $k \in [K]$ , there exists a predictor  $\Phi_{k,k^*}$  such that w.p.  $\geq 1 - \delta/2$  over the randomness of the data from the *original* game  $(\mathcal{H}, \mathbf{P}, \mathcal{K})$  that

$$\forall k \in [K], \sum_{t=1}^T 1\{h_{k^*}(\mathbf{x}_t) \neq \Phi_{k,k^*}(\mathbf{x}^t, \tilde{y}^{t-1})\} \leq C(\delta/(2K)). \quad (8)$$

---

**Algorithm 1** Multi-Label Noisy Predictor via Pairwise Comparison
 

---

**Input:** Class  $\mathcal{H} = \{h_1, \dots, h_K\}$ , predictor  $\Phi_{i,j}$  for  $i, j \in [K]$  and error bound  $C$

Set  $S^1 = \{1, \dots, K\}$ ;

**for**  $t = 1, \dots, T$  **do**

    Receive  $\mathbf{x}_t$ ;

    Sampling index  $\hat{k}_t$  from  $S^t$  *uniformly* and make prediction:

$$\hat{y}_t = h_{\hat{k}_t}(\mathbf{x}_t);$$

    Receive noisy label  $\tilde{y}_t$ ;

    Set  $S^{t+1} = \emptyset$ ;

**for**  $i \in S^t$  **do**

        Compute  $l_t^i = \max_{j \in [K]} \sum_{r=1}^t \mathbf{v}_r^i[j]$ , where  $\mathbf{v}_t^i[j]$  is defined in (6);

**if**  $l_t^i \leq C$  **then**

            Update  $S^{t+1} = S^{t+1} \cup \{i\}$ ;

**end**

**end**

**end**

---

Note that for any other  $\{i, j\} \not\supset k^*$ , we do not assume any guarantees for the predictor  $\Phi_{i,j}$  in the definition of pairwise testability.

We now condition on the event defined in (8). Let  $\mathbf{v}_t^k$  with  $k \in [K]$  and  $t \in [T]$  be the *surrogate loss* vector as defined in (6). We now observe the following key properties

1. We have for all  $t \in [T]$  that

$$\max_{j \in [K]} \sum_{r=1}^t \mathbf{v}_r^{k^*}[j] \leq C(\delta/(2K)); \quad (9)$$

2. For any  $k \neq k^*$ , we have for all  $t \in [T]$ :

$$\max_{j \in [K]} \sum_{r=1}^t \mathbf{v}_r^k[j] \geq \sum_{r=1}^t 1\{h_k(\mathbf{x}_r) \neq h_{k^*}(\mathbf{x}_r)\} - C(\delta/(2K)). \quad (10)$$

The first property is straightforward by the definition of  $\mathbf{v}_t^k$  and (8). The second property holds since the lower bound is attained when  $j = k^*$ .

We now analyze the performance of Algorithm 1. By property (9) we know that  $k^* \in S^t$  for all  $t \in [T]$ , i.e.,  $|S^t| \geq 1$ . Let  $N_t = |S^t|$ . We define for all  $t \in [T]$  the following *potential*:

$$E_t = \sum_{k \in S^t} \max \left\{ 0, 2C(\delta/(2K)) - \sum_{r=1}^t 1\{h_k(\mathbf{x}_r) \neq h_{k^*}(\mathbf{x}_r)\} \right\}.$$

Clearly, we have  $E_t \leq 2C(\delta/(2K))N_t$ . Let  $D_t = |\{k \in S^t : h_k(\mathbf{x}_t) \neq h_{k^*}(\mathbf{x}_t)\}|$ . We have

$$D_t \leq N_t - N_{t+1} + E_t - E_{t+1}, \quad (11)$$

since for any  $k \in S_t$  such that  $h_k(\mathbf{x}_t) \neq h_{k^*}(\mathbf{x}_t)$  either  $k$  is removed from  $S^{t+1}$  (which contributes at most  $N_t - N_{t+1}$ ) or its contribution to  $E_{t+1}$  is decreased by 1 when compared to  $E_t$  (this is because by our construction of Algorithm 1 and property (10) that once the contributions of  $k$  to  $E_t$  equals 0 it must be excluded from  $S^{t+1}$ ). We have by definition of  $\hat{y}_t$  that

$$\mathbb{E}[1\{h_{k^*}(\mathbf{x}_t) \neq \hat{y}_t\}] = \frac{D_t}{|S^t|} \leq \frac{N_t - N_{t+1} + E_t - E_{t+1}}{N_t}. \quad (12)$$

By a standard argument (Kakade and Kalai, 2005, Theorem 2), we have

$$\sum_{t=1}^T \frac{N_t - N_{t+1}}{N_t} \leq \sum_{t=1}^T \left( \frac{1}{N_t} + \frac{1}{N_t - 1} + \cdots + \frac{1}{N_{t+1} + 1} \right) \leq \sum_{k=1}^K \frac{1}{k} \leq \log K.$$

Moreover, we observe that

$$\sum_{t=1}^T \frac{E_t - E_{t+1}}{N_t} \leq \frac{2C(\delta/(2K))N_1 - E_2}{N_1} + \sum_{t=2}^T \frac{E_t - E_{t+1}}{N_t} \quad (13)$$

$$\leq \frac{2C(\delta/(2K))(N_1 - N_2)}{N_1} + \frac{2C(\delta/(2K))N_2 - N_3}{N_2} + \sum_{t=3}^T \frac{E_t - E_{t+1}}{N_t} \quad (14)$$

$$\dots \quad (15)$$

$$\leq 2C(\delta/(2K)) \sum_{t=1}^T \frac{N_t - N_{t+1}}{N_t} \leq 2C(\delta/(2K)) \log K, \quad (16)$$

where the inequalities follow from the fact that  $E_t \leq 2C(\delta/(2K))N_t$  and  $N_t \geq N_{t+1}$ .

Therefore, we conclude

$$\mathbb{E} \left[ \sum_{t=1}^T 1\{h_{k^*}(\mathbf{x}_t) \neq \hat{y}_t\} \right] \leq (1 + 2C(\delta/(2K))) \log K,$$

where the randomness is on the selection of  $\hat{k}_t \sim S^t$ . Since our selection of  $\hat{k}_t$  are independent (conditioning on  $S^t$ ) for different  $t$  and the indicator is bounded by 1 and non-negative, we can invoke Lemma 36 (second part) to obtain a high probability guarantee of confidence  $\delta/2$  by introducing an extra  $\log(2/\delta)$  additive term. The theorem now follows by a union bound with the event (8). ■

For  $L^2$  loss and well-separated kernel, we can instantiate our previous results as presented next.

**Theorem 18** *Let  $\mathcal{H}$  be a finite class of size  $K$ ,  $\mathcal{P}$  be the class of all singleton distributions over  $\mathcal{X}^T$  and  $\mathcal{K}$  be a kernel that is well-separated at scale  $\gamma$  w.r.t.  $L^2$  divergence. Then, the high probability minimax risk with confidence  $\delta > 0$  is upper bounded by*

$$B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}) \leq O \left( \frac{\log(K) \log(K/(2\delta\gamma^2))}{\gamma^2} \right). \quad (17)$$

**Proof** Clearly, a simple application of the high probability minimax risk bound in Theorem 12 and Theorem 17 yields a high probability risk bound of order  $O(\log K (\log K + \sqrt{T \log(K/\delta)})/\gamma)$ .

We now prove a stronger concentration result for hypothesis class with 2 elements. Let  $\{h_1, h_2\}$  be a class with 2 elements. We may assume, w.l.o.g., that  $h_1, h_2$  are binary valued with labels in  $\{0, 1\}$  (relabeling if necessary). Let  $q_0^{\mathbf{x}}, q_1^{\mathbf{x}}$  and  $\mathcal{Q}_0^{\mathbf{x}}, \mathcal{Q}_1^{\mathbf{x}}$  be as defined in the proof of Theorem 12.

Let  $\mathbf{x}^T \in \mathcal{X}^T$  be any realization of features. Assume, w.l.o.g., that  $h_1$  is the underlying true function and  $y_t = h_1(\mathbf{x}_t)$  is the true label. We construct the predictor  $\hat{y}_t$  as follows:

1. If  $h_1(\mathbf{x}_t) = h_2(\mathbf{x}_t)$ , we predict  $\hat{y}_t = h_1(\mathbf{x}_t)$ ;
2. Let  $S^t \subset [t]$  be the subset of all  $r \in [t]$  satisfying  $h_1(\mathbf{x}_r) \neq h_2(\mathbf{x}_r)$ . Suppose at time step  $t$ , we have  $h_1(\mathbf{x}_t) \neq h_2(\mathbf{x}_t)$ . We compute the following quantity:

$$a_t = \sum_{r \in S^{t-1}} L^2(e_r, q_{h_1(\mathbf{x}_r)}^{\mathbf{x}_r}) - L^2(e_r, q_{h_2(\mathbf{x}_r)}^{\mathbf{x}_r}),$$

where  $e_r$  is the standard base in  $\mathbb{R}^M$  that assigns value 1 at position  $\tilde{y}_r$  and  $\tilde{y}_r$  is the noisy label. We now predict  $\hat{y}_t = h_1(\mathbf{x}_t)$  if  $a_t \leq 0$  and  $\hat{y}_t = h_2(\mathbf{x}_t)$  otherwise.

We next analyze the prediction performance of  $\hat{y}_t$ . Clearly we only make errors for time steps  $t \in S^T$ . Denote  $N_T = |S^T|$ , and we may assume, w.l.o.g., that  $S^T = \{1, 2, \dots, N_T\}$ , since a time step  $t \notin S^T$  does not affect the prediction performance. Let  $\tilde{p}_1, \dots, \tilde{p}_{N_T}$  be the noisy distributions selected by adversary. For any  $t \leq N_T$ , we denote  $X_t = L^2(e_t, q_{h_1(\mathbf{x}_t)}^{\mathbf{x}_t}) - L^2(e_t, q_{h_2(\mathbf{x}_t)}^{\mathbf{x}_t})$ . By Proposition 7, Lemma 13 and the fact that  $\tilde{p}_t \in \mathcal{Q}_{h_1(\mathbf{x}_t)}^{\mathbf{x}_t}$ , we have  $Y_t \stackrel{\text{def}}{=} \mathbb{E}_{\tilde{y}_t}[X_t] = L^2(\tilde{p}_t, q_{h_1(\mathbf{x}_t)}^{\mathbf{x}_t}) - L^2(\tilde{p}_t, q_{h_2(\mathbf{x}_t)}^{\mathbf{x}_t}) \leq -\gamma$ . Invoking Lemma 35 (second part) and denoting  $N_t = |S^{t-1}|$ , we have for any  $t \leq N_T$  and  $\epsilon > 0$  that

$$\Pr \left[ \frac{1}{t} \sum_{r=1}^t X_r < \frac{1}{t} \sum_{r=1}^t Y_r + \epsilon \right] \geq 1 - e^{-2t\epsilon^2}.$$

Since  $Y_t \leq -\gamma$  holds almost surely, taking  $\epsilon = \gamma$ , we find  $\Pr[a_t < 0] \geq 1 - e^{-2(t-1)\gamma^2}$ . Therefore, the probability that  $\hat{y}_t$  makes an error is upper bounded by  $e^{-2(t-1)\gamma^2}$ . By the sum of geometric series and simple approximation, we conclude that

$$\sum_{t=n}^{\infty} e^{-2(t-1)\gamma^2} \leq \frac{1}{4\gamma^2} e^{-2(n-1)\gamma^2}.$$

Taking  $n = \frac{\log(1/(4\delta\gamma^2))}{2\gamma^2}$ , we can make the RHS  $\leq \delta$ . This implies that, w.p.  $\geq 1 - \delta$ , the prediction  $\hat{y}_t$  makes *no* errors after step  $n$ , hence, the total number errors is upper bounded by  $\frac{\log(1/(4\delta\gamma^2))}{2\gamma^2}$ . The result now follows by Theorem 17 with  $C(\delta/(2K)) = \frac{\log(K/(2\delta\gamma^2))}{2\gamma^2}$ . ■

It is interesting to note that the bound in Theorem 18 is *independent* to both the size of label set  $\mathcal{Y}$  and the noisy observation set  $\tilde{\mathcal{Y}}$ , as well as the time horizon  $T$ .

**Remark 19** Note that, the risk bound in Theorem 18 is of order  $\frac{\log^2 K}{\gamma^2}$ , which differs by a  $\frac{\log K}{\gamma}$  factor compared to the expected minimax risk bound in Theorem 12. This is introduced by the union bound when applying our reduction to pairwise comparison and the application of Azuma's inequality. We leave it as an open problem to determine if this extra factor can be removed.

**Remark 20** As we demonstrated in Example 1 the robust online classification problem with two-function class can be viewed as a robust hypothesis testing problem, where the distribution at each step can vary arbitrarily but must be consistent with the same hypothesis function. Note that, the classical hypothesis testing problem can be viewed as the case with constant functions and the adversary selecting one distribution at the beginning and staying fixed for all time steps. It was demonstrated in Polyanskiy and Wu (2022, Chapter 32.2.1) via minimax theorem that the testing error decays as  $e^{-nH^2(\mathcal{Q}_0, \mathcal{Q}_1)}$ , where  $H^2(p, q)$  is the squared Hellinger divergence. We conjecture that such a guarantee may also hold for our changing distribution cases, and if so would result in tighter bounds than our  $L^2$  divergence result presented in Theorem 18.

### 3.3. Tighter Bounds for Specialized Settings via Log-loss

In this section, we consider a specific noisy kernel that can occur naturally in applications such as differential privacy. Let  $\mathcal{Y} = \tilde{\mathcal{Y}} = \{1, \dots, M\}$ . We denote by  $u$  the uniform distribution over  $\tilde{\mathcal{Y}}$ . For any  $\eta \geq 0$ , we define a homogeneous (i.e., independent of  $\mathbf{x}$ ) kernel:

$$\forall \mathbf{x} \in \mathcal{X}, y \in \mathcal{Y}, \mathcal{K}^\eta(\mathbf{x}, y) = \{(1 - \eta')e_y + \eta'u : \eta' \leq \eta\},$$

where  $e_y$  is defined as before, that is, it is 1 at position  $y$  and zero otherwise.

We now state our main result for this section:

**Theorem 21** Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be any finite class,  $\mathcal{P}$  be the class of all singleton distributions over  $\mathcal{X}^T$  and  $\mathcal{K}^\eta$  be as defined above with  $0 \leq \eta < 1$ . Then, the expected minimax risk is upper bounded by

$$\tilde{r}_T(\mathcal{H}, \mathcal{P}, \mathcal{K}^\eta) \leq \frac{\log |\mathcal{H}|}{(1 - \eta)^2/2}.$$

Moreover, the high probability minimax risk at confidence  $\delta > 0$  is upper bounded by

$$B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}^\eta) \leq \frac{\log |\mathcal{H}| + 2 \log(1/\delta)}{(1 - \eta)^2/4}.$$

Furthermore, for  $1 - \eta \ll \frac{1}{M}$  we have  $B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}^\eta) \leq O\left(\frac{\log |\mathcal{H}| + \log(1/\delta)}{M(1 - \eta)^2}\right)$ .

**Proof** Our proof follows a similar path as the proof of Theorem 12. For any  $h \in \mathcal{H}$ , we define a distribution-valued function  $f_h$  such that  $f_h(\mathbf{x}) = (1 - \eta)e_{h(\mathbf{x})} + \eta u$ . Let  $\mathcal{F} = \{f_h : h \in \mathcal{H}\}$ . Invoking Lemma 9 and using the fact the KL-divergence is Bregman and 1-Exp-concave, there exists a prediction rule  $\hat{p}^T$  such that

$$\sup_{\mathbf{x}^T \in \mathcal{X}^T, f \in \mathcal{F}, \tilde{p}^T \in \mathcal{D}(\tilde{\mathcal{Y}})^T} \mathbb{E}_{\tilde{y}^T} \left[ \sum_{t=1}^T \text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f(\mathbf{x}_t)) \right] \leq \log |\mathcal{H}|,$$

where  $\tilde{y}_t \sim \tilde{p}_t$  and  $\tilde{p}^T$  is any noisy distribution. We now define the following predictor:

$$\hat{y}_t = \arg \max_y \{\hat{p}_t[y] : y \in \mathcal{Y}\}.$$

Let now  $h^* \in \mathcal{H}$  be the underlying true classification function and  $\tilde{p}^T$  be the noisy distribution selected by the adversary. We claim that:

**Claim 1** *The following holds for all  $t \leq T$ :*

$$\text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \geq 0.$$

Moreover, if  $\hat{y}_t \neq h^*(\mathbf{x}_t)$  then

$$\text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \geq (1 - \eta)^2/2.$$

**Proof** [Proof of the Claim] Let  $y_t = h^*(\mathbf{x}_t)$  and  $e_t \in \mathcal{D}(\tilde{\mathcal{Y}})$  be the distribution that assigns probability 1 on  $y_t$ . By the definition  $f_{h^*}(\mathbf{x}_t) = \lambda e_t + (1 - \lambda)u$  and  $\tilde{p}_t = \lambda_t e_t + (1 - \lambda_t)u$ , where  $\lambda = 1 - \eta$  and  $\lambda_t = 1 - \eta_t$  for some  $\eta_t \leq \eta$ . Since  $0 \leq \eta_t \leq \eta$ , we have  $1 \geq \lambda_t \geq \lambda$ . Note that,  $\text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t))$  is a linear function w.r.t.  $\lambda_t$  (Proposition 7), and it takes the minimal value at  $\lambda_t \in \{1, \lambda\}$  and therefore

$$\text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \geq \min\{\log(f_{h^*}(\mathbf{x}_t)[y_t]/\hat{p}_t[y_t]), \text{KL}(f_{h^*}(\mathbf{x}_t), \hat{p}_t)\}.$$

Clearly, the second KL-divergence term is positive. We now show that  $\log(f_{h^*}(\mathbf{x}_t)[y_t]/\hat{p}_t[y_t]) \geq 0$ . To see this, we have by Lemma 9 that  $\hat{p}_t$  is a convex combination of  $\{f(\mathbf{x}_t) : f \in \mathcal{F}\}$  and therefore  $\hat{p}_t = \lambda a_t + (1 - \lambda)u$  for some  $a_t \in \mathcal{D}(\tilde{\mathcal{Y}})$ . This implies that  $\hat{p}_t[y_t] = \lambda a_t[y_t] + (1 - \lambda)\frac{1}{M}$  and  $f_{h^*}(\mathbf{x}_t)[y_t] = \lambda + (1 - \lambda)\frac{1}{M}$ . Since  $a_t[y_t] \leq 1$ , we have  $f_{h^*}(\mathbf{x}_t)[y_t] \geq \hat{p}_t[y_t]$ . The first part of the claim now follows.

We now prove the second part of the claim. Note that in order for  $\hat{y}_t \neq y_t$  we must have  $a_t[y_t] \leq \frac{1}{2}$ , since  $\hat{y}_t$  is defined to be the label with maximum probability mass under  $\hat{p}_t$ . Therefore,

$$\log(f_{h^*}(\mathbf{x}_t)[y_t]/\hat{p}_t[y_t]) \geq \log\left(\frac{\lambda + (1 - \lambda)/M}{\lambda/2 + (1 - \lambda)/M}\right) = \log\left(1 + \frac{\lambda/2}{\lambda/2 + (1 - \lambda)/M}\right) \geq \log(1 + \lambda)$$

where the second inequality follows from  $\lambda/2 + (1 - \lambda)/M \leq 1/2$ . Furthermore, we have

$$\text{KL}(f_{h^*}(\mathbf{x}_t), \hat{p}_t) \geq \frac{1}{2} \|f_{h^*}(\mathbf{x}_t) - \hat{p}_t\|_1^2 \geq \lambda^2/2,$$

where the first inequality is a consequence of Pinsker's inequality (Polyanskiy and Wu, 2022). The claim now follows by noting that  $\log(1 + \lambda) \geq \lambda^2/2$  for all  $0 \leq \lambda \leq 1$ . ■

The first part of the theorem now follows by the same argument as in the proof of Theorem 12. We defer the proof of the second and third parts to Appendix D. ■

**Remark 22** Taking  $M = 2$  and  $\eta = 2\gamma$ , our setup in Theorem 21 recovers the Bounded Stochastic Noise setting as in Ben-David et al. (2009). It was shown in Ben-David et al. (2009, Theorem 15) that the expected minimax risk is upper bounded by  $\frac{\log |\mathcal{H}|}{1 - 2\sqrt{\gamma(1 - \gamma)}}$ , using a non-intuitive backward induction. This has the same asymptotic rate w.r.t.  $\gamma$  compared to our  $\frac{\log |\mathcal{H}|}{(1 - 2\gamma)^{2/2}}$  bound when  $\gamma \rightarrow \frac{1}{2}$ . Our approach is more intuitive and resolves the multi-label case in a unified way. We will show in Section 3.5 that the  $O\left(\frac{\log |\mathcal{H}|}{M(1 - \eta)^2}\right)$  bound is actually tight for  $\eta \rightarrow 1$ .

**Tsybakov type noise:** We now deal with the *Tsybakov type noise* as introduced in Example 2. Recall that we take  $|\mathcal{Y}| = |\tilde{\mathcal{Y}}| = 2$ . For any  $\mathbf{x}^T$ , the Tsybakov type noise satisfies

$$\mathcal{K}(\mathbf{x}_t, y) = \{(1 - \eta_t)e_y + \eta_t u\},$$

subject to the condition that for some  $A > 0$  and  $0 \leq \alpha < 1$

$$\forall r \in (0, 1/2], \frac{1}{T} \sum_{t=1}^T 1 \left\{ \frac{\eta_t}{2} \geq \frac{1}{2} - r \right\} \leq Ar^{\frac{\alpha}{1-\alpha}}. \quad (18)$$

Slightly departing from our previous definition of noisy kernels, we *do not* assume the parameter  $\eta_t$  to be known to the learner. We now state our main result for this part:

**Theorem 23** *Let  $\mathcal{H} \subset \{0, 1\}^{\mathcal{X}}$  be any finite class,  $\mathcal{P}$  be the class of all singleton distributions over  $\mathcal{X}^T$  and  $\mathcal{K}$  be the Tsybakov type noise kernel with parameter  $A > 0, 0 \leq \alpha < 1$ . Then the expected minimax risk for the robust online classification problem is upper bounded by*

$$\tilde{r}_T(\mathcal{H}, \mathcal{P}, \mathcal{K}) \leq O(\log |\mathcal{H}| \cdot T^{\frac{2-2\alpha}{\alpha}}).$$

**Proof** [Sketch of Proof] We only sketch the proof as it essentially follows the proof of Theorem 21. Let  $\eta$  be determined later. Define for any  $h \in \mathcal{H}$ , a function  $f_h$  such that  $f_h(\mathbf{x}) = (1 - \eta)e_{h(\mathbf{x})} + \eta u$  and  $\mathcal{F} = \{f_h : h \in \mathcal{H}\}$ . Let  $\hat{p}_t$  be the prediction rule derived from Lemma 9 with KL-divergence. By the proof of Theorem 21 we have

$$\sum_{t=1}^T \text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \leq \log |\mathcal{H}|,$$

where  $\tilde{p}_t = (1 - \eta_t)e_{h^*(\mathbf{x}_t)} + \eta_t u$  and  $h^*$  is the underlying true function. We still define  $\hat{y}_t = \arg \max_y \{\hat{p}_t[y] : y \in \mathcal{Y}\}$ . We now observe that, for any  $\eta_t \leq \eta$ , Claim 1 still holds. Denote by  $S \subset [T]$  the time steps  $t$  such that  $\eta_t \leq \eta$ . For the time step  $t \notin S$ , we can bound

$$\text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) \geq -\log(2/\eta),$$

since  $f_h(\mathbf{x})[y], \hat{p}_t[y] \geq \eta/2$  for all  $y \in \{0, 1\}$  by construction. By condition (18), we have

$$|\bar{S}| \leq AT\gamma^{\frac{\alpha}{1-\alpha}},$$

where  $\gamma = (1 - \eta)/2$  and  $\bar{S}$  is the complementary set of  $S$ . Therefore, by the argument in the proof of Theorem 12 restricted to  $S$ , we have the expected minimax risk upper bounded by

$$\inf_{0 \leq \gamma < 1/2} AT\gamma^{\frac{\alpha}{1-\alpha}} + \frac{\log(2/(1 - 2\gamma))AT\gamma^{\frac{\alpha}{1-\alpha}} + \log |\mathcal{H}|}{2\gamma^2} \leq O(\log |\mathcal{H}| T^{\frac{2-2\alpha}{\alpha}})$$

by taking  $\gamma = \left(\frac{\log |\mathcal{H}|}{T}\right)^{\frac{1-\alpha}{\alpha}}$ . This follows from the fact that

$$\begin{aligned} \sum_{t \in S} \text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) &\leq - \sum_{t \in \bar{S}} \text{KL}(\tilde{p}_t, \hat{p}_t) - \text{KL}(\tilde{p}_t, f_{h^*}(\mathbf{x}_t)) + \log |\mathcal{H}| \\ &\leq |\bar{S}| \log(2/\eta) + \log |\mathcal{H}|, \end{aligned}$$

and Claim 1 holds for any time step  $t \in S$ . ■

**Remark 24** Note that Theorem 23 achieves sub-linear risk if  $\alpha \geq \frac{2}{3}$ . For instance, if  $\alpha = 0.7$ , our result gives an upper bound of order  $\tilde{O}(T^{0.86})$ . It is worth noting that, if  $\eta_t$  is assumed to be known, then we do not need to pay the extra penalty  $\log(2/\eta)$  for a time step  $t \notin S$ , since we can simply ignore such steps when constructing the predictor  $\hat{p}_t$ . The derived risk bound will be of order  $\tilde{O}(T^{\frac{2(1-\alpha)}{2-\alpha}})$ . This is tight for the worst class  $\mathcal{H}$ , see Remark 34. We leave it as an open problem to obtain the optimal bound for the unknown  $\eta_t$  cases.

### 3.4. Stochastic Feature Generation Process, Infinite Class, and Covering

We have demonstrated in the previous section that the minimax risk of our robust online classification problem can be effectively bounded for a finite hypothesis class  $\mathcal{H}$  and singleton feature generating distributions over  $\mathcal{X}^T$ . We now demonstrate how such result can be generalized to infinite classes and general feature generating processes via suitable covering of the class.

We first introduce the following notion of covering from Wu et al. (2023a), which generalizes a similar concept in Ben-David et al. (2009).

**Definition 25** Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be any hypothesis class and  $\mathbf{P}$  be any class of random processes over  $\mathcal{X}^T$ . We say a class of functions  $\mathcal{G} \subset \mathcal{Y}^{\mathcal{X}^*}$  (where  $\mathcal{X}^*$  is a set of all finite sequences of  $\mathcal{X}$ ) is a stochastic sequential covering of  $\mathcal{H}$  w.r.t.  $\mathbf{P}$  at scale 0 and confidence  $\delta$  if

$$\forall \nu^T \in \mathbf{P}, \Pr_{\mathbf{x}^T} [\exists h \in \mathcal{H} \forall g \in \mathcal{G} \exists t \in [T], h(\mathbf{x}_t) \neq g(\mathbf{x}^t)] \leq \delta.$$

We now have the following result that reduces the minimax risk of an infinite class to the size of the stochastic sequential cover.

**Theorem 26** Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be any hypothesis class,  $\mathbf{P}$  be any class of random processes over  $\mathcal{X}^T$  and  $\mathcal{K}$  be a noisy kernel that is well-separated w.r.t.  $L^2$  divergence at scale  $\gamma$ . If there exists a finite stochastic sequential cover  $\mathcal{G} \subset \mathcal{Y}^{\mathcal{X}^*}$  of  $\mathcal{H}$  w.r.t.  $\mathbf{P}$  at scale 0 and confidence  $\delta/2 > 0$ , then the high probability minimax risk at confidence  $\delta$  of problem  $(\mathcal{H}, \mathbf{P}, \mathcal{X})$  is upper bounded by

$$B^\delta(\mathcal{H}, \mathbf{P}, \mathcal{K}) \leq O\left(\frac{\log(|\mathcal{G}|) \log(|\mathcal{G}|/4\delta\gamma^2)}{\gamma^2}\right).$$

**Proof** Let  $A$  be the event over  $\mathbf{x}^T$  so that  $\forall h \in \mathcal{H}, \exists g \in \mathcal{G}$  such that  $\forall t \in [T], h(\mathbf{x}_t) = g(\mathbf{x}^t)$ . Let now  $\nu^T \in \mathbf{P}$  be the underlying true feature generating process. We have by the definition of stochastic sequential covering that  $\Pr_{\mathbf{x}^T}[A] \geq 1 - \delta/2$ . We now observe that Theorem 18 holds for sequential functions as well (simply replace any occurrence of  $h(\mathbf{x}_t)$  with  $g(\mathbf{x}^t)$  in the proof). Therefore, taking confidence parameter  $\delta/2$ , the prediction rule derived from Theorem 18 yields high probability minimax risk of problem  $(\mathcal{G}, \mathbf{P}, \mathcal{X})$ , which is upper bounded by

$$B^{\delta/2}(\mathcal{G}, \mathbf{P}, \mathcal{K}) \leq O\left(\frac{\log(|\mathcal{G}|) \log(|\mathcal{G}|/4\delta\gamma^2)}{\gamma^2}\right). \quad (19)$$

Let  $h^* \in \mathcal{H}$  be the underlying true function,  $\mathbf{x}^T \in A$  be any realization of the feature, and  $g^*$  be the sequential covering function of  $h^*$  at scale 0. Note that,  $g^*$  has the same labeling as  $h^*$  on  $\mathbf{x}^T$ . Therefore, any predictor has the same behaviours when running on  $h^*$  and  $g^*$ , and thus the high probability minimax risk of problem  $(\mathcal{H}, \mathbf{P}, \mathcal{X})$  at confidence  $\delta/2$  conditioning on event  $A$  is upper bounded by (19). The theorem now follows by a union bound.  $\blacksquare$

**Remark 27** Note that, any bounds that we have established in the previous sections for finite class can be extended to the infinite classes; these bounds depend only on the stochastic sequential cover size using a similar argument as in Theorem 26. We will not discuss all such cases in this paper in the interest of clarity of presentation.

**Corollary 28** Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be a class with finite Littlestone dimension  $\text{Ldim}(\mathcal{H})$  (Daniely et al., 2015) and  $|\mathcal{Y}| = N$ ,  $\mathcal{P}$  be the class of all singleton distributions, and  $\mathcal{K}$  be any noisy kernel that is well-separated w.r.t.  $L^2$  divergence at scale  $\gamma$ . Then, with high probability the minimax risk at confidence  $\delta$  of problem  $(\mathcal{H}, \mathcal{P}, \mathcal{X})$  is upper bounded by

$$B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}) \leq O\left(\frac{\text{Ldim}(\mathcal{H})^2 \log^2(TN) + \text{Ldim}(\mathcal{H}) \log(TN/4\delta\gamma^2)}{\gamma^2}\right).$$

Moreover, for the noisy kernel  $\mathcal{K}^\eta$  as in Theorem 21, the high probability minimax risk with confidence  $\delta > 0$  is upper bounded by

$$B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}^\eta) \leq \frac{(\text{Ldim}(\mathcal{H}) + 1) \log(TN) + 2 \log(1/\delta)}{(1 - \eta)^2/4}.$$

**Proof** The first part follows directly from Theorem 26 and the fact that the sequential covering of  $\mathcal{H}$  w.r.t. the singleton distributions over  $\mathcal{X}^T$  is of order  $(TN)^{\text{Ldim}(\mathcal{H})+1}$  by (Daniely et al., 2015, Theorem 25). The second part follows by Theorem 21.  $\blacksquare$

**Approximate Covering With Small Errors:** In our Definition 25 of the stochastic sequential covering, we assumed that the function  $h \in \mathcal{H}$  is *exactly* covered over  $\mathbf{x}^T$  with high probability. This may sometimes be too strong. One possible way of dealing with this issue is to consider the following *approximate covering* notion:

$$\forall \nu^T \in \mathcal{P}, \Pr_{\mathbf{x}^T} \left[ \exists h \in \mathcal{H} \forall g \in \mathcal{G}, \sum_{t=1}^T 1\{h(\mathbf{x}_t) \neq g(\mathbf{x}_t)\} \geq D \right] \leq \delta,$$

for some error bound  $D$ . We note that such a relaxation does not change our narrative substantially, since one can always expand any covering functions  $g \in \mathcal{G}$  to a set of functions  $\{g_I : I \subset [T] \text{ and } |I| \leq D, \{y_r\}_{r \in I} \in \mathcal{Y}^I\}$ , where

$$g_I(\mathbf{x}^t) = \begin{cases} g(\mathbf{x}^t), & \text{if } t \notin I, \\ y_r, & \text{if } t = r \in I \end{cases}.$$

The class  $\mathcal{G}'$  containing all such  $g_I$ s forms an *exact* sequential cover of  $\mathcal{H}$  and  $|\mathcal{G}'| \leq (TN)^{(D+1)}|\mathcal{G}|$ .

Finally, we generalize our results for a large class of distribution over  $\mathcal{X}$  known as  $\sigma$ -smooth distributions. For any given distribution  $\mu$  over  $\mathcal{X}$ , we say a distribution  $\nu$  over  $\mathcal{X}$  is  $\sigma$ -smooth w.r.t.  $\mu$  if for all measurable sets  $A \subset \mathcal{X}$ , we have  $\nu(A) \leq \mu(A)/\sigma$  (Haghtalab et al., 2020). A random process  $\nu^T$  over  $\mathcal{X}^T$  is said to be  $\sigma$ -smooth if the *conditional marginal*  $\nu^T(\cdot \mid X^{t-1})$  is  $\sigma$ -smooth w.r.t.  $\mu$  for all  $t \leq T$ , almost surely.

**Corollary 29** *Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be a class with finite VC-dimension  $\text{VC}(\mathcal{H})$  and  $|\mathcal{Y}| = 2$ ,  $S^\sigma(\mu)$  be the class of all  $\sigma$ -smooth processes w.r.t.  $\mu$ , and  $\mathcal{K}^\eta$  be the Bernoulli noisy kernel as in Theorem 21. Then the high probability minimax risk at confidence  $\delta > 0$  is upper bounded by*

$$B^\delta(\mathcal{H}, S^\sigma(\mu), \mathcal{K}) \leq O\left(\frac{\text{VC}(\mathcal{H}) \log(T/\sigma) + \log(1/\delta)}{(1-\eta)^2}\right).$$

**Proof** By (Wu et al., 2023a, Proposition 22),  $\mathcal{H}$  admits an approximate stochastic cover  $\mathcal{G}$  at confidence  $\delta/2 > 0$  and error  $D = O(\text{VC}(\mathcal{H}) + \log(1/\delta)/\log T)$  such that

$$\log |\mathcal{G}| \leq O(\text{VC}(\mathcal{H})(\log(T/\sigma) + \log \log(T/\delta))).$$

Let  $\mathcal{G}'$  be defined as above where  $\log |\mathcal{G}'| \leq O(D \log T + \log |\mathcal{G}|)$ . We know that  $\log |\mathcal{G}'| \leq O(\text{VC}(\mathcal{H}) \log(T/\sigma) + \log(1/\delta))$ . We now condition on the event of the exact covering. By Theorem 21 (second part), the high probability minimax risk at confidence  $\delta/2$  is upper bounded by

$$B^{\delta/2}(\mathcal{H}, S^\sigma(\mu), \mathcal{K}^\eta) \leq O\left(\frac{\log |\mathcal{G}'| + \log(2/\delta)}{(1-\eta)^2}\right) \leq O\left(\frac{\text{VC}(\mathcal{H}) \log(T/\sigma) + \log(1/\delta)}{(1-\eta)^2}\right).$$

The result now follows by a union bound. ■

**Remark 30** We refer to Wu et al. (2023a) for more results on the stochastic sequential covering estimates of various distribution classes.

**Remark 31** Note that, the case when the underlying true labels  $y^T$  are not realizable by  $\mathcal{H}$  but differ by a small error  $D$  can also be resolved using the same approach as above. This will introduce an additive  $D \log(TN)$  term in the derived minimax risk bounds. However, in the more general case where the noisy samples for which the differing positions do not even follow the prescribed noisy kernel  $\mathcal{K}$ , this approach will not apply. This can, however, be resolved by using a similar argument as in the proof of Theorem 23 by introducing a penalty at each error position.

### 3.5. A Lower Bound Based on Hellinger Gap

We now prove a general lower bound based on the squared Hellinger gap, which provides matching lower bounds for the settings in Section 3.3. Recall that for any  $p, q \in \mathcal{D}(\hat{\mathcal{Y}})$ , the squared Hellinger divergence is defined as  $H^2(p, q) = \sum_{\tilde{y} \in \hat{\mathcal{Y}}} (\sqrt{p[\tilde{y}]} - \sqrt{q[\tilde{y}]})^2$ .

**Proposition 32** *For any noisy kernel  $\mathcal{K}$  such that  $\forall \mathbf{x} \in \mathcal{X}$  there exists  $y \neq y' \in \mathcal{Y}$  such that  $\inf_{p \in \mathcal{Q}_{\tilde{y}}^{\mathbf{x}}, q \in \mathcal{Q}_{y'}^{\mathbf{x}}} \{H^2(p, q)\} \leq \gamma$  and  $K \geq 1$ . If  $|\mathcal{X}| \geq \log K$ , then there exists a hypothesis class  $\mathcal{H}$  with  $|\mathcal{H}| = K$  such that for  $\mathbf{P}$  being the singleton process over  $\mathcal{X}^T$ , we have the expected minimax risk lower bounded by*

$$\tilde{r}_T(\mathcal{H}, \mathbf{P}, \mathcal{K}) \geq \Omega\left(\frac{\log |\mathcal{H}|}{\gamma}\right).$$

**Proof** Denote  $\tau = \log K$  with  $K = |\mathcal{H}|$ , and  $\mathbf{x}_1, \dots, \mathbf{x}_\tau$  be  $\tau$  distinct elements in  $\mathcal{X}$ . We define for any  $\mathbf{b} \in \{0, 1\}^\tau$  a function  $h_{\mathbf{b}}$  such that for all  $i \in [\tau]$ ,  $h_{\mathbf{b}}(\mathbf{x}_i) = y_i$  if  $\mathbf{b}[i] = 0$  and  $h_{\mathbf{b}}(\mathbf{x}_i) = y'_i$  otherwise, where  $y_i \neq y'_i \in \mathcal{Y}$  are the elements that satisfy  $\inf_{p \in \mathcal{Q}_{y_i}^{\mathbf{x}_i}, q \in \mathcal{Q}_{y'_i}^{\mathbf{x}_i}} \{H^2(p, q)\} \leq \gamma$ . Let

$\mathcal{H}$  be the class consisting of all such  $h_{\mathbf{b}}$ . Let  $q_i \in \mathcal{Q}_{y_i}^{\mathbf{x}_i}$  and  $q'_i \in \mathcal{Q}_{y'_i}^{\mathbf{x}_i}$  be the elements satisfying  $H^2(q_i, q'_i) \leq \gamma$ . We now partition the features  $\mathbf{x}^T$  into  $\tau$  epochs, each of length  $T/\tau$ , such that each epoch  $i$  has constant feature  $\mathbf{x}_i$ . Let  $\mathbf{h}$  be a random function selected uniformly from  $\mathcal{H}$ . We claim that for any prediction rule  $\hat{y}_t$  and any epoch  $i$  we have

$$\mathbb{E}_{\mathbf{h}, \tilde{y}^T} \left[ \sum_{t=iT/\tau-1}^{(i+1)T/\tau} 1\{\mathbf{h}(\mathbf{x}_t) \neq \hat{y}_t\} \right] \geq \Omega\left(\frac{1}{\gamma}\right), \quad (20)$$

where  $\tilde{y}_t \sim q_i$  if  $\mathbf{h}(\mathbf{x}_i) = y_i$  and  $\tilde{y}_t \sim q'_i$  otherwise. The proposition now follows by counting the errors for all  $\tau$  epochs.

We now establish (20) using the Le Cam's two point method. Clearly, for each epoch  $i$ , the prediction performance depends only on the label  $\mathbf{y}_i = \mathbf{h}(\mathbf{x}_i)$ , which is uniform over  $\{y_i, y'_i\}$  and independent for different epochs by construction. For any time step  $j$  during the  $i$ th epoch, we denote by  $\tilde{y}^{j-1}$  and  $\tilde{y}'^{j-1}$  the samples generated from  $q_i$  and  $q'_i$ , respectively. By the Le Cam's two point method (Polyanskiy and Wu, 2022, Theorem 7.7) the expected error at step  $j$  is lower bounded by

$$\frac{1 - \text{TV}(\tilde{y}^{j-1}, \tilde{y}'^{j-1})}{2} \geq \frac{1 - \sqrt{H^2(\tilde{y}^{j-1}, \tilde{y}'^{j-1})(1 - H^2(\tilde{y}^{j-1}, \tilde{y}'^{j-1})/4)}}{2} \quad (21)$$

where the inequality follows from (Polyanskiy and Wu, 2022, Equation 7.20). Note that the RHS of (21) is *monotone decreasing* w.r.t.  $H^2(\tilde{y}^{j-1}, \tilde{y}'^{j-1})$ , since  $H^2(p, q) \leq 2$  for all  $p, q$ .

By the *tensorization* of Hellinger divergence (Polyanskiy and Wu, 2022, Equation 7.23), we have

$$H^2(\tilde{y}^{j-1}, \tilde{y}'^{j-1}) = 2 - 2(1 - H^2(q_i, q'_i)/2)^{j-1} \leq 2 - 2(1 - \gamma/2)^{j-1},$$

where the last inequality is implied by  $H^2(q_i, q'_i) \leq \gamma$ . Using the fact  $\log(1 - x) \geq \frac{-x}{1-x}$ , we have if  $\gamma \leq 1$  and  $j-1 \leq \frac{1}{\gamma}$  then  $2 - 2(1 - \gamma/2)^{j-1} \leq 2(1 - e^{-1}) < 2$ . Therefore, the RHS of (21) is lower bounded by an *absolute* positive constant for all  $j-1 \leq \frac{1}{\gamma}$ , and hence the expected cumulative error will be lower bounded by  $\Omega(1/\gamma)$  during epoch  $i$ . This completes the proof.  $\blacksquare$

**Remark 33** Specializing to the setting in Theorem 21, we know that the squared Hellinger gap is of order

$$\left( \sqrt{\frac{\eta}{M}} - \sqrt{1 - \frac{(M-1)\eta}{M}} \right)^2 \sim \frac{M(1-\eta)^2}{4},$$

when  $\eta \rightarrow 1$  (by Taylor expansion). Proposition 32 implies an  $\Omega\left(\frac{\log |\mathcal{H}|}{M(1-\eta)^2}\right)$  lower bound. This is a matching lower bound for Theorem 21 (third part) when  $1 - \eta \ll \frac{1}{M}$ .

**Remark 34** Note that for the Tsybakov type noise as in Theorem 23 we can derive a lower bound of order  $T^{\frac{2(1-\alpha)}{2-\alpha}}$ , matching the upper bound of the case when the parameters  $\eta_t$  are known. To see this, we select  $AT\gamma^{\frac{\alpha}{1-\alpha}}$  time steps with  $\eta_t = \frac{1}{2} - \gamma$ . It is easy to verify that the squared Hellinger divergence for Bernoulli distributions with parameter  $\eta_t$  and  $1 - \eta_t$  is of order  $\gamma^2$ . Therefore, by Proposition 32, if  $AT\gamma^{\frac{\alpha}{1-\alpha}} \geq \Omega\left(\frac{\log |\mathcal{H}|}{\gamma^2}\right)$  then the expected minimax risk will be lower bounded by  $\Omega\left(\frac{\log |\mathcal{H}|}{\gamma^2}\right)$ , which happens when  $\gamma$  is of order  $T^{-\frac{1-\alpha}{2-\alpha}}$ .

## References

- Shai Ben-David, Dávid Pál, and Shai Shalev-Shwartz. Agnostic online learning. In *Conference on Learning Theory*, volume 3, 2009.
- Alankrita Bhatt and Young-Han Kim. Sequential prediction under log-loss with side information. In *Algorithmic Learning Theory*, pages 340–344. PMLR, 2021.
- Blair Bilodeau, Dylan Foster, and Daniel Roy. Tight bounds on minimax regret under logarithmic loss via self-concordance. In *International Conference on Machine Learning*, pages 919–929. PMLR, 2020.
- Blair Bilodeau, Dylan J Foster, and Daniel M Roy. Minimax rates for conditional density estimation via empirical entropy. *arXiv preprint arXiv:2109.10461*, 2021.
- N. Cesa-Bianchi and G. Lugosi. *Prediction, Learning and Games*. Cambridge University Press, 2006.
- Amit Daniely, Sivan Sabato, Shai Ben-David, and Shai Shalev-Shwartz. Multiclass learnability and the erm principle. *J. Mach. Learn. Res.*, 16(1):2377–2404, 2015.
- Sam Efromovich. Conditional density estimation in a regression setting. *The Annals of Statistics*, 35:2504–2535, 2007.
- Dylan J Foster, Sham M Kakade, Jian Qian, and Alexander Rakhlin. The statistical complexity of interactive decision making. *arXiv preprint arXiv:2112.13487*, 2021.
- Peter D Grünwald and Nishant A Mehta. Fast rates for general unbounded loss functions: from erm to generalized bayes. *The Journal of Machine Learning Research*, 21(1):2040–2119, 2020.
- Nika Haghtalab, Tim Roughgarden, and Abhishek Shetty. Smoothed analysis of online and differentially private learning. *Advances in Neural Information Processing Systems*, 33:9203–9215, 2020.
- Elad Hazan et al. Introduction to online convex optimization. *Foundations and Trends® in Optimization*, 2(3-4):157–325, 2016.
- Sham Kakade and Adam T Kalai. From batch to transductive online learning. *Advances in Neural Information Processing Systems*, 18, 2005.
- Nick Littlestone. Learning quickly when irrelevant attributes abound: A new linear-threshold algorithm. *Machine learning*, 2:285–318, 1988.
- Nagarajan Natarajan, Inderjit S Dhillon, Pradeep K Ravikumar, and Ambuj Tewari. Learning with noisy labels. *Advances in neural information processing systems*, 26, 2013.
- Yury Polyanskiy and Yihong Wu. *Information Theory: From Coding to Learning*. Cambridge University Press, 2022.
- Alexander Rakhlin and Karthik Sridharan. Sequential probability assignment with binary alphabets and large classes of experts. *arXiv preprint arXiv:1501.07340*, 2015.

- Alexander Rakhlin, Karthik Sridharan, and Ambuj Tewari. Online learning: Random averages, combinatorial parameters, and learnability. In *Advances in Neural Information Processing Systems*, 2010.
- Changlong Wu, Mohsen Heidari, Ananth Grama, and Wojciech Szpankowski. Expected worst case regret via stochastic sequential covering. *arXiv preprint arXiv:2209.04417*, 2022a.
- Changlong Wu, Mohsen Heidari, Ananth Grama, and Wojciech Szpankowski. Precise regret bounds for log-loss via a truncated bayesian algorithm. In *Advances in Neural Information Processing Systems*, volume 35, pages 26903–26914, 2022b.
- Changlong Wu, Ananth Grama, and Wojciech Szpankowski. Online learning in dynamically changing environments. In *Conference on Learning Theory*, pages 325–358. PMLR 195, 2023a.
- Changlong Wu, Yifan Wang, Ananth Grama, and Wojciech Szpankowski. Learning functional distributions with private labels. In *International Conference on Machine Learning (ICML)*, volume 202 of *PMLR*, pages 37728–37744. PMLR, 23–29 Jul 2023b.
- Tong Zhang. *Mathematical analysis of machine learning algorithms*. Cambridge University Press, 2023.

## Appendix A. Martingale concentration inequalities

In this appendix, we present some standard concentration results for martingales, which will be useful for deriving high probability guarantees. We refer to (Zhang, 2023, Chapter 13.1) for the proofs.

**Lemma 35 (Azuma’s Inequality)** *Let  $X_1, \dots, X_T$  be an arbitrary random process adaptive to some filtration  $\{\mathcal{F}_t\}_{t \leq T}$  such that  $|X_t| \leq M$  for all  $t \leq T$ . Let  $Y_t = \mathbb{E}[X_t \mid \mathcal{F}_{t-1}]$  be the conditional expected random variable of  $X_t$ . Then for all  $\delta > 0$ , we have*

$$\Pr \left[ \sum_{t=1}^T Y_t < \sum_{t=1}^T X_t + M \sqrt{(T/2) \log(1/\delta)} \right] \geq 1 - \delta,$$

and

$$\Pr \left[ \sum_{t=1}^T Y_t > \sum_{t=1}^T X_t - M \sqrt{(T/2) \log(1/\delta)} \right] \geq 1 - \delta.$$

The following lemma provides a tighter concentration when  $X_t \geq 0$ , which can be viewed as an Martingale version of the multiplicative Chernoff bound.

**Lemma 36 ((Zhang, 2023, Theorem 13.5))** *Let  $X_1, \dots, X_T$  be an arbitrary random process adaptive to some filtration  $\{\mathcal{F}_t\}_{t \leq T}$  such that  $0 \leq X_t \leq M$  for all  $t \leq T$ . Let  $Y_t = \mathbb{E}[X_t \mid \mathcal{F}_{t-1}]$  be the conditional expected random variable of  $X_t$ . Then for all  $\delta > 0$  we have*

$$\Pr \left[ \sum_{t=1}^T Y_t < 2 \sum_{t=1}^T X_t + 2M \log(1/\delta) \right] \geq 1 - \delta,$$

and

$$\Pr \left[ \sum_{t=1}^T Y_t > \frac{1}{2} \sum_{t=1}^T X_t - (M/2) \log(1/\delta) \right] \geq 1 - \delta.$$

**Proof** Applying [Zhang \(2023, Theorem 13.5\)](#) with  $\xi_t = X_t/M$  and  $\lambda = 1$  in the theorem.  $\blacksquare$

**Remark 37** It should be noted that the assumption  $X_t \geq 0$  is required for [Lemma 36](#) to hold. To see this, we group  $X^T$  as  $X_1X_2, X_3X_4, \dots$  such that  $X_{2t-1}$  is uniform over  $\{-1, 1\}$  and  $X_{2t} = -X_{2t-1}$  for all  $t \in [T]$ . It is easy to verify that  $X_1 + \dots + X_T = 0$  almost surely. But  $Y_{2t-1} = 0$  and  $Y_{2t} = -X_{2t-1}$ , hence, we have  $Y_1 + \dots + Y_T$  is sum of  $T/2$  independent uniform distributions over  $\{-1, 1\}$ . Therefore, by the central limit theorem  $Y_1 + \dots + Y_T \geq \Omega(\sqrt{T})$  with constant probability. This, unfortunately, limits its application to random variables of form  $L(e_t, \hat{p}_t) - L(e_t, f(\mathbf{x}_t))$ . There are, however, special cases such as for log-loss in the realizable case that a tight concentration holds for Hellinger divergence, see e.g., [Theorem 41](#).

## Appendix B. Exponential Weighted Average under Exp-concave losses

We now introduce the *Exponential Weighted Average (EWA)* algorithm and its regret analysis under the Exp-concave losses, which is mostly standard ([Cesa-Bianchi and Lugosi, 2006](#), Chapter 3.3) and we include it here for completeness. Let  $\mathcal{F} = \{f_1, \dots, f_K\} \subset \mathcal{D}(\tilde{\mathcal{Y}})^{\mathcal{X}}$  be a  $\mathcal{D}(\tilde{\mathcal{Y}})$ -valued function class of size  $K$  and  $\ell : \tilde{\mathcal{Y}} \times \mathcal{D}(\tilde{\mathcal{Y}}) \rightarrow \mathbb{R}^{\geq 0}$  be an  $\alpha$ -Exp-concave loss (see definition in [Section 2.1](#)). The EWA algorithm is presented in [Algorithm 2](#).

---

### Algorithm 2 Exponential Weighted Average (EWA) predictor

---

**Input:** Class  $\mathcal{F} = \{f_1, \dots, f_K\}$  and  $\alpha$ -Exp-concave loss  $\ell$

Set  $\mathbf{w}^1 = \{1, \dots, 1\} \in \mathbb{R}^K$ ;

**for**  $t = 1, \dots, T$  **do**

    Receive  $\mathbf{x}_t$ ;

    Make prediction:

$$\hat{p}_t = \frac{\sum_{k=1}^K \mathbf{w}^t[k] f_k(\mathbf{x}_t)}{\sum_{k=1}^K \mathbf{w}^t[k]}.$$

    Receive noisy label  $\tilde{y}_t$ ;

**for**  $k \in [K]$  **do**

        Set  $\mathbf{w}^{t+1}[k] = \mathbf{w}^t[k] e^{-\alpha \ell(\tilde{y}_t, f_k(\mathbf{x}_t))}$ ;

**end**

**end**

---

[Algorithm 2](#) provides the following regret bound ([Cesa-Bianchi and Lugosi, 2006](#), Proposition 3.1).

**Proposition 38** Let  $\mathcal{F} \subset \mathcal{D}(\tilde{\mathcal{Y}})^{\mathcal{X}}$  be any finite class of size  $K$  and  $\ell$  be an  $\alpha$ -Exp-concave loss. If  $\hat{p}_t$  is the predictor in [Algorithm 2](#), then for any  $\mathbf{x}^T \in \mathcal{X}^T$  and  $\tilde{y}^T \in \tilde{\mathcal{Y}}^T$  we have

$$\sup_{f \in \mathcal{F}} \sum_{t=1}^T \ell(\tilde{y}_t, \hat{p}_t) - \ell(\tilde{y}_t, f(\mathbf{x}_t)) \leq \frac{\log |\mathcal{F}|}{\alpha}.$$

**Proof** We now fix  $\mathbf{x}^T, \tilde{y}^T$  and any  $f^* \in \mathcal{F}$ . Denote  $W^t = \sum_{k=1}^K \mathbf{w}^t[k]$ . We have

$$\begin{aligned} \frac{W^{t+1}}{W^t} &= \sum_{k=1}^K \frac{\mathbf{w}^t[k] e^{-\alpha \ell(\tilde{y}_t, f_k(\mathbf{x}_t))}}{W^t} \\ &= \sum_{k=1}^K \frac{\mathbf{w}^t[k]}{W^t} e^{-\alpha \ell(\tilde{y}_t, f_k(\mathbf{x}_t))} \\ &\leq e^{-\alpha \ell(\tilde{y}_t, \sum_{k=1}^K \mathbf{w}^t[k] f_k(\mathbf{x}_t) / W^t)} \\ &= e^{-\alpha \ell(\tilde{y}_t, \hat{p}_t)}, \end{aligned}$$

where the inequality follows by Jensen's inequality and definition of  $\alpha$ -Exp-concavity, and the last equality follows by definition of  $\hat{p}_t$ . Therefore, by telescoping the product we have

$$\log W^{T+1} - \log W^1 = \log \frac{W^{T+1}}{W^1} = \log \prod_{t=1}^T \frac{W^{t+1}}{W^t} \leq -\alpha \sum_{t=1}^T \ell(\tilde{y}_t, \hat{p}_t).$$

Note that  $\log W^{T+1} = \log \left( \sum_{k=1}^K e^{-\alpha \sum_{t=1}^T \ell(\tilde{y}_t, f_k(\mathbf{x}_t))} \right) \geq -\alpha \sum_{t=1}^T \ell(\tilde{y}_t, f^*(\mathbf{x}_t))$  and  $\log W^1 = \log K$ , we have

$$\sum_{t=1}^T \ell(\tilde{y}_t, \hat{p}_t) - \ell(\tilde{y}_t, f^*(\mathbf{x}_t)) \leq \frac{\log K}{\alpha},$$

as needed. ■

## Appendix C. Omitted Proofs in Section 2.1

In this appendix, we present the omitted proofs from Section 2.1.

**Proof** [Proof of Proposition 7] By definition of Bergman divergence, we have

$$L(p, q_1) - L(p, q_2) = F(q_2) - F(q_1) - p^\top (\nabla F(q_1) - \nabla F(q_2)) + q_1^\top \nabla F(q_1) - q_2^\top \nabla F(q_2).$$

Note that the above expression is a *linear* function w.r.t.  $p$ . Therefore, by taking expectation over  $p \sim P$  and using the linearity of expectation, one can verify the claimed identity holds. ■

**Proof** [Proof of Proposition 8] The 1-Exp-concavity of log-loss can be verified directly. To prove the  $1/4$ -Exp-concavity of Brier loss, we have by Hazan et al. (2016, Lemma 4.2) that a function  $f$  is  $\alpha$ -Exp-concave if and only if

$$\alpha \nabla f(p) \nabla f(p)^\top \preceq \nabla^2 f(p).$$

For any  $q \in \mathcal{D}(\tilde{\mathcal{Y}})$ , we denote  $f(p) = \|p - q\|_2^2$ . We have  $\nabla f(p) = 2(p - q)$  and  $\nabla^2 f(p) = 2I$ , where  $I$  is the identity matrix. Taking any  $u \in \mathbb{R}^M$ , we have  $\frac{1}{4} \langle u, 2(p - q) \rangle^2 \leq \|u\|_2^2 \|p - q\|_2^2 \leq 2\|u\|_2^2 = 2u^\top I u$ , where the first inequality follows by Cauchy-Schwarz inequality and the second inequality follows by:

$$\|p - q\|_2^2 = \sum_{\tilde{y} \in \tilde{\mathcal{Y}}} (p[\tilde{y}] - q[\tilde{y}])^2 \leq \sum_{\tilde{y} \in \tilde{\mathcal{Y}}} \max\{p[\tilde{y}], q[\tilde{y}]\}^2 \leq \sum_{\tilde{y} \in \tilde{\mathcal{Y}}} p[\tilde{y}]^2 + q[\tilde{y}]^2 \leq 2,$$

since  $p, q \in \mathcal{D}(\tilde{\mathcal{Y}})$ . This completes the proof.  $\blacksquare$

**Proof** [Proof of Lemma 9] Let  $\Phi$  be the *Exponentially Weighted Average (EWA)* predictor as Algorithm 2 with input  $\mathcal{F}$  and loss  $\ell(\tilde{y}, p) \stackrel{\text{def}}{=} L(e_{\tilde{y}}, p)$ . Let  $\tilde{y}^T$  be any realization of the labels, we denote  $e_t$  to be the standard base of  $\mathbb{R}^M$  with value 1 at position  $\tilde{y}_t$  and zeros otherwise. By  $\alpha$ -Exp-concave and the regret bound of Proposition 38, we have

$$\sup_{f \in \mathcal{F}, \mathbf{x}^T \in \mathcal{X}^T, \tilde{y}^T \in \mathcal{Y}^T} \sum_{t=1}^T L(e_t, \hat{p}_t) - L(e_t, f(\mathbf{x}_t)) \leq \frac{\log |\mathcal{F}|}{\alpha}. \quad (22)$$

Note that this bound holds *point-wise* w.r.t. any individual sequences  $\mathbf{x}^T, \tilde{y}^T$ .

We now denote  $\mathbb{E}_t$  as the conditional expectation over the randomness of  $\tilde{y}^T$  conditioning on  $\tilde{y}^{t-1}$ . By Proposition 7, we have for all  $t \in [T]$  that

$$\mathbb{E}_t [L(e_t, \hat{p}_t) - L(e_t, f(\mathbf{x}_t))] = L(\tilde{p}_t, \hat{p}_t) - L(\tilde{p}_t, f(\mathbf{x}_t)),$$

since  $\mathbb{E}[e_t] = \tilde{p}_t$  if  $\tilde{y}_t \sim \tilde{p}_t$ ,  $\hat{p}_t$  depending only on  $\tilde{y}^{t-1}$  and  $L$  is Bregman divergence. We now take  $\mathbb{E}_{\tilde{y}^T}$  on both sides of (22). The result then follows by  $\sup \mathbb{E} \leq \mathbb{E} \sup$  and the law of total probability (i.e.,  $\mathbb{E}_{\tilde{y}^T} [X_1 + \dots + X_T] = \mathbb{E}_{\tilde{y}^T} [\mathbb{E}_1[X_1] + \dots + \mathbb{E}_T[X_T]]$  for any random variables  $X^T$ ). The last part follows by the fact that the exponential weighted average predictor automatically ensures  $\hat{p}_t$  is a convex combination of  $\{f(\mathbf{x}_t) : f \in \mathcal{F}\}$  for all  $t \in [T]$ .  $\blacksquare$

**Proof** [Proof of Lemma 10] Go through the same argument as in the proof of Lemma 9 and apply Lemma 35 to (22).  $\blacksquare$

## Appendix D. Proof of high probability minimax risk of Theorem 21

We begin with the following concentration result:

**Lemma 39** Let  $X_1, \dots, X_T$  be a random process adaptive to some filtration  $\{\mathcal{F}_t\}_{t \leq T}$ , and  $\mathbb{E}_t$  be the conditional expectation on  $\mathcal{F}_{t-1}$ . Then, for any  $\alpha, \delta > 0$  we have

$$\Pr \left[ -\sum_{t=1}^T \log \mathbb{E}_t [e^{-\alpha X_t}] \leq \alpha \sum_{t=1}^T X_t + \log(1/\delta) \right] \geq 1 - \delta.$$

**Proof** This follows directly by (Zhang, 2023, Theorem 13.2)  $\blacksquare$

The following key inequality holds:

**Lemma 40** Let  $\tilde{p} = (1 - \eta')e_{\tilde{y}} + \eta'u$ ,  $p = (1 - \eta)e_{\tilde{y}} + \eta u$  and  $\hat{p} = (1 - \eta)a + \eta u$ , where  $e_{\tilde{y}}, a, u \in \mathcal{D}(\tilde{\mathcal{Y}})$  and  $\eta' \leq \eta < 1$ , such that  $e_{\tilde{y}}$  is the distribution assigning probability 1 on  $\tilde{y}$ ,  $u$  is uniform over  $\tilde{\mathcal{Y}}$  and  $a \in \mathcal{D}(\tilde{\mathcal{Y}})$  is arbitrary. Then

$$\sum_{\tilde{y} \in \tilde{\mathcal{Y}}} \tilde{p}[\tilde{y}] \sqrt{\frac{\hat{p}[\tilde{y}]}{p[\tilde{y}]}} \leq \sum_{\tilde{y} \in \tilde{\mathcal{Y}}} p[\tilde{y}] \sqrt{\frac{\hat{p}[\tilde{y}]}{p[\tilde{y}]}} = \sum_{\tilde{y} \in \tilde{\mathcal{Y}}} \sqrt{p[\tilde{y}] \hat{p}[\tilde{y}]}. \quad (23)$$

**Proof** Denote  $|\tilde{\mathcal{Y}}| = M$ , and let  $r \in \mathbb{R}^{\tilde{\mathcal{Y}}}$  be the vector such that  $r[\tilde{y}] = \sqrt{\hat{p}[\tilde{y}]/p[\tilde{y}]}$ . We have the LHS of (23) equals  $e_{\tilde{y}}^\top r + \eta'(u - e_{\tilde{y}})^\top r$ . We claim that  $f(\eta') \stackrel{\text{def}}{=} e_{\tilde{y}}^\top r + \eta'(u - e_{\tilde{y}})^\top r$  attains maximum when  $\eta' = \eta$ , which will finish the proof. It is sufficient to prove that  $(u - e_{\tilde{y}})^\top r \geq 0$  since  $f(\eta')$  is a linear function w.r.t.  $\eta'$ . We have

$$u^\top r = \frac{1}{M} \sum_{\tilde{y}' \in \tilde{\mathcal{Y}}} \sqrt{\frac{\hat{p}[\tilde{y}']}{p[\tilde{y}']}}, \quad e_{\tilde{y}}^\top r = \sqrt{\frac{\hat{p}[\tilde{y}]}{p[\tilde{y}]}}.$$

We only need to show that  $\forall \tilde{y}' \in \tilde{\mathcal{Y}}$  with  $\tilde{y}' \neq \tilde{y}$ , we have  $\sqrt{\hat{p}[\tilde{y}']/p[\tilde{y}']} \geq \sqrt{\hat{p}[\tilde{y}]/p[\tilde{y}]}$ , i.e.,

$$\frac{p[\tilde{y}]}{p[\tilde{y}']} \geq \frac{\hat{p}[\tilde{y}]}{\hat{p}[\tilde{y}']}.$$

Note that,  $p[\tilde{y}] = 1 - \eta + \frac{\eta}{M}$ ,  $p[\tilde{y}'] = \frac{\eta}{M}$ ,  $\tilde{p}[\tilde{y}] = (1 - \eta)a[\tilde{y}] + \frac{\eta}{M}$  and  $\hat{p}[\tilde{y}'] = (1 - \eta)a[\tilde{y}'] + \frac{\eta}{M}$ , i.e., we have  $p[\tilde{y}] \geq \hat{p}[\tilde{y}], \hat{p}[\tilde{y}'] \geq p[\tilde{y}]$ . The result now follows by the simple fact that for any  $a \geq b, c \geq d \geq 0$  we have  $\frac{a}{d} \geq \frac{b}{c}$ .  $\blacksquare$

We are now ready to state our main result of this appendix, which establishes a high probability version of Theorem 21.

**Theorem 41** *Let  $\mathcal{H} \subset \mathcal{Y}^{\mathcal{X}}$  be any finite class,  $\mathcal{P}$  be the class of all singleton distributions over  $\mathcal{X}^T$ , and  $\mathcal{K}^\eta$  be as defined in Section 3.3 with  $0 \leq \eta < 1$ . Then, the high probability minimax risk at confidence  $\delta$  is upper bounded by*

$$B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}^\eta) \leq \frac{\log |\mathcal{H}| + 2 \log(1/\delta)}{(1 - \eta)^2/4}.$$

Furthermore, for  $1 - \eta \ll \frac{1}{M}$  we have  $B^\delta(\mathcal{H}, \mathcal{P}, \mathcal{K}^\eta) \leq O\left(\frac{\log |\mathcal{H}| + \log(1/\delta)}{M(1 - \eta)^2}\right)$ .

**Proof** Let  $\mathcal{F}$  be the class as in the proof of Theorem 21 and  $\hat{p}_t$  be the *Exponential Weighted Average* algorithm under Log-loss, see Algorithm 2. We have by Proposition 38 that for any  $\tilde{y}^T \in \tilde{\mathcal{Y}}^T$

$$\sup_{\mathbf{x}^T \in \mathcal{X}^T} \sum_{t=1}^T \log \frac{f^*(\mathbf{x}_t)[\tilde{y}_t]}{\hat{p}_t[\tilde{y}_t]} \leq \log |\mathcal{F}|$$

where  $f^*$  is the corresponding function of the underlying truth  $h^* \in \mathcal{H}$  (see the proof of Theorem 21). We now assume  $\tilde{y}^T$  are sampled from  $\tilde{p}^T$ , where  $\tilde{p}^T$  are the noisy distributions selected by the adversary. Denote by  $\mathbb{E}_t$  the conditional expectation on  $\tilde{y}^{t-1}$ . We have

$$\mathbb{E}_t \left[ e^{-\frac{1}{2} \log \frac{f^*(\mathbf{x}_t)[\tilde{y}_t]}{\hat{p}_t[\tilde{y}_t]}} \right] = \mathbb{E}_{\tilde{y}_t \sim \tilde{p}_t} \sqrt{\frac{\hat{p}[\tilde{y}_t]}{f^*(\mathbf{x}_t)[\tilde{y}_t]}} \leq \sum_{\tilde{y}_t \in \tilde{\mathcal{Y}}} \sqrt{\hat{p}[\tilde{y}_t] f^*(\mathbf{x}_t)[\tilde{y}_t]},$$

where the inequality follows from Lemma 40. By a similar argument as in the proof of (Foster et al., 2021, Lemma A.14), we have

$$\log \sum_{\tilde{y}_t \in \tilde{\mathcal{Y}}} \sqrt{\hat{p}[\tilde{y}_t] f^*(\mathbf{x}_t)[\tilde{y}_t]} = \log \left( 1 - \frac{1}{2} H^2(\hat{p}_t, f^*(\mathbf{x}_t)) \right) \leq -\frac{1}{2} H^2(\hat{p}_t, f^*(\mathbf{x}_t)),$$

where the first equality follows by definition of squared Hellinger divergence. Taking  $X_t = \log \frac{f^*(\mathbf{x}_t)[\tilde{y}_t]}{\hat{p}_t[\tilde{y}_t]}$ ,  $\alpha = \frac{1}{2}$  and invoking Lemma 39 we have w.p.  $\geq 1 - \delta$

$$\Pr \left[ \sum_{t=1}^T H^2(\hat{p}_t, f^*(\mathbf{x}_t)) \leq \log |\mathcal{F}| + 2 \log(1/\delta) \right] \geq 1 - \delta.$$

Let now  $\hat{y}_t = \arg \max_{\tilde{y}} \{\hat{p}_t[\tilde{y}] : \tilde{y} \in \tilde{\mathcal{Y}}\}$ . We have, if  $\hat{y}_t \neq h^*(\mathbf{x}_t)$

$$H^2(\hat{p}_t, f^*(\mathbf{x}_t)) \geq \|\hat{p}_t - f^*(\mathbf{x}_t)\|_1^2/4 \geq (1 - \eta)^2/4,$$

where the first inequality follows from  $\sqrt{H^2(p, q)} \geq \|p - q\|_1/2$  (Polyanskiy and Wu, 2022, Equation 7.20) and the second inequality follows the same as in the proof of Claim 1. Since  $H^2(p, q) \geq 0$  for all  $p, q$ , we have w.p.  $\geq 1 - \delta$  that

$$\sum_{t=1}^T 1\{\hat{y}_t \neq h^*(\mathbf{x}_t)\} \leq \frac{\log |\mathcal{H}| + 2 \log(1/\delta)}{(1 - \eta)^2/4}.$$

To prove the second part, we observe that if  $\hat{y}_t \neq h^*(\mathbf{x}_t)$ , then  $\hat{p}_t = (1 - \eta)a_t + \eta u$  such that  $a_t[h^*(\mathbf{x}_t)] \leq \frac{1}{2}$ . Since  $f^*(\mathbf{x}_t) = (1 - \eta)e_{h^*(\mathbf{x}_t)} + \eta u$ , we have by direct computation that

$$H^2(\hat{p}_t, f^*(\mathbf{x}_t)) \geq \left( \sqrt{(1 - \eta)/2} + \frac{\eta}{M} - \sqrt{1 - \eta + \frac{\eta}{M}} \right)^2 \sim \frac{M(1 - \eta)^2}{16},$$

where the last asymptote follows by Taylor expansion  $\frac{M(\eta-1)^2}{16} + O(\sum_{n=3}^{\infty} M^{n-1}(1 - \eta)^n)$  and the remainder term converges when  $1 - \eta \ll \frac{1}{M}$ .  $\blacksquare$

**Remark 42** Note that, Lemma 40 is the key that allows us to reduce our miss-specified setting to the well-specified case, such as (Foster et al., 2021, Lemma A.14), for which a reduction to the Hellinger divergence is possible.

**Unknown parameter  $\eta$ :** We now briefly discuss how to extend our results to the case when the parameter  $\eta$  is *unknown* to the learner. Note that, in this case we will not be able to construct the class  $\mathcal{F}$ . To resolve this issue, we can leverage the reduction to the pairwise comparison of two hypotheses as discussed in Section 3.2. More precisely, let  $h_1, h_2$  be two hypotheses. We can assume, w.l.o.g., that  $h_1(\mathbf{x}_t) = 0$  and  $h_2(\mathbf{x}_t) = 1$  for all time steps  $t$  (relabeling if necessary). Now, at each time step  $t + 1$ , we can simply compute the mean  $a_t = \frac{1}{t} \sum_{r=1}^t \tilde{y}_r$ . If  $a_t \leq \frac{1}{2} - \frac{\log(T/\delta)}{\sqrt{t}}$  or  $a_t \geq \frac{1}{2} + \frac{\log(T/\delta)}{\sqrt{t}}$  at some step  $t$ , we complete the process and declare  $h_1$  or  $h_2$  "wins", respectively. Now, if the parameter is  $\eta \in [0, 1)$ , we have, by Hoeffding and union bounds, w.p.  $\geq 1 - \delta$ , the process must be "completed" within  $O\left(\frac{\log^2(T/\delta)}{(1 - \eta)^2/4}\right)$  steps and identifies  $h_1$  or  $h_2$  with error probability  $\leq \delta$ . Invoking the reduction from Theorem 17 (replacing the update of  $S^t$  with a *stopping rule* that eliminates index  $k$  for which it declared "complete" with some  $k'$  and "losses"), we will obtain an  $O\left(\frac{\log |\mathcal{H}| \log^2(|\mathcal{H}|T/\delta)}{(1 - \eta)^2/4}\right)$  high probability minimax risk upper bound. This is of order  $\frac{\log^3 |\mathcal{H}|}{(1 - \eta)^2}$  and we leave it as open problem to obtain the optimal dependency.