

# Pairwise Alignment of Protein Interaction Networks

MEHMET KOYUTÜRK,<sup>1</sup> YOHAN KIM,<sup>2</sup> UMUT TOPKARA,<sup>1</sup>  
SHANKAR SUBRAMANIAM,<sup>2,3</sup> WOJCIECH SZPANKOWSKI,<sup>1</sup> and ANANTH GRAMA<sup>1</sup>

## ABSTRACT

With an ever-increasing amount of available data on protein–protein interaction (PPI) networks and research revealing that these networks evolve at a modular level, discovery of conserved patterns in these networks becomes an important problem. Although available data on protein–protein interactions is currently limited, recently developed algorithms have been shown to convey novel biological insights through employment of elegant mathematical models. The main challenge in aligning PPI networks is to define a graph theoretical measure of similarity between graph structures that captures underlying biological phenomena accurately. In this respect, modeling of conservation and divergence of interactions, as well as the interpretation of resulting alignments, are important design parameters. In this paper, we develop a framework for comprehensive alignment of PPI networks, which is inspired by duplication/divergence models that focus on understanding the evolution of protein interactions. We propose a mathematical model that extends the concepts of match, mismatch, and gap in sequence alignment to that of match, mismatch, and duplication in network alignment and evaluates similarity between graph structures through a scoring function that accounts for evolutionary events. By relying on evolutionary models, the proposed framework facilitates interpretation of resulting alignments in terms of not only conservation but also divergence of modularity in PPI networks. Furthermore, as in the case of sequence alignment, our model allows flexibility in adjusting parameters to quantify underlying evolutionary relationships. Based on the proposed model, we formulate PPI network alignment as an optimization problem and present fast algorithms to solve this problem. Detailed experimental results from an implementation of the proposed framework show that our algorithm is able to discover conserved interaction patterns very effectively, in terms of both accuracies and computational cost.

**Key words:** protein–protein interactions, network alignment, evolutionary models.

## 1. INTRODUCTION

**I**NCREASING AVAILABILITY OF EXPERIMENTAL DATA relating to biological sequences, coupled with efficient tools such as BLAST and CLUSTAL, have contributed to fundamental understanding of a variety of

---

<sup>1</sup>Department of Computer Sciences, Purdue University, West Lafayette, IN 47907.

<sup>2</sup>Department of Chemistry and Biochemistry, University of California at San Diego, La Jolla, CA 92093.

<sup>3</sup>Department of Bioengineering, University of California at San Diego, La Jolla, CA 92093.

biological processes (Altschul *et al.*, 1997; Thompson *et al.*, 1994). These tools are used for discovering common subsequences and motifs, which convey functional, structural, and evolutionary information. Recent developments in molecular biology have resulted in a new generation of experimental data that bear relationships and interactions between biomolecules (Hartwell *et al.*, 1999). An important class of molecular interaction data is in the form of protein–protein interaction (PPI) networks. These networks provide the experimental basis for understanding modular organization of cells, as well as useful information for predicting the biological function of individual proteins (Titz *et al.*, 2004). High-throughput screening methods such as two-hybrid analysis (Ito *et al.*, 2001), mass spectrometry (Ho *et al.*, 2002), and TAP (Gavin *et al.*, 2002) provide large amounts of data on the *interactome* of an increasing number of species.

Recent studies suggest that PPI networks evolve at a modular level (Wuchty *et al.*, 2003), and consequently, understanding conserved substructures through comparative analysis of these networks can provide basic insights into a variety of biochemical processes. As is the case with sequences, key problems on graphs derived from biomolecular interactions include aligning multiple graphs (Dandekar *et al.*, 1999; Koyutürk *et al.*, 2005; Sharan *et al.*, 2004, 2005; Tohsato *et al.*, 2000), finding frequently occurring subgraphs in a collection of graphs (Hu *et al.*, 2005; Koyutürk *et al.*, 2004; Sharan *et al.*, 2005), discovering highly conserved subgraphs in a pair of graphs, and finding good matches for a subgraph in a database of graphs (Kelley *et al.*, 2004). A class of comparative analysis techniques explore the problem of identifying conserved topological motifs in different species (Lotem *et al.*, 2004; Wuchty *et al.*, 2003). These studies reveal that many topological motifs are significantly conserved within and across species and proteins that are organized in cohesive patterns tend to be conserved to a higher degree.

A publicly available tool, PathBLAST, adopts the ideas in sequence alignment to PPI networks to discover conserved protein pathways across species (Kelley *et al.*, 2003, 2004). By restricting the alignment to pathways, i.e., linear chains of interacting proteins, this algorithm simplifies the problem, while preserving the biological implication of discovered patterns. PathBLAST accounts for gaps and mismatches by allowing nonrepeated jumps and matching of nonorthologous proteins, based on the notion that the orthologous counterpart of a pair of interacting proteins in one species will likely be indirectly interacting in the other (Kelley *et al.*, 2003). Pinter *et al.* (2005) align metabolic pathways based on subtree homeomorphism, observing that this model not only simplifies the problem by avoiding cycles, but also can describe variations in metabolic pathways effectively.

In addition to pathways and trees, a more general pattern structure is in the form of subgraphs induced by a group of proteins. Such subgraphs may provide insight into the conservation of functional modules and protein complexes, since these building blocks of cellular processes manifest themselves as dense or highly connected subgraphs in the PPI network (Bader and Hogue, 2003; Tornow and Mewes, 2003). Indeed, in a recent study, Sharan *et al.* (2004) show that cross-species network comparison in terms of general subgraphs provides novel biological insights through incorporation of knowledge about two different networks. Specifically, they identify conserved complexes in bacteria and yeast by constructing an orthology graph with nodes that correspond to pairs of orthologous proteins, one from each species. The edges of the orthology graph are weighted according to a probabilistic framework that compares null and conserved complex models based on log likelihood, which takes into account the conservation and density of interactions. The idea of constructing product graphs by joining orthologous nodes is also applied to the comparative analysis of PPI networks that belong to multiple species (Sharan *et al.*, 2005).

Based on the understanding of the structure of PPI networks that are available for several species, theoretical models that focus on understanding the evolution of protein interactions have been developed (Chung *et al.*, 2003; Eisenberg and Levanon, 2003; Pastor-Satorras *et al.*, 2003; Vázquez *et al.*, 2003; Wagner, 2003). Among these, the duplication/divergence model has been shown to be promising in explaining the power-law nature of PPI networks (Vázquez *et al.*, 2003). In this paper, we propose a framework for alignment of PPI networks based on these evolutionary models. As in Kelley *et al.* (2003, 2004) and Sharan *et al.* (2004, 2005), we construct product graphs by matching pairs of orthologous nodes. In contrast to Sharan *et al.* (2004), however, our framework is based on concepts of matches, mismatches, and duplications, and edges are weighted in order to reward or penalize these evolutionary events. This can be viewed as an extension of the concept of *alignment* in the sequence domain to that in network domain. Hence, our model provides a general framework that allows selection of parameters based on existing information about the conservation and divergence of proteins and their interactions, which can be refined in the light of a diverse range of mathematical models for network evolution (Berg *et al.*, 2004; Fraser


*et al.*, 2002; Kunin *et al.*, 2004; Pastor-Satorras *et al.*, 2003; Qin *et al.*, 2003; Wuchty, 2004). We reduce the resulting alignment problem to a graph-theoretic optimization problem and propose efficient heuristics to solve this problem. Experimental results based on an implementation of our framework show that the proposed algorithm is able to discover conserved interaction patterns very effectively. Another important difference between the proposed method and existing conserved network identification algorithms is that, by adapting the concepts of evolutionary models to comparative network analysis, our model allows interpretation of the divergence of functional modules through protein duplications and elimination/emergence of interactions. The proposed algorithm can also be adapted to finding matches for a subnet query in a database of PPI networks. The source code of the software and results on the pairwise alignment of PPI networks that belong to a variety of organisms are available at [www.cs.purdue.edu/homes/koyuturk/mawish/](http://www.cs.purdue.edu/homes/koyuturk/mawish/).

The rest of this paper is organized as follows: we start with a brief overview of duplication/divergence models for the evolution of PPI networks in Section 2. In Section 3, we define the alignment problem based on these models of evolution, formulate the problem as a graph optimization problem, and propose efficient heuristics for the solution of the problem. We illustrate the effectiveness of the proposed framework on comprehensive pairwise alignment of the PPI networks for three eukaryotic species in Section 4. We conclude our discussion in Section 5.

## 2. THEORETICAL MODELS FOR EVOLUTION OF PPI NETWORKS

There have been a number of studies aimed at understanding the general structure of PPI networks. These studies suggest that PPI networks can generally be modeled by power-law graphs, i.e., the relative frequency of proteins that interact with  $k$  proteins is roughly proportional to  $k^{-\gamma}$ , where  $\gamma$  is a network-specific parameter (Barabási and Albert, 1999). In order to explain this power-law nature, Barabási and Albert (1999) have proposed a network growth model based on preferential attachment, which is able to generate networks with degree distribution similar to PPI networks. According to this model, networks expand continuously by addition of new nodes, and these new nodes prefer to attach to well-connected nodes when joining the network. Observing that older proteins are better connected, Eisenberg and Levanon (2003) explain the evolutionary mechanisms behind such preference by the strength of selective pressure on maintaining connectivity of strongly connected proteins and creating proteins to interact with them. Furthermore, in a relevant study, it is observed that the interactions between groups of proteins that are temporally close in the course of evolution are likely to be conserved, suggesting synergistic selection during network evolution (Qin *et al.*, 2003).

A common model of evolution that explains preferential attachment is the duplication/divergence model, which is based on gene duplications (Pastor-Satorras *et al.*, 2003; Vázquez *et al.*, 2003; Wagner, 2001, 2003). According to this model, when a gene is duplicated in the genome, the node corresponding to the product of this gene is also duplicated together with its interactions. An example of protein duplication is shown in Fig. 1. A protein loses many aspects of its functions rapidly after being duplicated. This translates to divergence of duplicated (paralogous) proteins in the interactome through elimination and



**FIG. 1.** Duplication/divergence model for evolution of PPI networks. Starting with three interactions between three proteins, protein  $u_1$  is duplicated to add  $u'_1$  into the network together with its interactions (dashed circle and lines). Then,  $u_1$  loses its interaction with  $u_3$  (dotted line). Finally, an interaction between  $u_1$  and  $u'_1$  is added to the network (dashed line).

emergence of interactions. Elimination of an interaction in a PPI network implies the loss of an interaction between two proteins due to structural and/or functional changes. Similarly, emergence of an interaction in a PPI network implies the introduction of a new interaction between two noninteracting proteins caused by mutations that change protein surfaces. Examples of elimination and emergence of interactions are also illustrated in Fig. 1. If an elimination or emergence is related to a recently duplicated protein, it is said to be correlated; otherwise, it is uncorrelated (Pastor-Satorras *et al.*, 2003). Since newly duplicated proteins are more tolerant to interaction loss because of redundancy, correlated elimination is generally more probable than emergence and uncorrelated elimination (Vázquez *et al.*, 2003). It is also theoretically shown that network growth models based on node duplications generate power-law distributions (Chung *et al.*, 2003).

Since the elimination of interactions is related to sequence-level mutations, one can expect a positive correlation between similarity of interaction profiles and sequence similarity for paralogous proteins (Wagner, 2001). Indeed, the interaction profiles of duplicated proteins tend to almost totally diverge in about 200 million years, as estimated on the yeast interactome. On the other hand, the correlation between interaction profiles of duplicated proteins is significant for up to 150 million years after duplication, with more than half of interactions being conserved for proteins that are duplicated less than 50 million years back (Wagner, 2001). Consequently, when we consider the PPI networks that belong to two separate species, the in-paralogs will be likely to have more common interactions than out-paralogs. Here, we use the terms *in-paralog* and *out-paralog* for proteins that are duplicated before and after speciation, respectively (Remm *et al.*, 2001). While comparatively analyzing the proteome and interactome, it is important to distinguish in-paralogs from out-paralogs since the former are more likely to be functionally related. This, however, is a difficult task since out-paralogs also show sequence similarity.

In order to accurately identify and interpret conservation of interactions, complexes, and modules across species, we base our framework for the local alignment of PPI networks on duplication/divergence models. While searching for highly conserved groups of interactions, we evaluate mismatched interactions and paralogous proteins in light of the duplication/divergence model. Introducing the concepts of match (conservation), mismatch (emergence or elimination), and duplication, which are in accordance with widely accepted models of evolution, we are able to discover alignments that also allow speculation about the structure of the network in the common ancestor.

### 3. PAIRWISE LOCAL ALIGNMENT OF PPI NETWORKS

In light of the theoretical models of evolution of PPI networks, we develop a framework for the comparison of PPI networks in two different species. We formally define a computational problem that captures the underlying biological phenomena using matches, mismatches, and duplications. We then formulate PPI network alignment as a graph optimization problem and propose efficient heuristics to effectively solve this problem.

#### 3.1. The PPI network alignment problem

A PPI network is conveniently modeled by an undirected graph  $G(U, E)$ , where  $U$  denotes the set of proteins and  $uu' \in E$  denotes an interaction between proteins  $u \in U$  and  $u' \in U$ . For pairwise alignment of PPI networks, we are given two PPI networks belonging to two different species, denoted by  $G(U, E)$  and  $H(V, F)$ . The homology between a pair of proteins is quantified by a similarity measure that is defined as a function  $S : (U \cup V) \times (U \cup V) \rightarrow \mathfrak{R}$ . For any  $u, v \in U \cup V$ ,  $S(u, v)$  measures the degree of confidence in  $u$  and  $v$  being orthologous, where  $0 \leq S(u, v) \leq 1$ . If  $u$  and  $v$  belong to the same species, then  $S(u, v)$  quantifies the likelihood that the two proteins are in-paralogs. Quantity  $S$  is expected to be sparse; i.e., each protein is expected to have only a few potential orthologs. We discuss the methodology for deriving similarity scores from sequence alignments in Section 3.1.3.

For PPI networks  $G(U, E)$  and  $H(V, F)$ , a *protein subset pair*  $P = \{\tilde{U}, \tilde{V}\}$  is defined as a pair of protein subsets  $\tilde{U} \subseteq U$  and  $\tilde{V} \subseteq V$ . Any protein subset pair  $P$  induces a local alignment  $\mathcal{A}(G, H, S, P) = \{\mathcal{M}, \mathcal{N}, \mathcal{D}\}$  of  $G$  and  $H$  with respect to  $S$ , characterized by a set of duplications  $\mathcal{D}$ , a set of matches  $\mathcal{M}$ , and a set of mismatches  $\mathcal{N}$ . The biological analog of a *duplication* is the duplication of a gene in the

course of evolution. Each duplication is associated with a score that reflects the divergence of function between the two proteins, estimated using their similarity. A *match* corresponds to a conserved interaction between two orthologous protein pairs, which is rewarded by a match score that reflects our confidence in both protein pairs being orthologous. A *mismatch*, on the other hand, is the lack of an interaction in the PPI network of one organism between a pair of proteins whose orthologs interact in the other organism. A mismatch may correspond to the emergence of a new interaction or the elimination of a previously existing interaction in one of the species after the split, or an experimental error. Thus, mismatches are penalized to account for the divergence from the common ancestor. We provide formal definitions for these three concepts to construct a basis for the formulation of local alignment as an optimization problem. Note that although PPI networks are undirected graphs, interactions are regarded as ordered pairs in the following definitions for convenience; i.e., for an interaction  $uu' \in E$ , there is also an interaction  $u'u \in E$ , which is essentially the same interaction.

**Definition 1. Local Alignment of PPI Networks.** *Given protein interaction networks  $G(U, E)$ ,  $H(V, F)$ , let functions  $\Delta_G(u, u')$  and  $\Delta_H(v, v')$  denote the distance between two corresponding proteins in the interaction graphs  $G$  and  $H$ , respectively. Given a pairwise similarity function  $S$  defined over the union of their protein sets  $U \cup V$ , and a distance cutoff  $\bar{\Delta}$ , any protein subset pair  $P = (\tilde{U}, \tilde{V})$  induces a local alignment  $\mathcal{A}(G, V, S, P) = \{\mathcal{M}, \mathcal{N}, \mathcal{D}\}$ , where*

$$\mathcal{M} = \{u, u' \in \tilde{U}, v, v' \in \tilde{V} : S(u, v) > 0, S(u', v') > 0, \\ ((uu' \in E \wedge \Delta_H(v, v') \leq \bar{\Delta}) \vee (vv' \in F \wedge \Delta_G(u, u') \leq \bar{\Delta}))\}, \quad (1)$$

$$\mathcal{N} = \{u, u' \in \tilde{U}, v, v' \in \tilde{V} : S(u, v) > 0, S(u', v') > 0, \\ ((uu' \in E \wedge \Delta_H(v, v') > \bar{\Delta}) \vee (vv' \in F \wedge \Delta_G(u, u') > \bar{\Delta}))\}, \quad (2)$$

$$\mathcal{D} = \{u, u' \in \tilde{U} : S(u, u') > 0\} \cup \{v, v' \in \tilde{V} : S(v, v') > 0\}. \quad (3)$$

Each match  $M \in \mathcal{M}$ , mismatch  $N \in \mathcal{N}$ , and duplication  $D \in \mathcal{D}$  are associated with scores  $\mu(M)$ ,  $\nu(N)$ , and  $\delta(D)$ , respectively.

Following the definition of match and mismatch, while assessing the conservation of interactions, we take into account not only direct but also indirect interactions. If two proteins directly interact with each other in one organism and their orthologs are reachable from each other via at most  $\bar{\Delta}$  interactions in the other, we consider this a match. Conversely, a mismatch corresponds to the situation in which two proteins cannot reach each other via  $\bar{\Delta}$  interactions in one network while their orthologs directly interact in the other. This approach is motivated by two observations. First, proteins that are linked by a short alternate path are more likely to tolerate losing their interaction because of relaxation of evolutionary pressure. Second, high-throughput methods such as TAP (Gavin *et al.*, 2002) identify complexes that are associated with a single central protein, and these complexes are recorded in the interaction database as star networks with the central protein serving as a hub. Therefore, all proteins that are part of a particular complex can be viewed as interacting by setting  $\bar{\Delta} = 2$ .

*3.1.1. Scoring match, mismatch, and duplications.* For scoring matches and mismatches, we define the similarity between two protein pairs as follows:

$$S(uu', vv') = S(u, v)S(u', v') \quad (4)$$

where  $S(uu', vv')$  quantifies the likelihood that the interactions between  $u$  and  $v$ , and  $u'$  and  $v'$  are orthologous. Consequently, a match that corresponds to a conserved pair of orthologous interactions is rewarded as follows:

$$\mu(uu', vv') = \bar{\mu}S(uu', vv'). \quad (5)$$

Here,  $\bar{\mu}$  is the match coefficient that is used to tune the relative weight of matches against mismatches and duplications, based on the evolutionary distance between the species that are being compared.

A mismatch may correspond to the functional divergence of either interacting partner after speciation. It might also be due to a false positive or negative in one of the networks that is caused by incompleteness of data or experimental error (Titz *et al.*, 2004). However, considering indirect interactions as matches compensates for the second case to a certain extent. According to Wagner (2001), after a duplication event, duplicate proteins that retain similar functions in terms of being part of similar processes are likely to be part of the *same subnet*. Moreover, since conservation of proteins in a particular *module* is correlated with interconnectedness (Wuchty *et al.*, 2003), we expect that interacting partners that are part of a common functional module will at least be linked by short alternative paths. Based on these observations, we penalize mismatches for possible divergence in function as follows:

$$v(uu', vv') = -\bar{v}S(uu', vv'). \quad (6)$$

As for match score, mismatch penalty is also normalized by a coefficient  $\bar{v}$  that determines the relative weight of mismatches w.r.t. matches and duplications.

While aligning PPI networks, the motivation is to identify conserved patterns of interactions between orthologous proteins. For assessing the likelihood of orthology between proteins, the similarity score defined above relies on sequence homology. However, out-paralogs, which are proteins that are duplicated before the species split and hence cannot be considered orthologs, often show sequence similarities as well (Remm *et al.*, 2001). Since duplicated proteins rapidly lose their interactions, it is more likely that in-paralogs, i.e., the proteins that are duplicated after a split, will share more interacting partners than out-paralogs do (Wagner, 2001). Therefore, penalizing mismatches implicitly favors *real* orthologs by penalizing the out-paralogs for each interaction that is lost after duplication. Furthermore, we employ sequence similarity as a means for distinguishing in-paralogs from out-paralogs. This is based on the observation that sequence similarity provides a crude approximation for the age of duplication (Wagner, 2003). With the expectation that recently duplicated proteins, which are more likely to be in-paralogs, show more significant sequence similarity than older paralogs, we define duplication score as follows:

$$\delta(u, u') = \bar{\delta}(S(u, u') - \bar{d}). \quad (7)$$

Here  $\bar{d}$  is the cut-off for being considered in-paralogs. If  $S(u, u') > \bar{d}$ , suggesting that  $u$  and  $u'$  are likely to be in-paralogs, the duplication is rewarded by a positive score. If  $S(u, u') < \bar{d}$ , on the other hand, the proteins are considered out-paralogs; therefore, the duplication is penalized.

*3.1.2. Alignment score and the optimization problem.* The above formulation of match, mismatch, and duplication translates the problem of distinguishing orthologs and in-paralogs from out-paralogs to an optimization problem that accounts for the tradeoff between conservation of sequences and interactions. This enables accurate identification of conserved interactions between ortholog protein pairs, while allowing us to define the pairwise local alignment for interspecies comparison of PPI networks as an optimization problem.


**Definition 2. Alignment Score and PPI Network Alignment Problem.** Given PPI networks  $G$  and  $H$ , the score of alignment  $\mathcal{A}(G, H, S, P) = \{\mathcal{M}, \mathcal{N}, \mathcal{D}\}$  is defined as

$$\sigma(\mathcal{A}) = \sum_{M \in \mathcal{M}} \mu(M) + \sum_{N \in \mathcal{N}} v(N) + \sum_{D \in \mathcal{D}} \delta(D). \quad (8)$$

The PPI network alignment problem is one of finding all maximal protein subset pairs  $P$  such that  $\sigma(\mathcal{A}(G, H, S, P))$  is locally maximal, i.e., the alignment score cannot be improved by adding individual proteins to or removing proteins from  $P$ .

We aim to find local alignments with locally maximal score (drawing an analogy to sequence alignment (Smith and Waterman, 1981), *high-scoring subgraph pairs*).

We illustrate the concepts of match, mismatch, and duplication using a simple example. Consider the two interaction networks  $G$  and  $H$  shown in Fig. 2a. The alignment induced by the protein subset pair  $\tilde{U} = \{u_1, u_2, u_3, u_4\}$  and  $\tilde{V} = \{v_1, v_2, v_3\}$  is shown in Fig. 2b, where we set  $\bar{\Delta} = 1$ . The only duplication



**FIG. 2.** (a) An instance of the pairwise local alignment problem. The proteins that have nonzero similarity scores (i.e., are potentially orthologous), are shaded the same. Note that  $S$  does not necessarily induce a disjoint grouping of proteins in practice. (b) A local alignment induced by the protein subset pair  $\{u_1, u_2, u_3, u_4\}$  and  $\{v_1, v_2, v_3\}$ . Ortholog and paralog proteins are vertically aligned. Existing interactions are shown by solid lines; missing interactions that have an existing ortholog counterpart are shown by dotted lines. Solid interactions between two aligned proteins in separate species correspond to a match; one solid, one dotted interaction between two aligned proteins in separate species correspond to a mismatch. Proteins in the same species that are on the same vertical line correspond to duplications.

in this alignment is  $(u_1, u_2)$ . If this alignment is chosen to be a “good” one, then, based on the existence of this duplication in the alignment, if  $S(u_2, v_1) < S(u_1, v_1)$ , we can speculate that  $u_1$  and  $v_1$  have evolved from the same gene in the common ancestor, while  $u_2$  is an in-paralog that emerged from duplication of  $u_1$  after split. The match set consists of interaction pairs  $(u_1u_1, v_1v_1)$ ,  $(u_1u_2, v_1v_1)$ ,  $(u_1u_3, v_1v_3)$ , and  $(u_2u_4, v_1v_2)$ . Observe that  $v_1$  is mapped to both  $u_1$  and  $u_2$  in the context of different interactions. This is associated with the functional divergence of  $u_1$  and  $u_2$  after duplication. Furthermore, the self-interaction of  $v_1$  in  $H$  is mapped to an interaction between paralogous proteins in  $G$ .

The mismatch set is composed of  $(u_1u_4, v_1v_2)$ ,  $(u_2u_2, v_1v_1)$ ,  $(u_2u_3, v_1v_3)$ , and  $(u_3u_4, v_3v_2)$ . The interaction  $u_3u_4$  in  $G$  is left unmatched by this alignment, since the only possible pair of proteins in  $\tilde{V}$  that are orthologous to these two proteins are  $v_3$  and  $v_2$ , which do not interact in  $H$ . One conclusion that can be derived from this alignment is the elimination or emergence of this interaction in one of the species after the split. The indirect path between  $v_3$  and  $v_2$  through  $v_1$  may also serve as a basis for the tolerance to the loss of this interaction. Indeed, if we set  $\tilde{\Delta} = 2$ , then this pair of a direct and an indirect interaction would be considered a match. However, if we include  $v_4$  in  $\tilde{V}$  as well, then the induced alignment is able to match  $u_3u_4$  and  $v_3v_4$ . This strengthens the likelihood that this interaction existed in the common ancestor. However,  $v_4$  comes with another duplication since it is paralogous to  $v_2$ . Hence, if  $S(v_2, v_4) > \tilde{d}$ , the alignment that includes  $v_4$  will be favored over the present one. However, if  $S(v_2, v_4) < \tilde{d}$ , then  $v_4$  must compensate for the duplication penalty with the strength of its matching interactions in order to be included in the alignment.

**3.1.3. Estimation of similarity scores.** The similarity score  $S(u, v)$  quantifies the likelihood that proteins  $u$  and  $v$  are orthologous. We approximate this likelihood using the BLAST (Altschul *et al.*, 1997)  $E$ -value taking existing ortholog databases as point of reference. Let  $\mathbf{O}$  be the set of all orthologous protein pairs derived from COG (Tatusov *et al.*, 2000), or any other ortholog database. For proteins  $u$  and  $v$  with BLAST  $E$ -value  $E(u, v) < \tilde{E}$ , we approximate the probability of  $u$  and  $v$  being orthologous by

$$S(u, v) = P(E(u, v) < \tilde{E} | O_{uv}) = \frac{|\{u'v' \in \mathbf{O} : E(u', v') < \tilde{E}\}|}{|\mathbf{O}|} \quad (9)$$

where  $O_{uv}$  represents the event that  $u$  and  $v$  are orthologous. If we assume that the probability of a protein pair being orthologous ( $P(O_{uv})$ ) is a monotonically decreasing function of the  $E$ -value, then this quantity provides a measure of the likelihood that two proteins with  $E$ -value  $\tilde{E}$  are orthologous.

## 3.2. Alignment graph and the maximum-weight induced subgraph problem

It is possible to represent information regarding matches and mismatches between two PPI networks using a single alignment graph. This graph is a modified version of the graph Cartesian product that takes orthology into account. Assigning appropriate weights to the edges of the alignment graph, the local alignment problem defined in the previous section can be reduced to an optimization problem on this alignment graph. We define the following alignment graph:

**Definition 3. Alignment Graph.** For a pair of PPI networks  $G(U, E)$ ,  $H(V, F)$ , and protein similarity function  $S$ , the corresponding weighted alignment graph  $\mathbf{G}(\mathbf{V}, \mathbf{E})$  is computed as follows:

$$\mathbf{V} = \{\mathbf{v} = \{u, v\} : u \in U, v \in V \text{ and } S(u, v) > 0\}. \quad (10)$$


In other words, we have a node in the alignment graph for each pair of ortholog proteins. Each edge  $\mathbf{v}\mathbf{v}' \in \mathbf{E}$ , where  $\mathbf{v} = \{u, v\}$  and  $\mathbf{v}' = \{u', v'\}$ , is assigned weight

$$w(\mathbf{v}\mathbf{v}') = \mu(uu', vv') + \nu(uu', vv') + \delta(u, u') + \delta(v, v'). \quad (11)$$

Here,  $\mu(uu', vv') = 0$  if  $(uu', vv') \notin \mathcal{M}$ , and similarly for mismatches and duplications.

Note that the *alignment graph* is conceptually equivalent to the *global alignment graph* of Kelley *et al.* (2003) and the *orthology graph* of Sharan *et al.* (2004), with slight differences in formulation. In all models, the nodes of the alignment/orthology graph is constructed from any pair of potentially orthologous proteins in the two networks. On the other hand, in the above-defined alignment graph, all evolutionary information is encoded into edge weights through the concepts of matches, mismatches, and duplications.

Consider the PPI networks in Fig. 2a. To construct the corresponding alignment graph, we first compute the product of these two PPI networks to obtain five nodes that correspond to five ortholog protein pairs. We then insert an edge between two nodes of this graph if the corresponding proteins interact in both networks (*match edge*), interact in only one of the networks (*mismatch edge*), or at least one of them is paralogous (*duplication edge*), resulting in the alignment graph of Fig. 3a. Note that the weights assigned to these edges, which are shown in the figure, are not constant, but are functions of their incident nodes. Observe that the edge between  $\{u_1, v_1\}$  and  $\{u_2, v_1\}$  acts a match and duplication edge at the same time, allowing analysis of the conservation of self-interactions of duplicated proteins. This construction of the alignment graph allows us to formulate the alignment problem as a graph optimization problem defined below.



**FIG. 3.** (a) Alignment graph corresponding to the instance of Fig. 2a. Note that match scores, mismatch penalties and duplication scores are functions of incident nodes, which is not explicitly shown in the figure for simplicity. (b) Subgraph induced by node set  $\tilde{\mathbf{V}} = \{\{u_1, v_1\}, \{u_2, v_1\}, \{u_3, v_3\}, \{u_4, v_2\}\}$ , which corresponds to the alignment shown in Fig. 2b.

**Definition 4. Maximum Weight Induced Subgraph Problem (MAWISH).** Given graph  $G(\mathbf{V}, \mathbf{E})$  and a constant  $\epsilon$ , find a subset of nodes,  $\tilde{\mathbf{V}} \in \mathbf{V}$  such that the sum of the weights of the edges in the subgraph induced by  $\tilde{\mathbf{V}}$  is at least  $\epsilon$ , i.e.,  $W(\tilde{\mathbf{V}}) = \sum_{\mathbf{v}, \mathbf{v}' \in \tilde{\mathbf{V}}} w(\mathbf{v}\mathbf{v}') \geq \epsilon$ .

Not surprisingly, this problem is equivalent to the decision version of the local alignment problem defined in the previous section, as formally stated in the following theorem:

**Theorem 1.** Given PPI networks  $G, H$ , and a protein similarity function  $S$ , let  $G(\mathbf{V}, \mathbf{E}, w)$  be the corresponding alignment graph. If  $\tilde{\mathbf{V}}$  is a solution to the maximum weight induced subgraph problem on  $G(\mathbf{V}, \mathbf{E}, w)$ , then  $P = \{\tilde{U}, \tilde{V}\}$  induces an alignment  $\mathcal{A}(G, H, S, P)$  with  $\sigma(\mathcal{A}) = W(\tilde{\mathbf{V}})$ , where  $\tilde{U} = \{u \in U : \exists v \in V \text{ s.t. } \{u, v\} \in \tilde{\mathbf{V}}\}$  and  $\tilde{V} = \{v \in V : \exists u \in U \text{ s.t. } \{u, v\} \in \tilde{\mathbf{V}}\}$ .

**Proof.** Follows directly from the construction of alignment graph. ■

The induced subgraph that corresponds to the local alignment in Fig. 2b is shown in Fig. 3b.

It can be shown that MAWISH is NP-complete by reduction from maximum-clique, by assigning unit weight to edges and  $-\infty$  to nonedges. This problem is closely related to the maximum edge subgraph (Feige *et al.*, 2001) and maximum dispersion problems (Hassin *et al.*, 1997), which are also NP-complete. However, the positive weight restriction on these problems limits the application of existing algorithms to the maximum weight induced subgraph problem. Nevertheless, the local PPI network alignment problem aims to find all locally maximal alignments; consequently, locally optimal solutions of MAWISH are sufficient. Observing the similarity between min-cut graph partitioning and MAWISH, we develop fast heuristics based on common graph partitioning algorithms to identify locally maximal heavy subgraphs in the alignment graph.

### 3.3. Algorithms for local alignment of PPI networks

In terms of protein–protein interactions, functional modules are likely to be densely connected while being separable from other modules, i.e., a protein in a particular module interacts with most proteins in the same module either directly or through a common module hub, while it is only loosely connected to the rest of the network (Tornow and Mewes, 2003). Since analysis of conserved motifs reveals that proteins in highly connected motifs are more likely to be conserved, suggesting that such dense motifs are parts of functional modules (Wuchty *et al.*, 2003), high-scoring local alignments are likely to correspond to functional modules. Therefore, in the alignment graph, we can expect that proteins that belong to a conserved module will induce heavy subgraphs, while being loosely connected to other parts of the graph. This observation motivates the process of greedily growing a subgraph seeded at heavy nodes. This approach is shown to perform well in discovering conserved (Sharan *et al.*, 2004) or dense (Bader, 2003) subnets in PPI networks.

For min-cut graph partitioning, the most commonly applied heuristics are based on starting with a seed partition and repeatedly moving or swapping nodes with maximum gain on the objective function (Kernighan and Lin, 1970). The key point here is that the move is performed even if it is associated with a negative gain in order to climb over poor local optima. Observe that minimizing the total weight of the cut edges (min-cut) in graph partitioning is equivalent maximizing the total weight of internal edges. This is very similar to the objective function of MAWISH. The difference is that the total weight of only one part is considered in MAWISH and node balance is not an issue. Therefore, we apply this iterative improvement based heuristic to MAWISH in order to find locally maximal heavy subgraphs. The initial heavy subgraph is constructed by selecting the node with maximum number of matched interactions (i.e., a *conserved hub*) and adding all nodes that share a match edge with this node to the subgraph.

A sketch of this iterative improvement based algorithm for finding a single conserved subgraph on the alignment graph is shown in Fig. 4. Each pass (i.e., the loop between lines 3–13) of this algorithm works in linear time. In practice, we also limit the number of contiguous moves with negative gain. This allows us to tune the locality of identified patterns.

To find all nonredundant heavy subgraphs, we start with the entire alignment graph and find a maximally heavy subgraph. If this subgraph is statistically significant, we record the alignment that corresponds to this subgraph and mark its nodes. We repeat this process by considering only unmarked nodes. Once a

```

procedure HEAVIESTSUBGRAPH(G)
  ▷ Input  $\mathbf{G}(\mathbf{V}, \mathbf{E}, w)$ : Alignment graph
  ▷ Output  $\tilde{\mathbf{V}}$ : Subset of nodes that induces a maximally heavy subgraph in  $\mathbf{G}$ 
1   $\tilde{v} \leftarrow \operatorname{argmax}_{v \in \mathbf{V}} |\{v' \in \mathbf{V} : (v, v') \text{ is a match edge}\}|$ 
2   $\tilde{\mathbf{V}} \leftarrow \{\tilde{v}\} \cup \{v \in \mathbf{V} : (\tilde{v}, v) \text{ is a match edge}\}$ 
3  repeat
4     $Q \leftarrow \{v \in \mathbf{V} : \operatorname{key}(v) = -\sum_{v' \in \tilde{\mathbf{V}}} w(v, v') \text{ if } v \in \tilde{\mathbf{V}}, \operatorname{key}(v) = \sum_{v' \in \tilde{\mathbf{V}}} w(v, v') \text{ else}\}$ 
5     $W_{max} \leftarrow W(\tilde{\mathbf{V}})$ 
6    while  $Q \neq \emptyset$ 
7       $v \leftarrow \operatorname{EXTRACTMAX}(Q)$ 
8      if  $v \in \tilde{\mathbf{V}}$  then  $\tilde{\mathbf{V}} \leftarrow \tilde{\mathbf{V}} \setminus \{v\}$  else  $\tilde{\mathbf{V}} \leftarrow \tilde{\mathbf{V}} \cup \{v\}$ 
9      if  $W(\tilde{\mathbf{V}}) > W_{max}$  then  $W_{max} \leftarrow W(\tilde{\mathbf{V}})$ ,  $bestmove \leftarrow v$ 
10     for all  $v'$  such that  $vv' \in \mathbf{E}$  update  $\operatorname{key}(v')$ 
11   endwhile
12   roll back all moves after bestmove
13 until  $bestmove = \text{NULL}$ 
14 return  $\tilde{\mathbf{V}}$ 

```

**FIG. 4.** Fast heuristic for finding a subset of nodes that induces a subgraph of maximal total weight on the alignment graph.

new significant subgraph is identified, we add the previously marked nodes that are positively connected to this subgraph one by one, unless the resulting subgraph becomes redundant. A subgraph is said to be redundant if there exists a subgraph which contains  $r\%$  of its nodes, where  $r$  is a user-defined threshold that determines the extent of allowed overlap. This method allows identification of overlapping alignments while avoiding redundancy. Finally, we rank all subgraphs based on their significance and report the corresponding alignments.

### 3.4. Statistical significance

To evaluate the statistical significance of discovered high-scoring alignments, we compare them with a reference model generated by a random source. In the reference model, it is assumed that the interaction networks of the two organisms are independent of each other. In order to assess the significance of conservation of interactions between orthologous proteins rather than the conservation of proteins itself, we assume that the orthology relationship between protein is already established, i.e., is not generated by a random source. To accurately capture the power-law nature of PPI networks, we assume that the

interactions are generated randomly from a distribution characterized by a given degree sequence. (Note that the power law nature of the graphs is not critical to our algorithm. The degree distribution can be computed explicitly from the database of interactions). If proteins  $u$  and  $u'$  are interacting with  $d_u$  and  $d_{u'}$  proteins, respectively, then the probability  $p_{uu'}$  of observing an interaction between  $u$  and  $u'$  can be estimated as  $p_{uu'} = d_u d_{u'} / \sum_{v \in U} d_v$  (Chung *et al.*, 2003).

Recall that the weight of a subgraph of the alignment graph is equal to the score of the corresponding alignment. Hence, in the reference model, the expected value of the score of an alignment induced by  $\tilde{\mathbf{V}} \subseteq \mathbf{V}$  is  $\mathbf{E}[W(\tilde{\mathbf{V}})] = \sum_{\mathbf{v}, \mathbf{v}' \in \tilde{\mathbf{V}}} \mathbf{E}[w(\mathbf{v}\mathbf{v}')]$ , where

$$\begin{aligned} \mathbf{E}[w(\mathbf{v}\mathbf{v}')] &= \bar{\mu}S(uu', vv')p_{uu'}p_{vv'} - \bar{v}S(uu', vv')(p_{uu'}(1 - p_{vv'})) \\ &\quad + (1 - p_{uu'})p_{vv'} + \delta(u, u') + \delta(v, v') \end{aligned} \quad (12)$$

is the expected weight of an edge in the alignment graph. With the simplifying assumption of independence of interactions, we have  $\mathbf{Var}[W(\tilde{\mathbf{V}})] = \sum_{\mathbf{v}, \mathbf{v}' \in \tilde{\mathbf{V}}} \mathbf{Var}[w(\mathbf{v}\mathbf{v}')]$ , enabling us to compute the  $z$ -score to evaluate the statistical significance of each discovered high-scoring alignment, under the normal approximation that we assume.

While the approach described above enables quick calculation of significance without repeated simulations or extensive numerical computations, it has a few shortcomings. First, the significance of an identified pattern is estimated for the proteins involved in that conserved subgraph, rather than computing the probability of the existence of the pattern anywhere in the networks. Second, the model does not take into account the variability in the distribution of orthologs. These cause low variability of alignment score in the reference model, leading to overestimated  $z$ -scores, since the observed variances in alignment score are fairly high, which indeed is statistically significant. As models and techniques for analytical assessment of statistical significance of network structures become available, the significance of conserved subnets can also be evaluated more reliably (Koyutürk *et al.*, 2006).

### 3.5. Extensions to the model

The proposed model can be extended to account for data quality as well as algorithm parameters.

*3.5.1. Accounting for experimental error.* PPI networks obtained from high-throughput screening are prone to errors in terms of both false negatives and positives (Titz *et al.*, 2004). While the proposed framework can be used to detect experimental errors through cross-species comparison to a certain extent, experimental noise can also degrade the performance of the alignment algorithm. In other words, mismatches should be penalized for lost interactions during evolution, not for experimental false negatives. To account for such errors while analyzing interaction networks, several methods have been developed to quantify the likelihood of an interaction or complex co-membership between proteins (Ashtana *et al.*, 2004; Gilchrist *et al.*, 2003; Jansen *et al.*, 2003). Given the prior probability distribution for protein interactions and a set of observed interactions, these methods compute the posterior probability of interactions based on Bayesian models. Hence, PPI networks can be modeled by weighted graphs to account for experimental error more accurately.

While the network alignment framework introduced in Section 3.1 assumes that interactions are represented by unweighted edges, it can be easily generalized to a weighted graph model as follows. Assuming that weight  $\varpi_{uv}$  represents the posterior probability of interaction between  $u$  and  $v$ , we can define match score and mismatch penalty in terms of their expected values derived from these posterior probabilities. Therefore, for any  $u, u' \in U$  and  $v, v' \in V$ , we have

$$\mu(uu', vv') = \bar{\mu}S(uu', vv')\varpi_{uu'}\varpi_{vv'} \quad (13)$$

$$v(uu', vv') = \bar{v}S(uu', vv')(\varpi_{uu'}(1 - \varpi_{vv'}) + (1 - \varpi_{uu'})\varpi_{vv'}). \quad (14)$$

Note that match and mismatch sets are not necessarily disjoint here in contrast to the unweighted graph model, which is a special case of this model.

3.5.2. *Alternative model components and parameters.* **Contracting Paralogs.** An alternate approach for handling duplications is contracting the proteins in the same species that are likely to be in-paralogs. This approach fits into the alignment graph model since in-paralogs are expected to be consistently orthologous to the same set of proteins in the other organism. It also reduces the computational complexity since the number of nodes will be decreased by node contraction and the edges that correspond to duplications will be eliminated. Contraction of nodes is also shown to be effective for multiple alignment of metabolic pathways using graph mining (Koyutürk *et al.*, 2004). However, clustering proteins in the same organism to identify in-paralogs requires preprocessing to solve a difficult problem. Clustering algorithms that are specifically designed for this purpose, such as INPARANOID (Remm *et al.*, 2001) serve as reliable tools. However, the resulting graphs may produce conservative alignments since the search space is narrowed down by the clustering of proteins (Koyutürk *et al.*, 2005). In contrast, accounting for duplications using duplication edges provides more flexibility and uses conservation of interactions as additional information to distinguish in-paralogs from out-paralogs, as discussed above.

**Shortest-path mismatch model.** In the above discussion, while we consider proteins that are linked by at most  $\bar{\Delta}$  interactions as interacting, we do not take into account the distance while penalizing mismatches. We can extend this to a shortest-path mismatch model, defined as follows:

$$v(uu', vv') = \bar{v}S(uu', vv')(\max\{\Delta_G(u, u'), \Delta_H(v, v')\} - \bar{\Delta}). \quad (15)$$

While this model may improve the alignment algorithm, it is computationally expensive since it requires solution of the all pairs shortest path problem on both PPI networks.

**Linear duplication model.** The alignment graph model forces each duplicate pair in an alignment to be scored. For example, if an alignment contains  $n$  paralogous proteins in one species,  $\binom{n}{2}$  duplications are scored to account for each duplicate pair. However, in the evolutionary process, each paralogous protein is the result of a single duplication, i.e.,  $n$  paralogous proteins are created in only  $n - 1$  duplications. Therefore, we refer to the current model as a *quadratic duplication model*, since the number of scored duplications is a quadratic function of the number of duplicates. While this might be desirable as being more restrictive on duplications, to be more consistent with the underlying biological processes, it can be replaced by a *linear duplication model*. In this model, each duplicate protein is penalized only once, based on its similarity with the paralog that is most similar to itself. This model can be incorporated into the alignment graph model of Section 3.3 with a simple modification of the algorithm that dynamically reassigns weights to edges that correspond to duplications.

## 4. EXPERIMENTAL RESULTS

### 4.1. Data and implementation

We implement the proposed algorithms in the C programming language and test on PPI networks that belong to three commonly studied eukaryotic organisms. The source code of the software is available at [www.cs.purdue.edu/homes/koyuturk/mawish/](http://www.cs.purdue.edu/homes/koyuturk/mawish/) along with detailed alignment results. The interaction data is downloaded from BIND (Bader *et al.*, 2001) and DIP (Xenarios *et al.*, 2002) molecular interaction databases. The statistics for the PPI networks of *S. cerevisiae* (yeast), *C. elegans* (nematode), and *D. melanogaster* (fruit fly) are shown in Table 1.

We align all pairs of these three organisms using a fixed set of parameters to be able to compare the results with each other. We set these parameters conservatively in order to obtain a compact set of

TABLE 1. DESCRIPTION OF ALIGNED PPI NETWORKS

<i>Organism</i>	<i># Proteins</i>	<i># Interactions</i>
<i>S. cerevisiae</i>	5157	18192
<i>C. elegans</i>	3345	5988
<i>D. melanogaster</i>	8577	28829

illustrative results. For any pair of PPI networks, we set the  $E$ -value threshold adaptively based on the estimated similarity scores so that the minimum similarity score for any pair of potential orthologs is 0.6. In other words, two proteins that belong to two different species are considered potentially orthologous only if they have a BLAST  $E$ -value less than 60% of ortholog pairs in COG. On the other hand, we set  $\bar{d} = 0.9$ ; i.e., two proteins in the same organism are considered potential in-paralogs only if they have BLAST  $E$ -value less than 90% of protein pairs in this organism that are in the same COG. For potential out-paralogs, we consider protein pairs that have a BLAST  $E$ -value less than 0.1 but greater than 10% of the ortholog pairs in COG. By setting these cut-off values on similarity score, we consider only the homologous protein pairs that have the highest positive or negative contribution on the alignment score. This eliminates noise to a certain extent while improving the computational efficiency. However, for more detailed analysis and discovery of loosely visible patterns, it may be necessary to relax and set these parameters based on the evolutionary distance between the two organisms being compared.

#### 4.2. Results and Discussion

We perform pairwise alignment of the three PPI networks by tuning the alignment parameters to  $\bar{\mu} = 1.0$ ,  $\bar{\nu} = 1.0$ , and  $\bar{\delta} = 0.1$ . Detailed statistics on alignment of the three pairs of eukaryotic PPI networks are shown in Table 2. In this table, we list the number of nodes in the alignment graph, nodes with at least one matched edge, matches, mismatches, and duplications in both organisms. The number of matches and the number of matched nodes are shown for two values of  $\bar{\Delta}$ , where only direct interactions  $\bar{\Delta} = 1$  and indirect interactions through a single protein  $\bar{\Delta} = 2$  are considered as matches. In practice, we eliminate all nodes that do not have any matching interactions from the alignment graph. As evident in the table, this improves the computational performance of the algorithm significantly.

Note that the parameters are set in an ad hoc manner for all results reported in this section, by manually adjusting the balance between normalizing parameters through repeated runs of the algorithm. A more reliable method for adjusting these parameters is to employ a learning heuristic that tunes these parameters in such a way that the conservation/divergence of known functional modules is captured by these parameters, as in the case of sequence alignment. However, this requires knowledge of biologically validated functional modules for both species being compared. We anticipate that as such data becomes available, an accepted set of parameters can easily be derived.

Alignment of *S. cerevisiae* PPI network with *D. melanogaster* PPI network results in identification of 412 conserved subnets. Ten of the conserved subnets with highest alignment scores are shown in Table 3. In this table, the rank of the identified subnet among all conserved subnets, its  $z$ -score computed according to the procedure in Section 3.4, the number of alignment nodes and corresponding number of proteins in each organism, total number of matches, mismatches, and duplications in each organism are shown in a row. In the subsequent row, we report the most dominant biological process identified according to the GO annotations of the proteins that are in the conserved network, along with the percentage of proteins that are associated with that biological process, for each organism. Similarly, sample high-scoring conserved subnets identified by the alignment of *S. cerevisiae* versus *C. elegans* and *C. elegans* versus *D. melanogaster* PPI networks are shown in Tables 4 and 5, respectively. In total, 83 conserved subnets are identified on *S. cerevisiae* and *C. elegans*, and 146 are identified on *C. elegans* and *D. melanogaster*.

TABLE 2. ALIGNMENT STATISTICS FOR THE THREE PAIRS OF EUKARYOTIC ORGANISMS<sup>a</sup>

Organism pair	# Nodes	# Matched nodes		# Matches		# Mismatches	# Duplications	
		$\bar{\Delta} = 1$	$\bar{\Delta} = 2$	$\bar{\Delta} = 1$	$\bar{\Delta} = 2$	$\bar{\Delta} = 1$	Org. 1	Org. 2
SC vs CE	2746	312	1230	412	3007	40262	6107	6886
SC vs DM	15884	1730	8622	2061	42781	1054241	6107	32670
CE vs DM	11805	491	3391	455	6626	205593	6886	32670

<sup>a</sup>For each alignment, the number of nodes in alignment graphs (# of orthologous pairs), number of nodes with at least one matched edge, number of matches, number of mismatches, and number of duplications for both organisms are shown. Number of mismatches for  $\bar{\Delta} = 2$  can be derived from other statistics.

TABLE 3. TEN HIGH-SCORING CONSERVED SUBNETS IDENTIFIED BY THE ALIGNMENT OF *S. Cerevisiae* AND *D. Melanogaster* PPI NETWORKS<sup>a</sup>

Rank	Score	z-Score	# Proteins	# Matches	# Mismatches	# Duplications
1	15.97	6.6	18 (16, 5)	28	6	(4, 0)
			Protein amino acid phosphorylation (69%)/JAK-STAT cascade (40%)			
2	13.93	3.7	13 (8, 7)	25	7	(3, 1)
			Endocytosis (50%)/calcium-mediated signaling (50%)			
5	8.22	13.5	9 (5, 3)	19	11	(1, 0)
			Invasive growth ( <i>sensu Saccharomyces</i> ) (100%)/oxygen and reactive oxygen species metabolism (33%)			
6	8.05	7.6	8 (5, 3)	12	2	(0, 1)
			Ubiquitin-dependent protein catabolism (100%)/mitosis (67%)			
8	6.83	12.4	6 (4, 4)	12	6	(0, 1)
			Protein amino acid phosphorylation (50%, 50%)			
10	6.75	13.7	10 (7, 3)	24	12	(0, 1)
			Ubiquitin-dependent protein catabolism (100%)			
14	5.69	8.7	11 (11, 2)	10	1	(0, 0)
			Regulation of progression through cell cycle (9%, 50%)			
21	4.36	6.2	9 (5, 4)	18	13	(0, 5)
			Cytokinesis (100%, 50%)			
22	4.22	3.9	7 (6, 6)	9	5	(1, 1)
			Protein folding (67%, 17%)			
30	3.76	39.6	6 (3, 5)	5	1	(0, 6)
			DNA replication initiation (100%, 80%)			

<sup>a</sup>The dominant biological process for each organism, in which the majority of proteins in the conserved subnet participate is shown in the second row.

While most of the conserved subnets are dominated by one particular process and the dominant processes are generally consistent across species, there also exist different processes in different organisms that are mapped to each other by the discovered alignments. This illustrates that the comparative analysis of PPI networks is effective in not only identifying particular functional modules, pathways, and complexes, but also in discovering relationships between different processes in separate organisms and crosstalk between known functional modules and pathways.

A selection of interesting conserved subnets is shown in Fig. 5. The alignments in the figure illustrate that the alignment algorithm takes into account the conservation of interactions in addition to sequence similarity while mapping orthologous proteins to each other. In all of the alignments shown in the figure, the interactions of proteins that belong to the same orthologous group are highly conserved, suggesting relatively recent duplications.

Detailed examination of the conserved subnets in *S. cerevisiae* and *D. melanogaster* shows that many of them do correspond to functional modules. There are multiple instances of 20S proteasome (10,11). All seven of the alpha subunits in the 20S proteasome, a subcomplex of the 26S proteasome involved in protein degradation, are present in the alignment #10 (Groll *et al.*, 1997). In addition, there is a subnet for the proteasome regulatory particle (6) (Fu *et al.*, 2001) as well as one for calcium induced pathways (2). The method also detected a number of components involved in calcium-dependent stress-activated signaling pathways (Cmd1, Cna1, Cna2, and Cnb1) as well as those associated with bud growth of yeast (Cmd1, Myo2, and Myo4) in alignment #2 (Cyert *et al.*, 2001). Many of the subnets found for yeast are overlapping, possibly reflecting the fact that drosophila uses a functional module in various contexts.

In some cases, the self-interaction of a single protein in one organism is aligned with a clique of interactions between its orthologs that are part of a particular module. For example, in one alignment, five proteasome regulatory particle proteins (Rpt1, Rpt3, Rpt4, Rpt5, Rpt6) are mapped to one protein (Rpt4) in drosophila, while in other alignments the same group of proteins are mapped to a different set of proteins in the drosophila network. This may be due to missing interactions in one of the networks because of incompleteness or irregularities in the interaction data, including coding of observed interactions

TABLE 4. FIVE HIGH-SCORING CONSERVED SUBNETS IDENTIFIED BY THE ALIGNMENT OF *S. Cerevisiae* AND *C. Elegans* PPI NETWORKS

Rank	Score	z-Score	# Proteins	# Matches	# Mismatches	# Duplications
1	36.14	7.8	13 (5, 3)	65	24	(0, 3)
	Ubiquitin-dependent protein catabolism (100%)/reproduction (100%)					
2	8.47	6.5	20 (11, 5)	19	4	(1, 1)
	Protein amino acid phosphorylation (82%, 40%)					
3	6.28	10.1	8 (6, 3)	21	12	(0, 0)
	Ubiquitin-dependent protein catabolism (100%, 100%)					
7	3.23	4.9	7 (7, 6)	7	2	(0, 0)
	Glyoxylate cycle (14%, 17%)					
8	3.23	80.1	4 (3, 3)	4	1	(1, 1)
	Mismatch repair (67%, 67%)					

TABLE 5. FIVE HIGH-SCORING CONSERVED SUBNETS IDENTIFIED BY THE ALIGNMENT OF *C. Elegans* AND *D. Melanogaster* PPI NETWORKS


Rank	Score	z-Score	# Proteins	# Matches	# Mismatches	# Duplications
1	26.75	19.9	17 (4, 9)	52	4	(1, 4)
	Thermosensory behavior (25%)/regulation of transcription from RNA polymerase II promoter (44%)					
2	4.65	31.6	9 (5, 3)	8	0	(2, 1)
	Translational initiation (60%, 67%)					
4	4.37	10.7	11 (3, 6)	10	1	(1, 4)
	Determination of adult life span (33%, 67%)					
5	4.29	16.4	5 (4, 4)	6	0	(1, 1)
	Regulation of transcription, DNA-dependent (50%, 25%)					
6	4.00	12.2	6 (4, 6)	8	2	(0, 2)
	Signal transduction (50%, 17%)					

into the databases (e.g., spoke versus matrix model [Scholtens *et al.*, 2005]). By adjusting the mismatch and duplication coefficients, however, it is possible to make positive duplications dominate the negative mismatches caused by these missing interactions. This may be considered as a desirable feature of our algorithm, in the sense that it allows flexibility for trading off conservation of interactions with conservation of proteins. However, it might also be considered a drawback, since setting the parameters in this manner causes overrepresentation of somewhat distant proteins in one conserved subnet on another side of the network.

Based on these results, we establish pairwise alignment of PPI networks as a tool for not only identifying conserved modules, but also assessing functional differences and similarities of homologous proteins based on shared and missing interactions. Moreover, alignment results provide a means for discovery of new functional modules in relatively less studied organisms through mapping of functions at a modular level rather than at the level of single protein homologies.

## 5. CONCLUSION

This paper presents a framework for local alignment of protein interaction networks. The framework is guided by theoretical models of evolution of these networks. The model is based on discovering sets of proteins that induce conserved subnets based on scoring match and mismatch of interactions, and duplication of proteins. An implementation of the proposed algorithm reveals that this framework is successful in uncovering conserved substructures in protein interaction data.



**FIG. 5.** Sample conserved subnets identified by the alignment algorithm. Orthologous and paralogous proteins are either vertically aligned or connected by blue dotted lines. Existing interactions are shown by green solid lines, missing interactions that have an orthologous counterpart are shown by red dashed lines. The rank of each alignment in the set of alignments discovered for the respective pair of organisms is indicated in its label.

## REFERENCES

- Altschul, S.F., Madden, T.L., Schffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucl. Acids Res.* 25(17), 3389–3402.
- Ashtana, S., King, O.D., Gibbons, F.D., and Roth, F.P. 2004. Predicting protein complex membership using probabilistic network reliability. *Genome Res.* 14, 1170–1175.
- Bader, G.D., Donalson, I., Wolting, C., Quellet, B.F., Pawson, T., and Hogure, C.W. 2001. BIND—The Biomolecular Interaction Network Database. *Nucl. Acids Res.* 29(1), 242–245.
- Bader, G.D., and Hogue, C.W.V. 2003. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* 4(2).
- Bader, J.S. 2003. Greedily building protein networks with confidence. *Bioinformatics* 19, 1869–1874.
- Barabási A., and Albert, R. 1999. Emergence of scaling in random networks. *Science* 286, 509–512.
- Berg, J., Lässig, M., and Wagner, A. 2004. Structure and evolution of protein interaction networks: A statistical model for link dynamics and gene duplications. *BMC Evol. Biol.* 5(51).

- Chung, F., Lu, L., Dewey, T.G., and Galas, D.J. 2003. Duplication models for biological networks. *J. Comp. Biol.* 10(5), 677–687.
- Chung, F., Lu, L., and Vu, V. 2003. Spectra of random graphs with given expected degrees. *PNAS* 100(11), 6313–6318.
- Cyert, M.S. *et al.* 2001. Genetic analysis of calmodulin and its targets in *Saccharomyces cerevisiae*. *Ann. Rev. Genet.* 35, 647–672.
- Dandekar, T., Schuster, S., Snel, B., Huynen, M., and Bork, P. 1999. Pathway alignment: Application to the comparative analysis of glycolytic enzymes. *Biochem. J.* 343, 115–124.
- Eisenberg, E., and Levanon, Y. 2003. Preferential attachment in the protein network evolution. *Phys. Rev. Lett.* 91(13), 138701.
- Feige, U., Peleg, D., and Kortsarz, G. 2001. The dense k-subgraph problem. *Algorithmica* 29(3), 410–421.
- Fraser, H.B., Hirsh, A.E., Steinmetz, L.M., Scharfe, C., and Feldman, M.W. 2002. Evolutionary rate in the protein interaction network. *Science* 296(5568), 750–752.
- Fu, H. *et al.* 2001. Subunit interaction maps for the regulatory particle of the 26S proteasome and the COP9 signalosome. *EMBO J.* 20(24), 7096–7107.
- Gavin, A.C. *et al.* 2002. Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* 415(6868), 141–147.
- Gilchrist, M.A., Salter, L.A., and Wagner, A. 2003. A statistical framework for combining and interpreting proteomic datasets. *Bioinformatics* 20(5), 689–700.
- Groll, M. *et al.* 1997. Structure of 20S proteasome from yeast at 2.4 Å resolution. *Nature* 386(6624), 463–471.
- Hartwell, L.H., Hopfield, J.J., Leibler, S., and Murray, A.W. 1999. From molecular to modular cell biology. *Nature* 402, C47–C51.
- Hassin, R., Rubinstein, S., and Tamir, A. 1997. Approximation algorithms for maximum dispersion. *Oper. Res. Lett.* 21, 133–137.
- Ho, Y. *et al.* 2002. Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* 415, 180–183.
- Hu, H., Yan, X., Huang, Y., Han, J., and Zhou, X.J. 2005. Mining coherent dense subgraphs across massive biological networks for functional discovery. *Bioinformatics* 21(Suppl. 1), i213–i221.
- Ito, T., Chiba, T., Ozawa, R., Yoshida, M., Hattori, M., and Sakaki, Y. 2001. A comprehensive two-hybrid analysis to explore the yeast protein interactome. *PNAS* 98(8), 4569–4574.
- Jansen, R., Yu, H., Greenbaum, D., *et al.* 2003. A Bayesian networks approach for predicting protein–protein interactions from genomic data. *Science* 302, 449–453.
- Kelley, B.P., Sharan, R., Karp, R.M., Sittler, T., Root, D.E., Stockwell, B.R., and Ideker, T. 2003. Conserved pathways withing bacteria and yeast as revealed by global protein network alignment. *PNAS* 100(20), 11394–11399.
- Kelley, B.P., Yuan, B., Lewitter, F., Sharan, R., Stockwell, B.R., and Ideker, T. 2004. PathBLAST: A tool for alignment of protein interaction networks. *Nucl. Acids Res.* 32, W83–W88.
- Kernighan, B.W., and Lin, S. 1970. An efficient heuristic procedure for partitioning graphs. *The Bell System Tech. J.* 49(2), 291–307.
- Koyutürk, M., Grama, A., and Szpankowski, W. 2004. An efficient algorithm for detecting frequent subgraphs in biological networks. *Bioinformatics Suppl. 12th Int. Conf. Intel. Sys. Mol. Bio. (ISMB '04)*, i200–i207.
- Koyutürk, M., Grama, A., and Szpankowski, W. 2005. Pairwise local alignment of protein interaction networks guided by models of evolution. *RECOMB 2005, Lecture Notes in Bioinformatics*, 3500, 48–65.
- Koyutürk, M., Grama, A., and Szpankowski, W. 2006. Assessing significance of connectivity and conservation in protein interaction networks. *RECOMB 2006, Lecture Notes in Bioinformatics*, 3909, 45–59.
- Kunin, V., Pereira-Leal, J.B., and Ouzounis, C.A. 2004. Functional evolution of the yeast protein interaction network. *Mol. Biol. Evol.* 21(7), 1171–1176.
- Lotem, E.Y., Sattath, S., Kashtan, N., Itzkovitz, S., Milo, R., Pinter, R.Y., Alon, U., and Margalit, H. 2004. Network motifs in integrated cellular networks of transcription-regulation and protein–protein interaction. *PNAS* 101(16), 5934–5939.
- Pastor-Satorras, R., Smith, E., and Solé, R.V. 2003. Evolving protein interaction networks through gene duplication. *J. Theor. Biol.* 222, 199–210.
- Pinter, R.Y., Rokhlenko, O., Yeger-Lotem, E., and Ziv-Ukelson, M. 2005. Alignment of metabolic pathways. *Bioinformatics*. In press.
- Qin, H., Lu, H.H.S., Wu, W.B., and Li, W. 2003. Evolution of the yeast protein interaction network. *PNAS* 100(22), 12820–12824.
- Remm, M., Storm, C.E.V., and Sonnhammer, E.L.L. 2001. Automatic clustering of orthologs and in-paralogs from pairwise species comparisons. *J. Mol. Biol.* 314, 1041–1052.
- Scholtens, D., Vidal, M., and Gentleman, R. 2005. Local modeling of global interactome networks. *Bioinformatics* 21(17), 3548–3557.

- Sharan, R., Ideker, T., Kelley, B.P., Shamir, R., and Karp, R.M. 2004. Identification of protein complexes by comparative analysis of yeast and bacterial protein interaction data. *8th Intl. Conf. Res. Comp. Mol. Biol. (RECOMB '04)*, 282–289.
- Sharan, R., Suthram, S., Kelley, R.M., Kuhn, T., McCuine, S., Uetz, P., Sittler, T., Karp, R.M., and Ideker, T. 2005. Conserved patterns of protein interaction in multiple species. *PNAS* 102(6), 1974–1979.
- Smith, T.F., and Waterman, M.S. 1981. Identification of common molecular subsequences. *J. Mol. Biol.* 147(1), 195–197.
- Tatusov, R.L., Galperin, M.Y., Natale, D.A., and Koonin, E.V. 2000. The COG database: A tool for genome-scale analysis of protein functions and evolution. *Nucl. Acids Res.* 28, 33–36.
- Thompson, J.D., Higgins, D.G., and Gibson, T.J. 1994. CLUSTAL-W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids Res.* 22, 4673–4680.
- Titz, B., Schlesner, M., and Uetz, P. 2004. What do we learn from high-throughput protein interaction data? *Exp. Rev. Prot.* 1(1), 111–121.
- Tohsato, Y., Matsuda, H., and Hashimoto, A. 2000. A multiple alignment algorithm for metabolic pathway analysis using enzyme hierarchy. *8th Int. Conf. Intel. Sys. Mol. Biol. (ISMB '00)*, 376–383.
- Tornow, S., and Mewes, H.W. 2003. Functional modules by relating protein interaction networks and gene expression. *Nucl. Acids Res.* 31(21), 6283–6289.
- Vázquez, A., Flammini, A., Maritan, A., and Vespignani, A. 2003. Modeling of protein interaction networks. *ComplexUs* 1, 38–44.
- Wagner, A. 2001. The yeast protein interaction network evolves rapidly and contains few redundant duplicate genes. *Mol. Biol. Evol.* 18(7), 1283–1292.
- Wagner, A. 2003. How the global structure of protein interaction networks evolves. *Proc. R. Soc. Lond. Biol. Sci.* 270(1514), 457–466.
- Wuchty, S. 2004. Evolution and topology in the yeast protein interaction network. *Genome Res.* 14, 1310–1314.
- Wuchty, S., Oltvai, Z.N., and Barabási, A.L. 2003. Evolutionary conservation of motif constituents in the yeast protein interaction network. *Nature Genet.* 35(2), 176–179.
- Xenarios, I., Salwinski, L., Duan, X.J., Higney, P., Kim, S., and Eisenberg, D. 2002. DIP: The Database of Interacting Proteins. A research tool for studying cellular networks of protein interactions. *Nucl. Acids Res.* 30, 303–305.

Address correspondence to:  
Mehmet Koyutürk  
Computer Sciences Building  
250 North University Street  
West Lafayette, IN 47907

E-mail: koyuturk@cs.purdue.edu