

Computing Bottleneck Distance for 2-D Interval Decomposable Modules

Tamal K. Dey* Cheng Xin*

Abstract

Computation of the interleaving distance between persistence modules is a central task in topological data analysis. For 1-D persistence modules, thanks to the isometry theorem, this can be done by computing the bottleneck distance with known efficient algorithms. The question is open for most n -D persistence modules, $n > 1$, because of the well recognized complications of the indecomposables. Here, we consider a reasonably complicated class called *2-D interval decomposable* modules whose indecomposables may have a description of non-constant complexity. We present a polynomial time algorithm to compute the bottleneck distance for these modules from indecomposables, which bounds the interleaving distance from above, and give another algorithm to compute a new distance called *dimension distance* that bounds it from below.

1 Introduction

Persistence modules have become an important object of study in topological data analysis in that they serve as an intermediate between the raw input data and the output summarization with persistence diagrams. The classical persistence theory [18] for \mathbb{R} -valued functions produces one dimensional (1-D) persistence modules, which is a sequence of vector spaces (homology groups with a field coefficient) with linear maps over \mathbb{R} seen as a poset. It is known that [16, 25], this sequence can be decomposed uniquely into a set

*Department of Computer Science and Engineering, The Ohio State University. dey.8@osu.edu, xin.108@buckeyemail.osu.edu

of intervals called *bars* which is also represented as points in \mathbb{R}^2 called the persistence diagrams [15]. The space of these diagrams can be equipped with a metric d_B called the *bottleneck distance*. Cohen-Steiner et al. [15] showed that d_B is bounded from above by the input function perturbation measured in infinity norm. Chazal et al. [12] generalized the result by showing that the bottleneck distance is bounded from above by a distance d_I called the *interleaving distance* between two persistence modules; see also [6, 8, 17] for further generalizations. Lesnick [21] (see also [2, 13]) established the isometry theorem which showed that indeed $d_I = d_B$. Consequently, d_I for 1-D persistence modules can be computed exactly by efficient algorithms known for computing d_B ; see e.g. [18, 19]. The status however is not so well settled for multidimensional (n -D) persistence modules [9] arising from \mathbb{R}^n -valued functions.

Extending the concept from 1-D modules, Lesnick defined the interleaving distance for multidimensional (n -D) persistence modules, and proved its stability and universality [21]. The definition of the bottleneck distance, however, is not readily extensible mainly because the bars for finitely presented n -D modules called *indecomposables* are far more complicated though are guaranteed to be essentially unique by Krull-Schmidt theorem [1]. Nonetheless, one can define d_B as the supremum of the pairwise interleaving distances between indecomposables, which in some sense generalizes the concept in 1-D due to the isometry theorem. Then, straightforwardly, $d_I \leq d_B$ as observed in [7], but the converse is not necessarily true. For some special cases, results in the converse direction have started to appear. Botnan and Lesnick [7] proved that, in 2-D, $d_B \leq \frac{5}{2}d_I$ for what they called block decomposable modules. Bjerkevic [4] improved this result to $d_B \leq d_I$. Furthermore, he extended it by proving that $d_B \leq (2n - 1)d_I$ for rectangle decomposable n -D modules and $d_B \leq (n - 1)d_I$ for free n -D modules. He gave an example for exactness of this bound when $n = 2$.

Unlike 1-D modules, the question of estimating d_I for n -D modules through efficient algorithms is largely open [5]. Multi-dimensional matching distance introduced in [10] provides a lower bound to interleaving distance [20] and can be approximated within any error threshold by algorithms proposed in [3, 11]. But, it cannot provide an upper bound like d_B . For free, block, rectangle, and triangular decomposable modules, one can compute d_B by computing pairwise interleaving distances between indecomposables in constant time because they have a description of constant complexity. Due to the results mentioned earlier, d_I can be estimated within a constant or dimension-dependent factors by

computing d_B for these modules. It is not obvious how to do the same for the larger class of interval decomposable modules mentioned in the literature [4, 7] where indecomposables may not have constant complexity. These are modules whose indecomposables are bounded by “stair-cases”. Our main contribution is a polynomial time algorithm that, given indecomposables, computes d_B exactly for 2-D interval decomposable modules. The algorithm draws upon various geometric and algebraic analysis of the interval decomposable modules that may be of independent interest. It is known that no lower bound in terms of d_B for d_I may exist for these modules [7]. To this end, we complement our result by proposing a distance d_0 called *dimension distance* that is efficiently computable and satisfies the condition $d_0 \leq d_I$.

2 Persistence modules

Our goal is to compute the bottleneck distance between two 2-D interval decomposable modules. The bottleneck distance, originally defined for 1-D persistence modules [15] (also see [2]), and later extended to multi-dimensional persistence modules [7] is known to bound the interleaving distance between two persistence modules from above.

Let \mathbb{k} be a field, \mathbf{Vec} be the category of vector spaces over \mathbb{k} , and \mathbf{vec} be the subcategory of finite dimensional vector spaces. In what follows, for simplicity, we assume $\mathbb{k} = \mathbb{Z}/2\mathbb{Z}$.

Definition 1 (Persistence module). *Let \mathbb{P} be a poset category. A \mathbb{P} -indexed persistence module is a functor $M : \mathbb{P} \rightarrow \mathbf{Vec}$. If M takes values in \mathbf{vec} , we say M is pointwise finite dimensional (p.f.d). The \mathbb{P} -indexed persistence modules themselves form another category where the natural transformations between functors constitute the morphisms.*

Here we consider the poset category to be \mathbb{R}^n with the standard partial order and all modules to be p.f.d. We call \mathbb{R}^n -indexed persistence modules as n -dimensional persistence modules, n -D modules in short. The category of n -D modules is denoted as $\mathbb{R}^n\text{-mod}$. For an n -D module $M \in \mathbb{R}^n\text{-mod}$, we use notation $M_x := M(x)$ and $\rho_{x \rightarrow y}^M := M(x \leq y)$.

Definition 2 (Shift). *For any $\delta \in \mathbb{R}$, we denote $\vec{\delta} = \delta \cdot \sum e_i$, where $\{e_i\}_{i=1}^n$ is the standard basis of \mathbb{R}^n . We define a shift functor $(\cdot)_{\rightarrow \delta} : \mathbb{R}^n\text{-mod} \rightarrow \mathbb{R}^n\text{-mod}$ where $M_{\rightarrow \delta} := (\cdot)_{\rightarrow \delta}(M)$ is given by $M_{\rightarrow \delta}(x) = M(x + \vec{\delta})$ and $M_{\rightarrow \delta}(x \leq y) = M(x + \vec{\delta} \leq y + \vec{\delta})$. In words, $M_{\rightarrow \delta}$ is the module M shifted diagonally by $\vec{\delta}$.*

The following definition of interleaving taken from [23] adapts the original definition designed for 1-D modules in [13] to n -D modules.

Definition 3 (Interleaving). *For two persistence modules M and N , and $\delta \geq 0$, a δ -interleaving between M and N are two families of linear maps $\{\phi_x : M_x \rightarrow N_{x+\delta}\}_{x \in \mathbb{R}^n}$ and $\{\psi_x : N_x \rightarrow M_{x+\delta}\}_{x \in \mathbb{R}^n}$ satisfying the following two conditions (see Appendix A for commutative diagrams):*

- $\forall x \in \mathbb{R}^n, \rho_{x \rightarrow x+2\delta}^M = \psi_{x+\delta} \circ \phi_x$ and $\rho_{x \rightarrow x+2\delta}^N = \phi_{x+\delta} \circ \psi_x$
- $\forall x \leq y \in \mathbb{R}^n, \phi_y \circ \rho_{x \rightarrow y}^M = \rho_{x \rightarrow y}^N \circ \phi_x$ and $\psi_y \circ \rho_{x \rightarrow y}^N = \rho_{x \rightarrow y}^M \circ \psi_x$ symmetrically

If such a δ -interleaving exists, we say M and N are δ -interleaved. We call the first condition triangular commutativity and the second condition square commutativity.

Definition 4 (Interleaving distance). *Define the interleaving distance between modules M and N as $d_I(M, N) = \inf_{\delta} \{M \text{ and } N \text{ are } \delta\text{-interleaved}\}$. We say M and N are ∞ -interleaved if they are not δ -interleaved for any $\delta \in \mathbb{R}^+$, and assign $d_I(M, N) = \infty$.*

Definition 5 (Matching). *A matching $\mu : A \dashrightarrow B$ between two multisets A and B is a partial bijection, that is, $\mu : A' \rightarrow B'$ for some $A' \subseteq A$ and $B' \subseteq B$. We say $\text{im } \mu = B'$, $\text{coim } \mu = A'$.*

For the next definition [7], we call a module δ -trivial if $\rho_{x \rightarrow x+\delta}^M = 0$ for all $x \in \mathbb{R}^n$.

Definition 6 (Bottleneck distance). *Let $M \cong \bigoplus_{i=1}^m M_i$ and $N \cong \bigoplus_{j=1}^n N_j$ be two persistence modules, where M_i and N_j are indecomposable submodules of M and N respectively. Let $I = \{1, \dots, m\}$ and $J = \{1, \dots, n\}$. We say M and N are δ -matched for $\delta \geq 0$ if there exists a matching $\mu : I \dashrightarrow J$ so that, (i) $i \in I \setminus \text{coim } \mu \implies M_i$ is 2δ -trivial, (ii) $j \in J \setminus \text{im } \mu \implies N_j$ is 2δ -trivial, and (iii) $i \in \text{coim } \mu \implies M_i$ and $N_{\mu(i)}$ are δ -interleaved.*

The bottleneck distance is defined as

$$d_B(M, N) = \inf\{\delta \mid M \text{ and } N \text{ are } \delta\text{-matched}\}.$$

The following fact observed in [7] is straightforward from the definition.

Fact 7. $d_I \leq d_B$.

2.1 Interval decomposable modules

Persistence modules whose indecomposables are interval modules (Definition 9) are called *interval decomposable modules*, see for example [7]. To account for the boundaries of free modules, we enrich the poset \mathbb{R}^n by adding points at $\pm\infty$ and consider the poset $\bar{\mathbb{R}}^n = \bar{\mathbb{R}} \times \dots \times \bar{\mathbb{R}}$ where $\bar{\mathbb{R}} = \mathbb{R} \cup \{\pm\infty\}$ with the usual additional rule $a \pm \infty = \pm\infty$.

Definition 8. *An interval is a subset $\emptyset \neq I \subset \bar{\mathbb{R}}^n$ that satisfies the following:*

1. *If $p, q \in I$ and $p \leq r \leq q$, then $r \in I$;*
2. *If $p, q \in I$, then there exists a sequence $(p_1, p_2, \dots, p_{2m}) \in I$ for some $m \in \mathbb{N}$ such that $p \leq p_1 \geq p_2 \leq p_3 \geq \dots \geq p_{2m} \leq q$. We call the sequence $(p = p_0, p_1, p_2, \dots, p_{2m}, p_{2m+1} = q)$ a path from p to q (in I).*


In what follows, we fix the dimension $n = 2$. Let \bar{I} denote the closure of an interval I in the standard topology of $\bar{\mathbb{R}}^2$. The lower and upper boundaries of I are defined as

$$L(I) = \{x = (x_1, x_2) \in \bar{I} \mid \forall y = (y_1, y_2) \text{ with } y_1 < x_1 \text{ and } y_2 < x_2 \implies y \notin I\}$$

$$U(I) = \{x = (x_1, x_2) \in \bar{I} \mid \forall y = (y_1, y_2) \text{ with } y_1 \succeq x_1 \text{ and } y_2 \succeq x_2 \implies y \notin I\}.$$

See the figure below. Let $B(I) = L(I) \cup U(I)$.

We say an interval I is discretely presented if its boundary consists of a finite set of horizontal and vertical line segments called edges, with end points called vertices, which satisfy the following conditions: (i) every vertex is incident to either a single edge or to a horizontal and a vertical edge, (ii) no vertex appears in the interior of an edge. We denote the set of edges and vertices with $E(I)$ and $V(I)$ respectively.



According to this definition, $\bar{\mathbb{R}}^2$ is an interval with boundary $B(\bar{\mathbb{R}}^2)$ that consists of all the points with at least one coordinate ∞ . The vertex set $V(\bar{\mathbb{R}}^2)$ consists of four corners of the infinitely large square $\bar{\mathbb{R}}^2$ with coordinates $(\pm\infty, \pm\infty)$.

Definition 9 (Interval module). *A 2-D **interval persistence module**, or interval module in short, is a persistence module M that satisfies the following*

condition: for some interval $I_M \subseteq \bar{\mathbb{R}}^2$, called the interval of M ,

$$M_x = \begin{cases} \mathbb{k} & \text{if } x \in I_M \\ 0 & \text{otherwise} \end{cases} \quad \rho_{x \rightarrow y}^M = \begin{cases} \mathbb{1} & \text{if } x, y \in I_M \\ 0 & \text{otherwise} \end{cases}$$

It is known that an interval module is indecomposable [21].

Definition 10 (Interval decomposable module). *A 2-D interval decomposable module is a persistence module that can be decomposed into interval modules. We say a 2-D interval decomposable module is finitely presented if it can be decomposed into finitely many interval modules whose intervals are discretely presented.*

3 Algorithm to compute d_B

Given the intervals of the indecomposables (interval modules) as input, an approach based on bipartite-graph matching is well known for computing the bottleneck distance $d_B(M, N)$ between two 1-D persistence modules M and N [18]. This approach constructs a bi-partite graph G out of the intervals of M and N and their pairwise interleaving distances including the distances to zero modules. If these distance computations take $O(C)$ time in total, the algorithm for computing d_B takes time $O(m^{\frac{5}{2}} \log m + C)$ if M and N together have m indecomposables altogether. Given indecomposables (say computed by Meat-Axe [22]), this approach is readily extensible to the n -D modules if one can compute the interleaving distance between any pair of indecomposables including the zero modules. To this end, we present an algorithm to compute the interleaving distance between two interval modules M_i and N_j with t_i and t_j vertices respectively on their intervals in $O((t_i + t_j) \log(t_i + t_j))$ time. This gives a total time of $O(m^{\frac{5}{2}} \log m + \sum_{i,j} (t_i + t_j) \log(t_i + t_j)) = O(m^{\frac{5}{2}} \log m + t^2 \log t)$ where t is the number of vertices over all input intervals.

Now we focus on computing the interleaving distance between two given intervals. Given two intervals I_M and I_N with t vertices, this algorithm searches a value δ so that there exists two families of linear maps from M to $N_{\rightarrow \delta}$ and from N to $M_{\rightarrow \delta}$ respectively which satisfy both triangular and square commutativity. This search is done with a binary probing. For a chosen δ from a candidate set of $O(t)$ values, the algorithm determines the direction of the search by checking two conditions called *trivializability* and *validity* on the intersections of modules M and N .

Definition 11 (Intersection module). For two interval modules M and N with intervals I_M and I_N respectively let $I_Q = I_M \cap I_N$, which is a disjoint union of intervals, $\coprod I_{Q_i}$. The intersection module Q of M and N is $Q = \bigoplus Q_i$, where Q_i is the interval module with interval I_{Q_i} . That is,

$$Q_x = \begin{cases} \mathbb{k} & \text{if } x \in I_M \cap I_N \\ 0 & \text{otherwise} \end{cases} \quad \text{and for } x \leq y, \rho_{x \rightarrow y}^Q = \begin{cases} \mathbb{1} & \text{if } x, y \in I_M \cap I_N \\ 0 & \text{otherwise} \end{cases}$$

From the definition we can see that the support of Q , $\text{supp}(Q)$, is $I_M \cap I_N$. We call each Q_i an intersection component of M and N . Write $I := I_{Q_i}$ and consider $\phi : M \rightarrow N$ to be any morphism in the following proposition which says that ϕ is constant on I .

Proposition 12. $\phi|_I \equiv a \cdot \mathbb{1}$ for some $a \in \mathbb{k} = \mathbb{Z}/2$.

Proof.

$$\begin{array}{ccc} M_{p_i} & \xrightarrow{\mathbb{1}} & M_{p_{i+1}} & & M_{p_i} & \xleftarrow{\mathbb{1}} & M_{p_{i+1}} \\ \phi_{p_i} \downarrow & & \downarrow \phi_{p_{i+1}} & & \phi_{p_i} \downarrow & & \downarrow \phi_{p_{i+1}} \\ N_{p_i} & \xrightarrow{\mathbb{1}} & N_{p_{i+1}} & & N_{p_i} & \xleftarrow{\mathbb{1}} & N_{p_{i+1}} \end{array}$$

For any $x, y \in I$, consider a path $(x = p_0, p_1, p_2, \dots, p_{2m}, p_{2m+1} = y)$ in I from x to y and the commutative diagrams above for $p_i \leq p_{i+1}$ (left) and $p_i \geq p_{i+1}$ (right) respectively. Observe that $\phi_{p_i} = \phi_{p_{i+1}}$ in both cases due to the commutativity. Inducting on i , we get that $\phi(x) = \phi(y)$. \square


Definition 13 (Valid intersection). An intersection component Q_i is (M, N) -valid if for each $x \in I_{Q_i}$ the following two conditions hold (see figure below):

(i) $y \leq x$ and $y \in I_M \implies y \in I_N$, and (ii) $z \geq x$ and $z \in I_N \implies z \in I_M$

Proposition 14. Let $\{Q_i\}$ be a set of intersection components of M and N with intervals $\{I_{Q_i}\}$. Let $\{\phi_x\} : M \rightarrow N$ be the family of linear maps defined as $\phi_x = \mathbb{1}$ for all $x \in I_{Q_i}$ and $\phi_x = 0$ otherwise. Then ϕ is a morphism if and only if every Q_i is (M, N) -valid.

See the proof in Appendix A.

We focus on the interval modules with discretely presented intervals (figure on right). They belong to persistence modules called finitely presented persistence modules studied previously in [21]. For an interval module M , let \overline{M} be the interval module defined on the closure $\overline{I_M}$. To avoid complication in this exposition, we assume that the upper and lower boundaries of every interval module meet exactly at two points. We also assume that every interval module has closed intervals which is justified by the following proposition (proof in Appendix A).



Proposition 15. $d_I(M, N) = d_I(\overline{M}, \overline{N})$.

From the definition of boundaries of intervals, the following proposition is immediate.

Proposition 16. *Given an interval I and any point $x = (x_1, x_2) \in I \setminus (I \cap B(\overline{\mathbb{R}}^2))$, we have $x \in L(I) \iff \forall \epsilon > 0, x - \vec{\epsilon} \notin I$. Similarly, we have $x \in U(I) \iff \forall \epsilon > 0, x + \vec{\epsilon} \notin I$.*

Definition 17 (Diagonal projection and distance). *Let I be an interval and $x \in \overline{\mathbb{R}}^2$. For $x \in \mathbb{R}^2 \subseteq \overline{\mathbb{R}}^2$, let Δ_x denote the line called diagonal with slope 1 that passes through x . We define (see Figure 1)*

$$\text{dl}(x, I) = \begin{cases} \min_{y \in \Delta_x \cap I} \{d_\infty(x, y) := |x - y|_\infty\} & \text{if } \Delta_x \cap I \neq \emptyset \\ +\infty & \text{otherwise.} \end{cases}$$

In case $\Delta_x \cap I \neq \emptyset$, define $\pi_I(x)$, called the projection point of x on I , to be the point $y \in \Delta_x \cap I$ where $\text{dl}(x, I) = d_\infty(x, y)$. For $x \in B(\overline{\mathbb{R}}^2) \setminus V(\overline{\mathbb{R}}^2)$, Δ_x is defined to be the edge in $E(\overline{\mathbb{R}}^2)$ containing x . Define $\text{dl}(x, I)$ and $\pi_I(x)$ accordingly. For $x \in V(\overline{\mathbb{R}}^2)$, we set $\pi_I(x) = x$ if and only if $x \in I$. Then, $\text{dl}(x, I) = 0$ if $x \in I$ and $\text{dl}(x, I) = +\infty$ otherwise.

Notice that upper and lower boundaries of an interval are also intervals by definition. With this understanding, following properties of dl are obvious from the above definition.

Fact 18. (i) *For any $x \in I_M$,*

$$\text{dl}(x, U(I_M)) = \sup_{\delta \in \overline{\mathbb{R}}} \{x + \vec{\delta} \in I_M\} \text{ and } \text{dl}(x, L(I_M)) = \sup_{\delta \in \overline{\mathbb{R}}} \{x - \vec{\delta} \in I_M\}.$$




Figure 1: $d = \text{dl}(x, I)$, $y = \pi_I(x)$, $d' = \text{dl}(x', L(I))$ (left); $d = \text{dl}(x, I)$ and $d' = \text{dl}(x', U(I))$ are defined on the left edge of $B(\bar{\mathbb{R}}^2)$ (middle); Q is $d'_{(M,N)}$ - and $d_{(N,M)}$ -trivializable (right)

(ii) Let $L = L(I_M)$ or $U(I_M)$ and let x, x' be two points such that $\pi_L(x), \pi_L(x')$ both exist. If x and x' are on some same horizontal, vertical, or diagonal line, then $|\text{dl}(x, L) - \text{dl}(x', L)| \leq d_\infty(x, x')$.

Set $VL(I) := V(I) \cap L(I)$, $EL(I) := E(I) \cap L(I)$, $VU(I) := V(I) \cap U(I)$, and $EU(I) := E(I) \cap U(I)$. Following proposition is proved in Appendix A.

Proposition 19. For an intersection component Q of M and N with interval I , the following conditions are equivalent:

- (1) Q is (M, N) -valid.
- (2) $L(I) \subseteq L(I_M)$ and $U(I) \subseteq U(I_N)$.
- (3) $VL(I) \subseteq L(I_M)$ and $VU(I) \subseteq U(I_N)$.

Definition 20 (Trivializable intersection). Let Q be a connected component of the intersection of two modules M and N . For each point $x \in I_Q$, define

$$d_{triv}^{(M,N)}(x) = \max\{\text{dl}(x, U(I_M))/2, \text{dl}(x, L(I_N))/2\}.$$

For $\delta \geq 0$, we say a point x is $\delta_{(M,N)}$ -trivializable if $d_{triv}^{(M,N)}(x) < \delta$. We say an intersection component Q is $\delta_{(M,N)}$ -trivializable if each point in I_Q is $\delta_{(M,N)}$ -trivializable (Figure 1).

Following proposition discretizes the search for trivialisability (proof in Appendix A).

Proposition 21. *An intersection component Q is $\delta_{(M,N)}$ -trivializable if and only if every vertex of Q is $\delta_{(M,N)}$ -trivializable.*

Recall that for two modules to be δ -interleaved, we need two families of linear maps satisfying both triangular commutativity and square commutativity. For a given δ , Theorem 23 below provides criteria which ensure that such linear maps exist. In our algorithm, we make sure that these criteria are verified.

Given an interval module M and the diagonal line Δ_x for any $x \in \bar{\mathbb{R}}^2$, there is a 1-dimensional persistence module $M|_{\Delta_x}$ which is the functor restricted on the poset Δ_x as a subcategory of $\bar{\mathbb{R}}^2$. We call it a *1-dimensional slice* of M along Δ_x . Define

$$\delta^* = \inf_{\delta \in \bar{\mathbb{R}}} \{ \delta : \forall x \in \bar{\mathbb{R}}^2, M|_{\Delta_x} \text{ and } N|_{\Delta_x} \text{ are } \delta\text{-interleaved} \}.$$

Proposition 22 follows from the observation that $\delta^* = \sup_{x \in \bar{\mathbb{R}}^2} \{ d_I(M|_{\Delta_x}, N|_{\Delta_x}) \}$.

Proposition 22. *For two interval modules M, N and $\delta \in \mathbb{R}^+$, we have $\delta > \delta^*$ if and only if there exist two families of linear maps $\phi = \{ \phi_x : M_x \rightarrow N_{(x+\delta)} \}$ and $\psi = \{ \psi_x : N_x \rightarrow M_{(x+\delta)} \}$ such that for each $x \in \bar{\mathbb{R}}^2$, the 1-dimensional slices $M|_{\Delta_x}$ and $N|_{\Delta_x}$ are δ -interleaved by the linear maps $\phi|_{\Delta_x}$ and $\psi|_{\Delta_x}$.*

Theorem 23. *Two interval modules M and N are δ -interleaved if and only if*

- $\delta > \delta^*$, and
- each intersection component of M and $N_{\rightarrow\delta}$ is either $(M, N_{\rightarrow\delta})$ -valid or $\delta_{(M, N_{\rightarrow\delta})}$ -trivializable, and each intersection component of $M_{\rightarrow\delta}$ and N is either $(N, M_{\rightarrow\delta})$ -valid or $\delta_{(N, M_{\rightarrow\delta})}$ -trivializable.

Proof. \implies direction: Suppose M and N are δ -interleaved. By definition, we have two families of linear maps $\{ \phi_x \}$ and $\{ \psi_x \}$ which satisfy both triangular and square commutativities. Let the morphisms between the two persistence modules constituted by these two families of linear maps be $\phi = \{ \phi_x \}$ and $\psi = \{ \psi_x \}$ respectively. By Proposition 22, we get the first part of the claim that $\delta > \delta^*$. For each intersection component Q of M and $N_{\rightarrow\delta}$ with interval $I := I_Q$, consider the restriction $\phi|_I$. By Proposition 12, $\phi|_I$ is constant, that is, $\phi|_I \equiv 0$ or $\mathbb{1}$. If $\phi|_I \equiv \mathbb{1}$, by Proposition 14, Q is $(M, N_{\rightarrow\delta})$ -valid. If $\phi|_I \equiv 0$, by the triangular commutativity of ϕ , we have that $\rho_{x \rightarrow x+2\delta}^M = \psi_{x+\delta} \circ \phi_x = 0$ for

each point $x \in I$. That means $x + 2\vec{\delta} \notin I_M$. By Fact 18(i), $\text{dl}(x, U(I_M))/2 < \delta$. Similarly, $\rho_{x-\vec{\delta} \rightarrow x+\vec{\delta}}^N = \phi_x \circ \psi_{x-\vec{\delta}} = 0 \implies x - \vec{\delta} \notin I_N$, which is the same as to say $x - 2\vec{\delta} \notin I_{N \rightarrow \delta}$. By Fact 18(i), $\text{dl}(x, L(I_{N \rightarrow \delta}))/2 < \delta$. So $\forall x \in I$, we have $d_{triv}^{(M, N \rightarrow \delta)}(x) < \delta$. This means Q is $\delta_{(M, N \rightarrow \delta)}$ -trivializable. Similar statement holds for intersection components of $M \rightarrow \delta$ and N .

\Leftarrow direction: We construct two families of linear maps $\{\phi_x\}, \{\psi_x\}$ as follows: On the interval $I := I_{Q_i}$ of each intersection component Q_i of M and $N \rightarrow \delta$, set $\phi|_I \equiv \mathbb{1}$ if Q_i is $(M, N \rightarrow \delta)$ -valid and $\phi|_I \equiv 0$ otherwise. Set $\phi_x \equiv 0$ for all x not in the interval of any intersection component. Similarly, construct $\{\psi_x\}$. Note that, by Proposition 14, $\phi := \{\phi_x\}$ is a morphism between M and $N \rightarrow \delta$, and $\psi := \{\psi_x\}$ is a morphism between N and $M \rightarrow \delta$. Hence, they satisfy the square commutativity. We show that they also satisfy the triangular commutativity. We claim that $\forall x \in I_M$, $\rho_{x \rightarrow x+2\vec{\delta}}^M = \mathbb{1} \implies x + \vec{\delta} \in I_N$ and similar statement holds for I_N . From condition that $\delta > \delta^*$ and by proposition 22, we know that there exist two families of linear maps satisfying triangular commutativity everywhere, especially on the pair of 1-dimensional persistence modules $M|_{\Delta_x}$ and $N|_{\Delta_x}$. From triangular commutativity we know that $x + \vec{\delta} \in I_N$ since otherwise one cannot construct a δ -interleaving between $M|_{\Delta_x}$ and $N|_{\Delta_x}$. Now for each $x \in I_M$ with $\rho_{x \rightarrow x+2\vec{\delta}}^M = \mathbb{1}$, we have $\text{dl}(x, U(I_M))/2 \geq \delta$ by Fact 18, and $x + \vec{\delta} \in I_N$ by our claim. This implies that $x \in I_M \cap I_{N \rightarrow \delta}$ is a point in an interval of an intersection component Q_x of $M, N \rightarrow \delta$ which is not $\delta_{(M, N \rightarrow \delta)}$ -trivializable. Hence, it is $(M, N \rightarrow \delta)$ -valid by the assumption. So, by our construction of ϕ on valid intersection components, $\phi_x = \mathbb{1}$. Symmetrically, we have that $x + \vec{\delta} \in I_N \cap I_{M \rightarrow \delta}$ is a point in an interval of an intersection component of N and $M \rightarrow \delta$ which is not $\delta_{(N, M \rightarrow \delta)}$ -trivializable since $\text{dl}(x + \vec{\delta}, L(I_M))/2 \geq \delta$. So by our construction of ψ on valid intersection components, $\psi_{x+\vec{\delta}} = \mathbb{1}$. Then, we have $\rho_{x \rightarrow x+2\vec{\delta}}^M = \psi_{x+\vec{\delta}} \circ \phi_x$ for every nonzero linear map $\rho_{x \rightarrow x+2\vec{\delta}}^M$. The statement also holds for any nonzero linear map $\rho_{x \rightarrow x+2\vec{\delta}}^N$. Therefore, the triangular commutativity holds. \square

Note that the above proof provides a construction of the interleaving maps for a specific δ if it exists. Furthermore, the interleaving distance $d_I(M, N)$ is the infimum of all δ satisfying the two conditions in the theorem, which means $d_I(M, N)$ is the infimum of all $\delta > \delta^*$ satisfying condition 2 in Theorem 23. Based on this observation, we propose a search algorithm for computing the interleaving distance $d_I(M, N)$ for interval modules M and N .

Definition 24 (Candidate set). For two interval modules M and N , and for each point x in $I_M \cup I_N$, let

$$\begin{aligned} D(x) &= \{\text{dl}(x, L(I_M)), \text{dl}(x, L(I_N)), \text{dl}(x, U(I_M)), \text{dl}(x, U(I_N))\} \text{ and} \\ S &= \{d \mid d \in D(x) \text{ or } 2d \in D(x) \text{ for some vertex } x \in V(I_M) \cup V(I_N)\} \text{ and} \\ S_{\geq \delta} &:= \{d \mid d \geq \delta, d \in S\}. \end{aligned}$$

Algorithm INTERLEAVING (output: $d_I(M, N)$, input: I_M and I_N with t vertices in total)

1. Compute the candidate set S and let ϵ be the smallest difference between any two numbers in S . /* $O(t)$ time */
2. Compute δ^* ; Let $\delta = \delta^*$. /* $O(t)$ time */
3. Output δ after a binary search in $S_{\geq \delta^*}$ by following steps /* $O(\log t)$ probes */
 - let $\delta' = \delta + \epsilon$
 - Compute intersections $I_M \cap I_{N \rightarrow \delta'}$ and $I_N \cap I_{M \rightarrow \delta'}$. /* $O(t)$ time */
 - For each intersection component, check if it is valid or trivializable according to Theorem 23. /* $O(t)$ time */

In the above algorithm, the following generic task of computing *diagonal span* is performed for several steps. Let L and U be any two chains of vertical and horizontal edges that are both x - and y -monotone. Assume that L and U have at most t vertices. Then, for a set X of $O(t)$ points in L , one can compute the intersection of Δ_x with U for every $x \in X$ in $O(t)$ total time. The idea is to first compute by a binary search a point x in X so that Δ_x intersects U if at all. Then, for other points in X , traverse from x in both directions while searching for the intersections of the diagonal line with U in lock steps.

Now we analyze the complexity of the algorithm INTERLEAVING. The candidate set, by definition, has only $2t$ values which can be computed in $O(t)$ time by the diagonal span procedure. Proposition 25 shows that δ^* is in S and can be determined by computing the one dimensional interleaving distances $d_I(M|_{\Delta_x}, N|_{\Delta_x})$ for diagonal lines passing through $O(t)$ vertices of I_M and I_N . This can be done in $O(t)$ time by diagonal span procedure. Once we determine δ^* , we search for $\delta = d_I(M, N)$ in the truncated set $S_{\geq \delta^*}$ to satisfy

the first condition of Theorem 23. Intersections between two polygons I_M and I_N bounded by x - and y -monotone chains can be computed in $O(t)$ time by a simple traversal of the boundaries. The validity and trivializability of each intersection component can be determined in time linear in the number of its vertices due to Proposition 19 and Proposition 21 respectively. Since the total number of intersection points is $O(t)$, validity check takes $O(t)$ time in total. The check for trivializability also takes $O(t)$ time if one uses the diagonal span procedure.

Proposition 25 below says that δ^* is determined by a vertex in I_M or I_N and $\delta^* \in S$. Its proof appears in Appendix A.

Proposition 25. (i) $\delta^* = \max_{x \in V(I_M) \cup V(I_N)} \{d_I(M|_{\Delta_x}, N|_{\Delta_x})\}$, (ii) $\delta^* \in S$.

The correctness of the algorithm INTERLEAVING already follows from Theorem 23 as long as the candidate set contains the distance $d_I(M, N)$. The following concept of stable intersections helps us to establish this result.

Definition 26 (Stable intersection). *Let Q be an intersection component of M and N . We say Q is stable if every intersection point $x \in I_Q \cap B(I_M) \cap B(I_N)$ is non-degenerate, that is, x is in the interior of two edges $e_1 \in E(I_M)$ and $e_2 \in E(I_N)$, and $e_1 \perp e_2$ at x .*

From Proposition 42 and Corollary 43 in Appendix A, we have the following claim.

Proposition 27. $d \notin S$ if and only if each intersection component of $M, N_{\rightarrow d}$, and $N_{\rightarrow d}, M$ is stable.

The main property of a stable intersection component Q of M and N is that if we shift one of the interval module, say N , to $N_{\rightarrow \epsilon}$ continuously for some small value $\epsilon \in \mathbb{R}^+$, the interval I_{Q^ϵ} of the intersection component Q^ϵ of M and $N_{\rightarrow \epsilon}$ changes continuously. Next proposition follows directly from the stability of intersection components.

Proposition 28. *For a stable intersection component Q of M and N , there exists a positive real $\delta \in \mathbb{R}^+$ so that the following holds:*

For each $\epsilon \in (-\delta, +\delta)$, there exists a unique intersection component Q^ϵ of M and $N_{\rightarrow \epsilon}$ so that it is still stable and $I_{Q^\epsilon} \cap I_Q \neq \emptyset$. Furthermore, there is a bijection $\mu_\epsilon : V(I_Q) \rightarrow V(I_{Q^\epsilon})$ so that $\forall x \in V(I_Q)$, x and $\mu_\epsilon(x)$ are on the same horizontal, vertical, or diagonal line, and $d_\infty(\mu_\epsilon(x), x) = \epsilon$. We call the set $\{Q^\epsilon \mid \epsilon \in (-\delta, +\delta)\}$ a stable neighborhood of Q .

Corollary 29. *For a stable intersection component Q , we have:*

(i) Q is (M, N) -valid iff each Q^ϵ in the stable neighborhood is $(M, N_{\rightarrow\epsilon})$ -valid.

(ii) If Q is $d_{(M,N)}$ -trivializable, then Q^ϵ is $(d + 2\epsilon)_{(M, N_{\rightarrow\epsilon})}$ -trivializable.

Proof. (i): Let Q^ϵ be any intersection component in a stable neighborhood of Q . We know that if Q is (M, N) -valid, then $VL(I_Q) \subseteq L(I_M)$ and $VU(I_Q) \subseteq U(I_N)$. By Proposition 28, $\mu_\epsilon(VL(I_Q)) = VL(I_{Q^\epsilon}) \subseteq L(I_M)$ and $\mu_\epsilon(UV(I_Q)) = UV(I_{Q^\epsilon}) \subseteq U(I_N)$. So Q^ϵ is $(M, N_{\rightarrow\epsilon})$ -valid. Other direction of the implication can be proved by switching the roles of Q and Q^ϵ in the above argument.

(ii): From Proposition 28, we have that $\forall x' \in V(I_{Q^\epsilon})$, there exists a point $x \in V(I_Q)$ so that x and x' are on some horizontal, vertical, or diagonal line (Δ_x) , and $d_\infty(x, x') \leq \epsilon$. Then, by Fact 18(ii), one observes

$$d_{triv}^{(M, N_{\rightarrow\epsilon})}(x) \leq d_{triv}^{(M, N_{\rightarrow\epsilon})}(x') + \epsilon \leq d_{triv}^{(M, N)}(x) + 2\epsilon < d + 2\epsilon.$$

Therefore, Q^ϵ is $(d + 2\epsilon)_{(M, N_{\rightarrow\epsilon})}$ -trivializable. \square

Theorem 30. $d_I(M, N) \in S$.

Proof. Suppose that $d = d_I(M, N) \notin S$. Let d^* be the largest value in S satisfying $d^* \leq d$. Note that $d \in S$ if and only if $d = d^*$. Then, $d^* < d$ by our assumption that $d \notin S$.

By definition of interleaving distance, we have $\forall d' > d$, there is a d' -interleaving between M and N , and $\forall d'' < d$, there is no d'' -interleaving between M and N . By Proposition 25(ii), one can see that $\delta^* \leq d^* < d$. So, to get a contradiction, we just need to show that there exists d'' , $d^* < d'' < d$, satisfying the condition 2 in Theorem 23.

Let Q be any intersection component of $M, N_{\rightarrow d}$ or $N, M_{\rightarrow d}$. Without loss of generality, assume Q is an intersection component of M and $N_{\rightarrow d}$. By Proposition 27, Q is stable. We claim that there exists some $\epsilon > 0$ such that $Q^{-\epsilon}$ is an intersection component of M and $N_{\rightarrow d-\epsilon}$ in a stable neighborhood of Q , and $Q^{-\epsilon}$ is either $(M, N_{\rightarrow d-\epsilon})$ -valid or $(d - \epsilon)_{(M, N_{\rightarrow d-\epsilon})}$ -trivializable.

Let $\epsilon > 0$ be small enough so that $Q^{+\epsilon}$ is a stable intersection component of M and $N_{\rightarrow d+\epsilon}$ in a stable neighborhood of Q . By Theorem 23, $Q^{+\epsilon}$ is either $(M, N_{\rightarrow(d+\epsilon)})$ -valid or $(d + \epsilon)_{(M, N_{\rightarrow(d+\epsilon)})}$ -trivializable. If $Q^{+\epsilon}$ is $(M, N_{\rightarrow(d+\epsilon)})$ -valid, then by Corollary 29(i), any intersection component in a stable neighborhood of Q is valid, which means there exists $Q^{-\epsilon}$ that is

$(M, N_{\rightarrow d-\epsilon})$ -valid for some $\epsilon > 0$. Now assume $Q^{+\epsilon}$ is not $(M, N_{\rightarrow(d+\epsilon)})$ -valid. Then, $\forall \epsilon > 0$, $Q^{+\epsilon}$ is $(M, N_{\rightarrow(d+\epsilon)})$ -trivializable, By Proposition 21 and 29(ii), we have $\forall x \in V(I_Q)$, $d_{triv}^{(M, N_{\rightarrow d+\epsilon})}(x) < d + 3\epsilon$, $\forall \epsilon > 0$. Taking $\epsilon \rightarrow 0$, we get $\forall x \in V(I_Q)$, $d_{triv}^{(M, N_{\rightarrow d})}(x) \leq d$. We claim that, actually, $\forall x \in V(I_Q)$, $d_{triv}^{(M, N_{\rightarrow d})}(x) < d$. If the claim were not true, some point $x \in V(I_Q)$ would exist so that $d_{triv}^{(M, N_{\rightarrow d})}(x) = d$. There are two cases. If $x \in V(I_M) \cup V(I_N)$, then obviously $d = d_{triv}^{(M, N_{\rightarrow d})}(x) \in S$ contradicting $d \neq d^*$. The other case is that x is the intersection point of two perpendicular edges $e_1 \in E(I_M)$ and $e_2 \in E(I_N)$ since Q is a stable intersection component. But, then x and $\pi_L(x)$ are always on two parallel edges where L is either $U(I_M)$ or $L(I_N)$. By Proposition 41(ii), we have $d = d^*$, reaching a contradiction. Now by our claim and Proposition 21, Q is $d_{(M, N_{\rightarrow d})}$ -trivializable where $d > d^* \geq \max_{x \in V(I_Q)} \{d_{triv}^{(M, N_{\rightarrow d})}(x)\}$. Let $\delta = d - d^*$ and $\epsilon = \delta/4$. Since $d - \epsilon = d - \delta/4 > d - \delta/2 = d - \delta + 2 \cdot \delta/4 = d^* + 2\epsilon$ and $d^* \geq \max_{x \in V(I_Q)} \{d_{triv}^{(M, N_{\rightarrow d})}(x)\}$, we have $d > d^*$ and $d - \epsilon > \max_{x \in V(I_Q)} \{d_{triv}^{(M, N_{\rightarrow d})}(x)\} + 2\epsilon$. Therefore, by Corollary 21, $Q^{-\epsilon}$ is $(d - \epsilon)_{(M, N_{\rightarrow d-\epsilon})}$ -trivializable.

The above argument shows that there exists a d'' -interleaving where $d'' = d - \epsilon < d$, reaching a contradiction. \square

4 A lower bound on d_I

In this section we propose a distance between two persistence modules that bounds the interleaving distance from below. This distance is defined for n -D modules and not necessarily only for 2-D modules. It is based on dimensions of the vectors involved with the two modules and is efficiently computable.

Let $[n] = \{1, 2, \dots, n\}$ be the set of all the integers from 1 to n . Let $\binom{[n]}{k} = \{s \subseteq [n] : |s| = k\}$ be the set of all subset in $[n]$ with cardinality k .

Definition 31. For a right continuous function $f : \mathbb{R}^n \rightarrow \mathbb{Z}$, define the differential of f to be $\Delta f : \mathbb{R}^n \rightarrow \mathbb{Z}$ where

$$\Delta f(x) = \sum_{k=0}^n (-1)^k \cdot \sum_{s \in \binom{[n]}{k}} \lim_{\epsilon \rightarrow 0^+} f(x - \epsilon \cdot \sum_{i \in s} e_i)$$

Note that for $k = 0$, $\sum_{s \in \binom{[n]}{k}} \lim_{\epsilon \rightarrow 0^+} f(x - \epsilon \cdot \sum_{i \in s} e_i) = f(x)$. We say f is nice if the support $\text{supp}(\Delta f)$ is finite and $\text{supp}(f) \subseteq \{x \mid x \geq \vec{a}\}$ for some $a \in \mathbb{R}$.

The differential Δf is a function recording the change of function values of f at each point, especially at 'jump points'. For $n = 1$, $\Delta f(x) = f(x) - \lim_{\epsilon \rightarrow 0_+} f(x - \epsilon)$. For $n = 2$, which is the case we deal with, we have

$$\Delta f(x) = f(x) - \lim_{\epsilon \rightarrow 0_+} f(x - (\epsilon, 0)) - \lim_{\epsilon \rightarrow 0_+} f(x - (0, \epsilon)) + \lim_{\epsilon \rightarrow 0_+} f(x - (\epsilon, \epsilon)).$$

See Figure 2 and 3 for illustrations in 1- and 2-D cases respectively.

Proposition 32. *For a nice function f , $f(x) = \sum_{y \leq x} \Delta f(y)$ (Proof in Appendix B).*

We also define $\Delta f_+ = \max\{\Delta f, 0\}$, $\Delta f_- = \min\{\Delta f, 0\}$ and $f_{\Sigma^+}(x) = \sum_{y \leq x} \Delta f_+(y)$, $f_{\Sigma^-}(x) = \sum_{y \leq x} \Delta f_-(y)$. Note that $f_{\Sigma^+} \geq 0$, $f_{\Sigma^-} \leq 0$, and are both monotonic functions. By definition and property of Δf , we have $f = f_{\Sigma^+} + f_{\Sigma^-}$.

Definition 33. *For any $\delta > 0$, we define the δ -extension of f as $f^{+\delta} = f_+(x + \delta) + f_-(x - \delta)$. Similarly we define the δ -shrinking of f as $f^{-\delta} = f_-(x + \delta) + f_+(x - \delta)$ (see Figure 2).*

Proposition 34 below follows from the definition.




Figure 2: A nice function and its differential (left), its δ -extension (middle), δ -shrinking (right)

Proposition 34. *For any $\delta > 0 \in \mathbb{R}$, we have $f^{\pm\delta}(x) = f(x \mp \delta) + \sum_{y \leq x \pm \delta, y \not\leq x \mp \delta} \Delta f_{\pm}(y)$.*

That is to say, for any $\delta \in \mathbb{R}$, the extended (shrunk) function f^{δ} can be computed by adding to $f(x - |\delta|)$ the positive (negative) difference values of Δf in $(x - |\delta|, x + |\delta|]$. From this, it follows:

Corollary 35. *Given $0 \leq \delta \leq \delta' \in \mathbb{R}$, we have $f^{+\delta} \leq f^{+\delta'}$ and $f^{-\delta} \geq f^{-\delta'}$.*

Definition 36. For any two nice functions $f, g : \mathbb{R}^n \rightarrow \mathbb{Z}$ and $\delta \geq 0$, we say f, g are within δ -extension, denoted as $f_{\leftarrow\delta\rightarrow}g$, if $f \leq g^{+\delta}$ and $g \leq f^{+\delta}$. Similarly, we say f, g are within δ -shrinking, denoted as $f_{\rightarrow\delta\leftarrow}g$, if $f \geq g^{-\delta}$ and $g \geq f^{-\delta}$.

Let d_+, d_-, d_0 be defined as follows on the space of all nice real-valued functions on \mathbb{R}^n :

$$d_-(f, g) = \inf_{\delta} \{ \delta \mid f_{\rightarrow\delta\leftarrow}g \}, \quad d_+(f, g) = \inf_{\delta} \{ \delta \mid f_{\leftarrow\delta\rightarrow}g \}, \quad d_0(f, g) = \min(d_-, d_+)$$

One can verify that d_0 is indeed a distance function. Also, note that when $f, g \geq 0$ (for example, f, g are dimension functions as defined below), we have $d_- \leq d_+$, hence $d_0 = d_-$. It seems that the definition of d_- has a similar connotation as the erosion distance defined by Patel [24] in 1-D case.

4.1 Dimension distance




Figure 3: Dimension function (left), its differential is non-zero only at vertices (right)

Given a persistence module M , let the *dimension function* $\text{dm}M : \mathbb{R}^n \rightarrow \mathbb{Z}$ be defined as $\text{dm}M(x) = \dim(M_x)$. The distance $d_0(\text{dm}M, \text{dm}N)$ for two modules M and N is called the dimension distance. Our main result in theorem 38 is that this distance is stable with respect to the interleaving distance and thus provides a lower bound for it.

Definition 37. A persistence module M is nice if there exists a value $\epsilon_0 \in \mathbb{R}^+$ so that for every $\epsilon < \epsilon_0$, each linear map $\rho_{x \rightarrow x+\bar{\epsilon}}^M : M_x \rightarrow M_{x+\bar{\epsilon}}$ is either injective or surjective (or both).

For example, a persistence module generated by a simplicial filtration defined on a grid with at most one additional simplex being introduced between two adjacent grid points satisfies this nice condition above.

Theorem 38. For nice persistence modules M and N , $d_0(\text{dm}M, \text{dm}N) \leq d_I(M, N)$.

Proof. Let $d_I(M, N) = \delta$. There exists δ -interleaving, $\phi = \{\phi_x\}, \psi = \{\psi_x\}$ which satisfy both triangular and square commutativity. We claim $(\text{dm}M)^{-\delta} \leq \text{dm}N$ and $(\text{dm}N)^{-\delta} \leq \text{dm}M$.

Let $x \in \mathbb{R}^n$ be any point. By Proposition 34, we know that $(\text{dm}M)^{-\delta}(x) = \text{dm}M(x-\delta) + \sum_{y \leq x+\delta, y \not\leq x-\delta} (\Delta \text{dm}N_-)(y)$. If $\text{dm}M(x-\delta) \leq \text{dm}N(x)$, then we get $(\text{dm}M)^{-\delta}(x) \leq \text{dm}M(x-\delta) \leq \text{dm}N(x)$, because $\sum_{y \leq x+\delta, y \not\leq x-\delta} (\Delta \text{dm}N_-)(y) \leq 0$.

Now assume $\text{dm}M(x-\delta) > \text{dm}N(x)$. From triangular commutativity, we have $\text{rank}(\psi_x \circ \phi_{x-\delta}) = \text{rank}(\rho_{x-\delta \rightarrow x+\delta}^M)$, which gives $\dim(\text{im}(\rho_{x-\delta \rightarrow x+\delta}^M)) \leq \dim(\text{im}(\phi_{x-\delta})) \leq \text{dm}N(x)$.

There exists a collection of linear maps $\{\rho_i : M_{x_i} \rightarrow M_{x_{i+1}}\}_{i=0}^k$ such that $\rho_{x-\delta \rightarrow x+\delta}^M = \rho_k \circ \rho_{k-1} \circ \dots \circ \rho_1 \circ \rho_0$ and each ρ_i is either injective or surjective. Let $\text{im}_i = \text{im}(\rho_i \circ \dots \circ \rho_0)$. Note that $\text{im}_k = \text{im}(\rho_{x-\delta \rightarrow x+\delta}^M)$. Let $\epsilon_i = \dim(\text{im}_i) - \dim(\text{im}_{i-1})$. Then note that $\epsilon_i = 0$ if ρ_i is injective and $\dim(\text{im}_k) - \dim(M_{x_0}) = \sum_{i=1}^k \epsilon_i$. Since $\dim(\text{im}_k) - \dim(M_{x_0}) < 0$, there exists a collection of ρ_{i_j} 's such that $\epsilon_{i_j} < 0$. This means these ρ_{i_j} 's are non-isomorphic surjective linear maps with $\dim(M_{x_{i_j}}) - \dim(M_{x_{i_j-1}}) < 0$. By definition of Δdm , this means that, for each pair (x_{i_j-1}, x_{i_j}) , there exists a collection y_1, y_2, \dots such that $y_l \leq x_{i_j}, y_l \not\leq x_{i_j-1}$ and $\sum_l (\Delta \text{dm}M_-)(y_l) \leq \epsilon_{i_j}$. All these y 's also satisfy that $y \leq x + \delta, y \not\leq x - \delta$. So,

$$\sum_{y \leq x+\delta, y \not\leq x-\delta} (\Delta \text{dm}M_-)(y) \leq \sum_j \epsilon_j = \dim(\text{im}_k) - \dim(M_{x_0}) \leq \dim(N_x) - \dim(M_{x-\delta}),$$

which gives $(\text{dm}M)^{-\delta}(x) \leq \text{dm}N(x)$. Similarly, we can show $(\text{dm}N)^{-\delta}(x) \leq \text{dm}M(x)$. \square

4.2 Computation

For computational purpose, assume that two input persistence modules M and N are finite in that they are functors on the subcategory $\{1, \dots, k\}^n \subset \mathbb{R}^n$ and the dimension functions $f := \text{dm}M, g := \text{dm}N$ have been given as input on an n -dimensional k -ary grid.

First, for the dimension functions f, g , we compute $\Delta f, \Delta g, \Delta f_{\pm}, \Delta g_{\pm}, f_{\pm}, g_{\pm}$ in $O(k^2)$ time. By Proposition 34, for any $\delta \in \mathbb{Z}^+$, we can also compute

$f^{\pm\delta}$, $g^{\pm\delta}$ in $O(k^2)$ time. Then we can apply the binary search to find the minimal value δ within a bounded region such that f, g are within δ -extension or δ -shrinking. This takes $O(\log k)$ time. So the entire computation takes $O(k^2 \log k)$ time.

5 Conclusions

In this paper, we presented an efficient algorithm to compute the bottleneck distance of two 2-D persistence modules given by indecomposables that may have non-constant complexity. No such algorithm for such case is known. Making the algorithm more efficient will be one of our future goals. Extending the algorithm or its modification to larger classes of modules such as the n -D modules or exact pfd bi-modules considered in [14] will be interesting. The definition of valid and trivializable intersection component and Theorem 21 can be extended easily for n -D modules. So is the algorithm. But, further work is necessary to establish the correctness of the algorithm for this general case.

The assumption of nice modules for dimension distance d_0 is needed so that the dimension function, which is a weaker invariant compared to the rank invariants or barcodes in one dimensional case, provides meaningful information without ambiguity. There are cases where the dimension distance can be larger than interleaving distance if the assumption of nice modules is dropped. Of course, one can adjust the definition of dimension distance to incorporate more information so that it remains bounded from above by the interleaving distance.

Acknowledgments

This research is supported by NSF grants CCF-1526513, 1740761, and 1733798. We thank anonymous referees for insightful comments on a previous version of this paper.

References

- [1] Michael Atiyah. On the Krull-Schmidt theorem with application to sheaves. *Bulletin de la Société Mathématique de France*, 84:307–317,

1956. URL: <http://eudml.org/doc/86907>.

- [2] Ulrich Bauer and Michael Lesnick. Induced matchings of barcodes and the algebraic stability of persistence. In *Proceedings of the Thirtieth Annual Symposium on Computational Geometry*, SoCG'14, pages 355:355–355:364, 2014.
- [3] Silvia Biasotti, Andrea Cerri, Patrizio Frosini, and Daniela Giorgi. A new algorithm for computing the 2-dimensional matching distance between size functions. *Pattern Recognition Letters*, 32(14):1735–1746, 2011.
- [4] Håvard Bjerkevik. Stability of higher-dimensional interval decomposable persistence modules. *arXiv preprint arXiv:1609.02086*, 2016.
- [5] Håvard Bjerkevik and Magnus Botnan. Computational complexity of the interleaving distance. *arXiv preprint arXiv:1712.04281*, 2017.
- [6] Magnus Botnan, Justin Curry, and Elizabeth Munch. The poset interleaving distance, 2016. URL: https://jointmathematicsmeetings.org/amsmtgs/2180_abstracts/1125-55-1151.pdf.
- [7] Magnus Botnan and Michael Lesnick. Algebraic stability of zigzag persistence modules. *arXiv preprint arXiv:1604.00655*, 2016.
- [8] Peter Bubenik and Jonathan Scott. Categorification of persistent homology. *Discrete & Computational Geometry*, 51(3):600–627, 2014.
- [9] Gunnar Carlsson and Afra Zomorodian. The theory of multidimensional persistence. *Discrete & Computational Geometry*, 42(1):71–93, Jul 2009.
- [10] Andrea Cerri, Barbara Di Fabio, Massimo Ferri, Patrizio Frosini, and Claudia Landi. Betti numbers in multidimensional persistent homology are stable functions. *Mathematical Methods in the Applied Sciences*, 36(12):1543–1557, 2013.
- [11] Andrea Cerri and Patrizio Frosini. A new approximation algorithm for the matching distance in multidimensional persistence. Technical report, February 2011. URL: <http://amsacta.unibo.it/2971>.
- [12] Frédéric Chazal, David Cohen-Steiner, Marc Glisse, Leonidas Guibas, and Steve Oudot. Proximity of persistence modules and their diagrams.

- In *Proceedings of the Twenty-fifth Annual Symposium on Computational Geometry*, SoCG '09, pages 237–246, 2009.
- [13] Frédéric Chazal, Vin de Silva, Marc Glisse, and Steve Oudot. The structure and stability of persistence modules. *arXiv preprint arXiv:1207.3674*, 2012.
 - [14] Jérémy Cochoy and Steve Oudot. Decomposition of exact pfd persistence bimodules. *arXiv preprint arXiv:1605.09726*, 2016.
 - [15] David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. *Discrete & Computational Geometry*, 37(1):103–120, 2007.
 - [16] William Crawley-Boevey. Decomposition of pointwise finite-dimensional persistence modules. *Journal of Algebra and Its Applications*, 14(05):1550066, 2015. doi:10.1142/S0219498815500668.
 - [17] Vin de Silva, Elizabeth Munch, and Amit Patel. Categorified reeb graphs. *Discrete & Computational Geometry*, 55(4):854–906, Jun 2016.
 - [18] Herbert Edelsbrunner and John Harer. *Computational Topology: An Introduction*. Applied Mathematics. American Mathematical Society, 2010.
 - [19] Michael Kerber, Dmitriy Morozov, and Arnur Nigmatov. Geometry helps to compare persistence diagrams. *Journal of Experimental Algorithmics (JEA)*, 22(1):1–4, 2017.
 - [20] Claudia Landi. The rank invariant stability via interleavings. *arXiv preprint arXiv:1412.3374*, 2014.
 - [21] Michael Lesnick. The theory of the interleaving distance on multidimensional persistence modules. *Foundations of Computational Mathematics*, 15(3):613–650, 2015.
 - [22] Klaus Lux and Magdolna Szöke. Computing decompositions of modules over finite-dimensional algebras. *Experimental Mathematics*, 16(1):1–6, 2007.
 - [23] Steve Oudot. *Persistence theory: from quiver representations to data analysis*, volume 209. American Mathematical Society, 2015.

- [24] Amit Patel. Generalized persistence diagrams. *arXiv preprint arXiv:1601.03107*, 2016.
- [25] Carry Webb. Decomposition of graded modules. *Proc. American Math. Soc.*, 94(4):565–571, 1985.

Appendix

A Missing details in section 3

Triangular and square commutative diagrams.

$$\begin{array}{ccc}
 M_x & \xrightarrow{\rho_{x \rightarrow x+2\bar{\delta}}^M} & M_{x+2\bar{\delta}} \\
 \phi_x \searrow & & \nearrow \psi_{x+\bar{\delta}} \\
 & & N_{x+\bar{\delta}}
 \end{array}
 \qquad
 \begin{array}{ccc}
 N_x & \xrightarrow{\rho_{x \rightarrow x+2\bar{\delta}}^N} & N_{x+2\bar{\delta}} \\
 \psi_x \searrow & & \nearrow \phi_{x+\bar{\delta}} \\
 & & M_{x+\bar{\delta}}
 \end{array}$$

$$\begin{array}{ccc}
 M_x & \xrightarrow{\rho_{x \rightarrow y}^M} & M_y \\
 \phi_x \downarrow & & \downarrow \phi_y \\
 N_{x+\bar{\delta}} & \xrightarrow{\rho_{x+\bar{\delta} \rightarrow y+\bar{\delta}}^N} & N_{y+\bar{\delta}}
 \end{array}
 \qquad
 \begin{array}{ccc}
 M_{x+\bar{\delta}} & \xrightarrow{\rho_{x+\bar{\delta} \rightarrow y+\bar{\delta}}^M} & M_{y+\bar{\delta}} \\
 \psi_x \uparrow & & \uparrow \psi_y \\
 N_x & \xrightarrow{\rho_{x \rightarrow y}^N} & N_y
 \end{array}$$

Proposition 14 and its proof.

Let $\{Q_i\}$ be a set of intersection components of M and N with intervals $\{I_{Q_i}\}$. Let $\{\phi_x\} : M \rightarrow N$ be the family of linear maps defined as $\phi_x = \mathbb{1}$ for all $x \in I_{Q_i}$ and $\phi_x = 0$ otherwise. Then ϕ is a morphism if and only if every Q_i is (M, N) -valid.

Proof. \implies direction: Let $x \in I_{Q_i}$ and $y, z \in \bar{\mathbb{R}}^2$ be such that $y \leq x \leq z$. Then,

$$\begin{aligned}
 y \in I_M & \implies \rho_{y \rightarrow x}^M = \mathbb{1} \\
 & \implies \phi_x \circ \rho_{y \rightarrow x}^M = \mathbb{1} = \rho_{y \rightarrow x}^N \circ \phi_y \text{ because } \phi \text{ is a morphism} \\
 & \implies \phi_y = \mathbb{1} \\
 & \implies N_y = \mathbb{k} \\
 & \implies y \in I_N.
 \end{aligned}$$

Similarly, we have $z \in I_N \implies z \in I_M$. So, we get Q_i is (M, N) -valid.

\Leftarrow direction: We want to show that the square commutativity $\phi_y \circ \rho_{x \rightarrow y}^M = \rho_{x \rightarrow y}^N \circ \phi_x$ holds for any $x \leq y \in \bar{\mathbb{R}}^n$ as depicted in the diagram below:

$$\begin{array}{ccc} M_x & \xrightarrow{\rho_{x \rightarrow y}^M} & M_y \\ \phi_x \downarrow & & \downarrow \phi_y \\ N_x & \xrightarrow{\rho_{x \rightarrow y}^N} & N_y \end{array}$$

First, assume that M and N have a single intersection component Q with $I := I_Q$. There are several cases.

Case 1: $x, y \in I$: By assumption, every linear map in the square commutative diagram is the identity map. So, it commutes with ρ as required.

Case 2: $x, y \notin I$: By assumption we have $\phi_x = 0, \phi_y = 0$. So, it commutes with ρ trivially.

Case 3: $x \in I$: If $y \in I_N$, then by the assumption that Q is (M, N) -valid, we have $y \in I_M$. It reduces to case 1. If $y \in I_M \setminus I_N$, we have $\phi_x = \mathbb{1}, \phi_y = 0, \rho_{x \rightarrow y}^M = \mathbb{1}, \rho_{x \rightarrow y}^N = 0$, which imply $\phi_y \circ \rho_{x \rightarrow y}^M = 0 = \rho_{x \rightarrow y}^N \circ \phi_x$ as required.

Case 4: $y \in I$: If $x \in I_M$, then by assumption that Q is (M, N) -valid, we have $x \in I_N$. It reduces to case 1. If $x \in I_N \setminus I_M$, we have $\phi_x = 0, \phi_y = \mathbb{1}, \rho_{x \rightarrow y}^M = 0, \rho_{x \rightarrow y}^N = \mathbb{1}$, which imply $\phi_y \circ \rho_{x \rightarrow y}^M = 0 = \rho_{x \rightarrow y}^N \circ \phi_x$ as required.

Now for the case when M and N intersect in a set $\{Q_i\}$ that has more than one element, let ϕ_i be the morphism constructed for Q_i only. Then we let $\phi = \{\phi_x\}$ where $\phi_x = \sum_i (\phi_i)_x$. Since each $(\phi_i)_x$ is a scalar function, either $\mathbb{1}$ or 0 in $\mathbb{k} = \mathbb{Z}/2$, the sum of such morphisms is still a morphism. We can also see that $\phi_x = \mathbb{1}$ for any x in any I_{Q_i} in the set $\{Q_i\}$ and $\phi_x = 0$ if x is not in any I_{Q_i} . Hence, ϕ is a morphism as required. \square

Proposition 15 and its proof.

$$d_I(M, N) = d_I(\overline{M}, \overline{N}).$$

Proof. With the triangular inequality of the interleaving distance, the proposition follows straightforwardly from the claim that $d_I(M, \overline{M}) = 0$ which we prove below.

By definition of \overline{M} , we have $I_{\overline{M}} = \overline{I_M}$. First, note that each pair of one dimensional slices $M|_{\Delta_x}$ and $\overline{M}|_{\Delta_x}$ are δ -interleaved for any $\delta > 0$. That means $\delta^* = 0$. Let $\delta > 0$ be a small enough number and $I = I_M \cap I_{\overline{M} \rightarrow \delta}$, $J = I_{M \rightarrow \delta} \cap I_{\overline{M}}$.

We claim that $\forall x \in I, \forall y < x, y \in I_M \implies y \in I_{\overline{M} \rightarrow \delta}$. This is because $\exists w$ such that $y - \vec{\delta} < w < y$ and $w \in I_{\overline{M} \rightarrow \delta}$. By the property of interval,

$$w < y < x \text{ and } w, x \in I_{\overline{M} \rightarrow \delta} \implies y \in I_{\overline{M} \rightarrow \delta}.$$

Similarly, we have $\forall x \in I, \forall z > x, z \in I_{\overline{M} \rightarrow \delta} \implies z \in I_M$. Now we construct $\phi = \{\phi_x : M_x \rightarrow \overline{M}_{x+\delta}\}$ by setting $\forall x \in I, \phi_x \equiv \mathbb{1}$ and $\forall x \notin I, \phi_x \equiv 0$. We define $\psi = \{\psi_x : \overline{M} \rightarrow M_{x+\delta}\}$ in a similar way. Applying similar argument as in the proof of Proposition 14, one can obtain that these two maps satisfy square commutativity, and hence are morphisms.

Now we claim that ϕ and ψ provide a δ -interleaving for each pair of one-dimensional slices $M|_{\Delta_x}$ and $\overline{M}|_{\Delta_x}$, which means they also follow the triangular commutativity. Observe that $\forall x \in I_M, \forall \epsilon > 0, x + 2\vec{\epsilon} \in I_M \implies x + \vec{\epsilon} \in I_{\overline{M}}$. Symmetrically, we have $\forall x \in I_{\overline{M}}, \forall \epsilon > 0, x + 2\vec{\epsilon} \in I_{\overline{M}} \implies x + \vec{\epsilon} \in I_M$. Now let $\epsilon = \delta$ and consider any nonzero linear map $\rho_{x \rightarrow x+2\vec{\delta}}^M = \mathbb{1}$ in M . Since $x, x + 2\vec{\delta} \in I_M \implies x + \vec{\delta} \in I_{\overline{M}}$, we have $x \in I$ and $x + \vec{\delta} \in J$, which imply $\phi_x = \psi_{x+\delta} = \mathbb{1}$ by our construction of ϕ and ψ . So, $\forall x$ so that $\rho_{x \rightarrow x+2\vec{\delta}}^M = \mathbb{1}$, we have $\rho_{x \rightarrow x+2\vec{\delta}}^M = \mathbb{1} = \psi_{x+\delta} \circ \phi_x$. For those x so that $\rho_{x \rightarrow x+2\vec{\delta}}^M = 0$, observe that the commutativity holds trivially. Therefore, $\forall x, \rho_{x \rightarrow x+2\vec{\delta}}^M = \psi_{x+\delta} \circ \phi_x$. Symmetrically, we also have the commutativity $\rho_{x \rightarrow x+2\vec{\delta}}^{\overline{M}} = \phi_{x+\delta} \circ \psi_x$.

Therefore, the morphisms ϕ and ψ provide δ -interleaving on the interval modules M, \overline{M} . Since this is true for any $\delta > 0$, we get $d_I(M, \overline{M}) = 0$. \square

Proposition 19 and its proof.

For an intersection component Q of M and N with interval $I := I_Q$, the following conditions are equivalent:

- (1) Q is (M, N) -valid.
- (2) $L(I) \subseteq L(I_M)$ and $U(I) \subseteq U(I_N)$.
- (3) $VL(I) \subseteq L(I_M)$ and $VU(I) \subseteq U(I_N)$.

Proof. (1) \iff (2): Assume (1) is true. Let $x \in L(I)$. For any $y = (y_1, y_2)$ with $y_1 < x_1$ and $y_2 < x_2$, we have $y \notin I_M$ or $y \notin I_N$ because no such

point y can belong to the intersection I as x is on the boundary $L(I)$. Also, by definition of (M, N) -validity, $y \notin I_N \implies y \notin I_M$. These two conditions on y imply that $y \notin I_M$. Therefore, $x \in L(I_M)$, that is $L(I) \subseteq L(I_M)$. Similarly, we get $U(I) \subseteq U(I_N)$ proving (1) \implies (2).

Assume (2). Let $x \in I$. For any $y \leq x$, we want to show that $y \in I_M \implies y \in I_N$, which is equivalent to the condition $y \notin I \implies y \notin I_M$ since $I = I_N \cap I_M$. Observe that $y \notin I \implies y < y' = \pi_{L(I)}(y)$. By assumption that $L(I) \subseteq L(I_M)$, we have $y' \in L(I_M)$, which implies $y < \pi_{L(I_M)}(y) = y'$. So we get $y \notin I_M$. In a similar way, we can get $\forall z \geq x$, $z \notin I \implies z \notin I_N$, or equivalently, $z \in I_N \implies z \in I_M$. Therefore, by definition of (M, N) -validity, we obtain (1).

(2) \iff (3): $L(I)$ and $U(I)$ are uniquely determined by their vertices. \square

Proposition 21 and its proof.

An intersection component Q is $\delta_{(M,N)}$ -trivializable if and only if each vertex in $V(I_Q)$ is $\delta_{(M,N)}$ -trivializable.

Proof. The \implies direction is trivial. For the \impliedby direction, observe that an intersection component Q is $\delta_{(M,N)}$ -trivializable if and only if every point in $B(I_Q)$ is $\delta_{(M,N)}$ -trivializable. Now assume Q is not $\delta_{(M,N)}$ -trivializable. That means there exists a point $x \in B(I_Q)$ such that $d_{triv}^{(M,N)}(x) \geq \delta$. If x is in $V(I_Q)$, we are done. So, assume that $x \notin V(I_Q)$. Let \overline{st} , $s \leq t$, be the edge in $E(I_Q)$ containing x . Let $L \in \{U(I_M), L(I_N)\}$ be the one on which $d_{triv}^{(M,N)}(x) = \max\{\text{dl}(x, U(I_M))/2, \text{dl}(x, L(I_N))/2\}$ attains the maximum. Without loss of generality, let $L = U(I_M)$. By assumption, we have $x + 2\vec{\delta} \in I_M$. Consider $s \leq x$, we have $s \leq s + 2\vec{\delta} \leq x + 2\vec{\delta}$. Since $s, x + 2\vec{\delta} \in I_M$, by the definition of interval modules, we have $s + 2\vec{\delta} \in I_M$, which implies $d_{triv}^{(M,N)}(s) \geq \delta$ giving the necessary claim. \square

Next proposition is used to prove Proposition 25.

Proposition 39. *Let M and N be two one-dimensional interval modules with intervals $I_M = \overline{st}$ and $I_N = \overline{uv}$ respectively. We have $\delta \geq d_I(M, N)$ if and only if*

$$\begin{aligned} |s - t|_\infty > 2\delta &\implies s + \vec{\delta} \in I_N \text{ and } t - \vec{\delta} \in I_N, \text{ and} \\ |u - v|_\infty > 2\delta &\implies u + \vec{\delta} \in I_M \text{ and } v - \vec{\delta} \in I_M. \end{aligned}$$

Proof. The \implies direction is obvious by the definition of δ -interleaving. For the \impliedby direction, we split the premise into two cases.

Case(1): both $|s - t|_\infty \leq 2\delta$ and $|u - v|_\infty \leq 2\delta$ so that the premise holds vacuously. In this case M, N are two bars with length less than or equal to 2δ and one can observe that $d_I(M, N) \leq \delta$.

Case(2): there is at least one of $|s - t|_\infty$ and $|u - v|_\infty$ which is greater than 2δ . We want to show that M and N are δ -interleaved by constructing the linear maps $\phi = \{\phi_x : M_x \rightarrow N_{x+\vec{\delta}}\}$ and $\psi = \{\psi_x : N_x \rightarrow M_{x+\vec{\delta}}\}$ explicitly that satisfy both the square commutativity and triangle commutativity.

Let ϕ and ψ be defined as follows:

$$\phi_x = \begin{cases} \mathbb{1}, & x \in I_M \cap I_{N \rightarrow \delta} \\ 0, & \text{otherwise} \end{cases} \quad \psi_x = \begin{cases} \mathbb{1}, & x \in I_N \cap I_{M \rightarrow \delta} \\ 0, & \text{otherwise} \end{cases}$$

By assumption, one can easily verify that for each nonzero linear map $\rho_{x \rightarrow x+2\vec{\delta}}^M$, we have $\rho_{x \rightarrow x+2\vec{\delta}}^M = \mathbb{1} = \psi_{x+\vec{\delta}} \circ \phi_x$. Similarly, we have $\rho_{x \rightarrow x+2\vec{\delta}}^N = \mathbb{1} = \phi_{x+\vec{\delta}} \circ \psi_x$. So, ϕ and ψ satisfy the triangular commutativity. Now we show that they also satisfy the square commutativity. By Proposition 14, it is equivalent to showing that $I_M \cap I_{N \rightarrow \delta}$ is $(M, N_{\rightarrow \delta})$ -valid and $I_N \cap I_{M \rightarrow \delta}$ is $(N, M_{\rightarrow \delta})$ -valid. We show the first validity, that is, $I_M \cap I_{N \rightarrow \delta}$ is $(M, N_{\rightarrow \delta})$ -valid. The second validity can be proved in a similar way.

Observe that, for one dimensional interval modules, $I_M \cap I_{N \rightarrow \delta}$ being $(M, N_{\rightarrow \delta})$ -valid is equivalent to saying that $u - \vec{\delta} \leq s$ and $v - \vec{\delta} \leq t$. By assumption of case 2, we know that at least one of $|s - t|_\infty$ and $|u - v|_\infty$ is greater than 2δ . Consider the case when $|s - t|_\infty > 2\delta$. The other case can be argued similarly. By assumption, we have $s + \vec{\delta} \in I_N$. This means $u \leq s + \vec{\delta}$, or equivalently, $u - \vec{\delta} \leq s$. Then, the only thing remaining to be shown is that $v - \vec{\delta} \leq t$. Assume on the contrary that $v - \vec{\delta} > t$, which is equivalent to saying $v > t + \vec{\delta}$. Again, by assumption, $t - \vec{\delta} \in I_N$. This means $u \leq t - \vec{\delta}$, which implies $|v - u|_\infty > |t + \vec{\delta} - (t - \vec{\delta})|_\infty = 2\vec{\delta}$. Now by assumption, we have $v - \vec{\delta} \in I_M$, which is contradictory to $v - \vec{\delta} > t$. \square

Note that the above proof also works for interval modules with unbounded intervals.

Proposition 25 and its proof.

- (i) $\delta^* = \max_{x \in V(I_M) \cup V(I_N)} \{d_I(M|_{\Delta_x}, N|_{\Delta_x})\}$, (ii) $\delta^* \in S$.

Proof. First, we show (i). By definition of δ^* , the claim is equivalent to showing that

$$\delta \geq \delta^* \iff \delta \geq \max_{x \in V(I_M) \cup V(I_N)} \{d_I(M|_{\Delta_x}, N|_{\Delta_x})\}.$$

Let

$$\begin{aligned} \pi(V(I_M)) &= \{y \mid y \in \Delta_x \cap B(I_M), x \in V(I_M)\} \text{ and} \\ \pi(V(I_N)) &= \{y \mid y \in \Delta_x \cap B(I_N), x \in V(I_N)\}. \end{aligned}$$

We observe the following chain of equivalences.

$$\begin{aligned} \delta \geq \delta^* &\iff \text{for every pair } M|_{\Delta_x}, N|_{\Delta_x}, \delta \geq d_I(M|_{\Delta_x}, N|_{\Delta_x}) \\ &\iff \begin{cases} \forall x \in B(I_M), \\ \text{dl}(x, U(I_M)) > 2\delta \implies x + \vec{\delta} \in I_N \text{ and } \pi_{U(I_M)}(x) - \vec{\delta} \in I_N, \\ \text{dl}(x, L(I_M)) > 2\delta \implies x - \vec{\delta} \in I_N \text{ and } \pi_{L(I_M)}(x) + \vec{\delta} \in I_N. \\ \forall y \in B(I_N), \\ \text{dl}(y, U(I_N)) > 2\delta \implies y + \vec{\delta} \in I_M \text{ and } \pi_{U(I_N)}(y) - \vec{\delta} \in I_M, \\ \text{dl}(y, L(I_N)) > 2\delta \implies y - \vec{\delta} \in I_M \text{ and } \pi_{L(I_N)}(y) + \vec{\delta} \in I_M. \end{cases} \\ &\iff \begin{cases} \forall x \in \pi(V(I_M)), \\ \text{dl}(x, U(I_M)) > 2\delta \implies x + \vec{\delta} \in I_N \text{ and } \pi_{U(I_M)}(x) - \vec{\delta} \in I_N, \\ \text{dl}(x, L(I_M)) > 2\delta \implies x - \vec{\delta} \in I_N \text{ and } \pi_{L(I_M)}(x) + \vec{\delta} \in I_N. \\ \forall y \in \pi(V(I_N)), \\ \text{dl}(y, U(I_N)) > 2\delta \implies y + \vec{\delta} \in I_M \text{ and } \pi_{U(I_N)}(y) - \vec{\delta} \in I_M, \\ \text{dl}(y, L(I_N)) > 2\delta \implies y - \vec{\delta} \in I_M \text{ and } \pi_{L(I_N)}(y) + \vec{\delta} \in I_M. \end{cases} \\ &\iff \delta \geq \max_{x \in V(I_M) \cup V(I_N)} \{d_I(M|_{\Delta_x}, N|_{\Delta_x})\}. \end{aligned}$$

The first two and the last equivalences are clear by the definition of interleaving distances and Proposition 39. The \implies direction of the third equivalence is trivial since $\pi(V(I_M)) \subseteq B(I_M)$ and $\pi(V(I_N)) \subseteq B(I_N)$. For the \impliedby direction, we show that if the implications $x \pm 2\vec{\delta} \in I_M \implies x \pm \vec{\delta} \in I_N$ hold for every point $x \in \pi(V(I_M))$ then they also hold for every point in $B(I_M)$. Similarly, one can show if the implications $x \pm 2\vec{\delta} \in I_N \implies x \pm \vec{\delta} \in I_M$ hold for every point $x \in \pi(V(I_N))$, then they also hold for every point in $B(I_N)$.

Without loss of generality, assume that $x \in L(I_M)$ with $\text{dl}(x, U(I_M)) > 2\delta$. We want to show that $x + \vec{\delta} \in I_N$.

Let \overline{st} , $s \leq t$, be the edge containing x in $L(I_M)$.

If x is a vertex, then we get $x + \vec{\delta} \in I_N$ directly from the assumption. Now assume x is not a vertex, we have $s, t \in V(I_M)$ such that $s < x < t$. Note that $\text{dl}(x, U(I_M)) > 2\delta$ implies that $\pi_{U(I_M)}(x) = x + 2\vec{\delta} + \vec{\epsilon}$ for some $\epsilon > 0$, and $x + 2\vec{\delta}, x + 2\vec{\delta} + \vec{\epsilon} \in I_M$. Since $s \leq s + 2\vec{\delta} + \vec{\epsilon} \leq x + 2\vec{\delta} + \vec{\epsilon}$ and $s, x + 2\vec{\delta} + \vec{\epsilon} \in I_M$ by assumption, we have $s + 2\vec{\delta} + \vec{\epsilon} \in I_M$ by the property of interval modules, which implies that $\text{dl}(s, U(I_M)) > 2\delta$. Then, by the assumption on vertices, $\text{dl}(s, U(I_M)) > 2\delta \implies s + \vec{\delta} \in I_N$. Note that $s + \vec{\delta} \leq x + \vec{\delta}$. So, to show that $x + \vec{\delta} \in I_N$, we just need to find a point in I_N which is greater than or equal to $x + \vec{\delta}$. Then, by the property of intervals, we have that $x + \vec{\delta} \in I_N$.




Figure 4: The yellow region and black points (circles and the cross) are contained in I_M . The goal is to show that the point indicated with the cross (cross-point) is also in I_N . Since the red points are in I_N , the entire green square is contained in I_N , which implies that the cross-point $x + \vec{\delta}$ is in I_N (here s' is chosen to be z).

Let $x' = \pi_{U(I_M)}(x)$, which is greater than $x + 2\vec{\delta}$. There are two cases:

Case 1: $x' \in V(I_M)$. By the assumption, we have $x + \vec{\delta} \in I_N$.

Case 2: $x' \in B(I_M) \setminus V(I_M)$. Let \overline{uv} , $u < v$, be the edge containing x' in $U(I_M)$. Let $s' = \pi_{\overline{uv}}(s)$, $t' = \pi_{\overline{uv}}(t)$ (not necessarily exist). Let z be the minimum element in the set $\{u, v, s', t'\}$ such that $z > x'$. Then either z is a vertex or z is a projection of a vertex. That is $z \in \pi(V(I_M))$. In either case, we have $\text{dl}(z, L(I_M)) > 2\delta$ and $I_M \ni z > x'$. So, we have $z - \vec{\delta} > x' - \vec{\delta} \geq x + \vec{\delta}$ and $z - \vec{\delta} \in I_N$. Therefore, from $I_N \ni s + \vec{\delta} \leq x + \vec{\delta} < z - \vec{\delta} \in I_N$, we get $x + \vec{\delta} \in I_N$ (See Figure 4 for an example).

This completes the proof of (i). Now we argue for (ii) $\delta^* \in S$.

From the definition and isometry theorem [13] of one dimensional persistence modules, we have the following fact.

Fact 40. *Let M and N be two one-dimensional interval modules with intervals $I_M = \overline{st}$, $I_N = \overline{uv}$ respectively. We have $d_I(M, N) = \min\{\max\{|s - u|_\infty, |t - v|_\infty\}, \max\{|s - t|_\infty, |u - v|_\infty\}\}$.*

The claim (ii) follows from the above fact and claim (i). \square

For the proposition below, recall that

$$\begin{aligned} D(x) &= \{\text{dl}(x, L(I_M)), \text{dl}(x, L(I_N)), \text{dl}(x, U(I_M)), \text{dl}(x, U(I_N))\} \\ S &= \{d : d \in D(x) \text{ or } 2d \in D(x) \text{ for some vertex } x \in V(I_M) \cup V(I_N)\}. \end{aligned}$$

Proposition 41. *Let M and N be two interval modules. Given any point $x \in B(I_M)$ and any $L \in \{L(I_M), U(I_M), L(I_N), U(I_N)\}$, let $d = \text{dl}(x, L)$.*

(i) *There exist (not necessarily distinct) two values $d_1 \in D(y)$, $d_2 \in D(z)$ for (not necessarily distinct) two vertices $y, z \in V(I_M) \cup V(I_N)$ such that $d_1 \leq d \leq d_2$.*

(ii) *Furthermore, if x and $\pi_L(x)$ are on two parallel edges, then $d_1 = d_2$. In that case, $d = d_1 = d_2 = d^* \in S$.*

Proof. (i) Given $x \in B(I_M)$, let $x' = \pi_L(x)$ if exists. If either x or x' is a vertex, then we just let $y = z = x$ or x' respectively, which provides the conclusion. Now assume neither x nor x' is a vertex.

If $x \in B(\overline{\mathbb{R}^2})$, without loss of generality, let $x = (a, +\infty)$ and \overline{st} , $s \leq t$, be the edge in $E(I_M)$ containing x , then we have the following cases.

Case 1: Assume $x' = \pi_L(x)$ does not exist. That means there is no point with the second coordinate being equal to $+\infty$ in L . Then we have $\text{dl}(s, L) = +\infty = \text{dl}(x, L)$.

Assume x' exists. Let $\overline{s't'}$, $s' \leq t'$, be the edge in L containing x' .

Case 2: If $d = d_\infty(x, x') < +\infty$, then $x' = (a', +\infty)$ for some $a' \in \mathbb{R}$. If $a' = a$, then $d = 0 \in S$. If $a' \neq a$, then $x' = s'$ or t' , that is, x' is a vertex, which has been considered before.

Case 3: If $d = d_\infty(x, x') = +\infty$, then $x' = (\pm\infty, +\infty)$. But, in that case, either s or t has the first coordinate different from x' , which means either $d_\infty(s, x') = +\infty = d$ or $d_\infty(t, x') = +\infty = d$.

Now assume $x \in \mathbb{R}^2$. Let $\overline{st} \in E(I_M)$ be the edge containing x and $\overline{uv} \in E(L)$ be the edge containing x' . Also, let $l_0 = \overline{xx'}$ be the line segment

with ends x, x' . By construction, l_0 is contained in the line Δ_x passing through x that has slope 1. For any line segment l in \mathbb{R}^2 , let $|l|_\infty$ be the d_∞ distance between the two end points of l . By definition, we know that $x' = \pi_L(x) = \Delta_x \cap L$. So $dl(x, L) = d_\infty(x, x') = |l_0|$.

Consider the five lines $\Delta_x, \Delta_s, \Delta_t, \Delta_u, \Delta_v$ with slope 1. We can order these five lines by their intercepts on the axis of the first coordinate. Note that Δ_x is ordered third (in the middle) in this sequence. We pick the second and fourth ones in this sequence and observe that they necessarily intersect both edges \overline{uv} and \overline{st} . Let l_1, l_2 be the line segments on these lines with end points on \overline{uv} and \overline{st} . Without loss of generality, we assume $|l_1|_\infty \leq |l_2|_\infty$. Then we have $|l_1|_\infty \leq |l_0|_\infty \leq |l_2|_\infty$. (See Figure 5 for an example).




Figure 5: Five diagonal lines (black dotted lines), $\Delta_x, \Delta_s, \Delta_t, \Delta_u, \Delta_v$, and three line segments (blue solid line segments), l_1, l_0, l_2 .

Note that one of the end points of l_1 is in the set $\{s, t, u, v\}$, which is a subset of vertices in $V(I_M) \cup V(I_N)$. Let that vertex be y . Similarly one of the end points of l_2 is a vertex, which we take as z . We have $|l_1|_\infty \in D(y)$ and $|l_2|_\infty \in D(z)$ and $d_1 = |l_1|_\infty \leq d = |l_0|_\infty \leq d_2 = |l_2|_\infty$ for $y, z \in V(I_M) \cup V(I_N)$. This completes the first part of the claim.

(ii) For the second part, it is clear that if x and x' are on two parallel edges, then one has $|l_0|_\infty = |l_1|_\infty = |l_2|_\infty$, proving the claim. \square

Proposition 42. *Given two interval modules M and N , and $d \geq 0$. If there exists an intersection point $x \in B(I_M) \cap B(I_{N \rightarrow d})$ and two parallel edges $e_1 \in E(I_M)$ and $e_2 \in E(I_{N \rightarrow d})$ both containing x , then $d \in S$.*

Proof. Let $\nu : \bar{\mathbb{R}}^2 \rightarrow \bar{\mathbb{R}}^2$ be the shift function defined as $\nu(x) = x + \vec{d}$. Then

$I_N = \nu(I_{N \rightarrow d})$. Let $x' = \nu(x) = x + \vec{d}$ and $e'_2 = \nu(e_2)$. Then e'_2 and e_1 are two parallel edges containing x' and x in $B(I_N)$ and $B(I_M)$ respectively. We know that $e'_2 \subseteq L$ for some $L = L(I_N)$ or $U(I_N)$. Then we have $x' = \pi_L(x)$ with $\text{dl}(x, L) := d_\infty(x, x') = d$. By Proposition 41(ii), we have $d \in S$. \square

From the above proposition, we get the following corollary.

Corollary 43. *Let M and N be two interval modules and $d \notin S$. Then, for all intersection points $x \in B(I_M) \cap B(I_{N \rightarrow d})$, any two edges containing x in $B(I_M)$ and $B(I_{N \rightarrow d})$ are perpendicular, and these intersection points are in the interior of the edges containing them.*

B Missing proof in section 4

Proposition 32 and its proof.

For a nice function f , $f(x) = \sum_{y \leq x} \Delta f(y)$.

Proof. For a nice function f , we extend Δf to be a function $\overline{\Delta f}$ defined on $\text{Pow}(\mathbb{R}^n)$ as $\overline{\Delta f}(U) = \sum_{x \in U} \Delta f(x)$ for any $U \subseteq \mathbb{R}^n$. Note that $\overline{\Delta f}(\emptyset) = 0$ and $\overline{\Delta f}(\{x\}) = \Delta f(x)$. First, we observe the following property of the function $\overline{\Delta f}$:

$$\overline{\Delta f}(U_1 \cup U_2) = \overline{\Delta f}(U_1) + \overline{\Delta f}(U_2) - \overline{\Delta f}(U_1 \cap U_2) \quad (\star)$$

For any $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, define $R_x = \{y : y \leq x\} \subseteq \mathbb{R}^n$ and $R_x^i = R_x \setminus \{y : y_i = x_i\} = \{y : y \leq x, y_i \neq x_i\}$. For any $k = 0, \dots, n$ and $s \in \binom{[n]}{k}$, let $R_x^s = \bigcap_{i \in s} R_x^i = \{y : y \leq x, y_i \neq x_i, \forall i \in s\}$. We prove the proposition by induction on x .

Assume it is true for any $y < x$, that is $\forall y < x, f(y) = \sum_{z \leq y} \Delta f(z) = \overline{\Delta f}(R_y)$. Since $R_x = \{x\} \amalg (R_x \setminus \{x\}) = \{x\} \amalg \bigcup_i R_x^i$, by the property (\star) , we have $\sum_{y \leq x} \Delta f(y) = \overline{\Delta f}(R_x) = \overline{\Delta f}(\{x\} \amalg \bigcup_i R_x^i) = \Delta f(x) + \overline{\Delta f}(\bigcup_i R_x^i)$. By the inclusion-exclusion principle, we have

$$\begin{aligned} \overline{\Delta f}\left(\bigcup_i R_x^i\right) &= \sum_i \overline{\Delta f}(R_x^i) - \sum_{ij} \overline{\Delta f}(R_x^{\{i,j\}}) + \dots \\ &= (-1) \cdot \sum_{k=1}^n (-1)^k \sum_{s \in \binom{[n]}{k}} \overline{\Delta f}(R_x^s) \end{aligned}$$

Note that by inductive hypothesis, for any $s \in \binom{[n]}{k}$, $\lim_{\epsilon \rightarrow 0^+} f(x - \epsilon \cdot \sum_{i \in s} e_i) = \lim_{\epsilon \rightarrow 0^+} \overline{\Delta f}(R_{(x - \epsilon \cdot \sum_{i \in s} e_i)}) = \overline{\Delta f}(\bigcup_{\epsilon > 0} R_{(x - \epsilon \cdot \sum_{i \in s} e_i)}) = \overline{\Delta f}(R_x^s)$. Therefore, we have $\overline{\Delta f}(\bigcup_i R_x^i) = (-1) \cdot \sum_{k=1}^n (-1)^k \cdot \sum_{s \in \binom{[n]}{k}} \lim_{\epsilon \rightarrow 0^+} f(x - \epsilon \cdot \sum_{i \in s} e_i)$. By definition of $\Delta f(x)$, we have $f(x) = \Delta f(x) + \overline{\Delta f}(\bigcup_i R_x^i) = \overline{\Delta f}(R_x) = \sum_{y \leq x} \Delta f(y)$. \square